



TGP/8/1 Draft 3

ORIGINAL: English only

DATE: March 27, 2006

**INTERNATIONAL UNION FOR THE PROTECTION OF NEW VARIETIES OF PLANTS**  
GENEVA

**DRAFT**

Associated Document  
to the  
General Introduction to the Examination  
of Distinctness, Uniformity and Stability and the  
Development of Harmonized Descriptions of New Varieties of Plants (document TG/1/3)

**DOCUMENT TGP/8**

**[USE OF STATISTICAL PROCEDURES IN DISTINCTNESS, UNIFORMITY AND  
STABILITY TESTING]/[TRIAL DESIGN AND TECHNIQUES USED IN THE  
EXAMINATION OF DISTINCTNESS, UNIFORMITY AND STABILITY]**

*Document prepared by the Office of the Union*

*to be considered by the Technical Committee at its forty-second session  
to be held in Geneva, from April 3 to 5, 2006*

**TABLE OF CONTENTS**

INTRODUCTION .....	4
PART I: DUS TRIAL DESIGN AND DATA ANALYSIS.....	5
1. Reason for using statistics .....	5
2. Trial Design.....	6
2.1 Experimental design practices .....	6
2.1.1 Introduction.....	6
2.1.2 Trials, experimental units and test of hypotheses .....	7
2.1.3 Pair-wise comparisons of some varieties .....	8
2.1.4 Completely randomized design and randomized complete block design.....	9
2.1.5 Randomized incomplete block designs .....	12
2.1.6 Trial elements .....	13
2.1.7 Sample size .....	15
2.1.8 Analyses over years or cycles .....	16
2.2 Validation of data and assumptions .....	17
2.2.1 Introduction.....	17
2.2.2 Check on data quality (before doing analyses) .....	17
2.2.3 Assumptions.....	19
2.2.3.1 Introduction .....	19
2.2.3.2 Independent observations .....	20
2.2.3.3 Variance homogeneity .....	20
2.2.3.4 Normal distributed observations .....	21
2.2.3.5 Additivity of block and variety effects .....	21
2.2.4 Validation.....	22
2.2.4.1 Introduction .....	22
2.2.4.2 Looking through the data .....	23
2.2.4.3 Using Figures.....	23
2.3 Types of characteristics and their scale levels.....	27
2.3.1 Introduction.....	27
2.3.2 Different levels to look at a characteristic.....	28
2.3.2.1 Understanding the need for process levels:.....	29
2.3.3 Types of expression of characteristics .....	30
2.3.4 Types of scales of data .....	30
2.3.4.1 Quantitatively scaled data (metric or ordinal scaled data) .....	30
2.3.4.1.1 Ratio scale.....	31
2.3.4.1.2 Interval scale .....	32
2.3.4.2 Qualitatively scaled data .....	33
2.3.4.2.1 Ordinal scale .....	33
2.3.4.2.2 Nominal scale.....	33
2.3.5 Scale levels for variety description .....	35
2.3.6 Relation between types of expression of characteristics and scale levels of data .....	35
2.3.7 Relation between method of observation of characteristics, scale levels of data and recommended statistical procedures .....	36
PART II: TECHNIQUES USED IN DUS EXAMINATION .....	41
1. Methods for Assessing Uniformity on the Basis of Off-Types.....	41
1.1 Fixed Population Standard.....	41
1.1.1 Introduction.....	41
1.1.2 Recommendations on the fixed population standard method of assessing uniformity by the number of off-types.....	41
1.1.3 Errors in testing for off-types.....	41
1.1.4 Examples.....	43
1.1.5 Introduction to the tables and figures.....	46
1.1.6 Detailed description of the method for one single test.....	48
1.1.7 More than one single test (year).....	49
1.1.8 Detailed description of the methods for more than one single test .....	49
1.1.8.1 Combined Test.....	49
1.1.8.2 Two-stage Test .....	49
1.1.8.3 Sequential tests .....	50
1.1.9 Note on type I and type II errors.....	51
1.1.10 Definition of statistical terms and symbols.....	51
3.1.11 Tables and figures .....	53

2.	LSD .....	74
3.	The Combined Over-Years Criterion for Distinctness and Uniformity .....	74
3.1	The Combined Over-Years Distinctness Criterion (COYD) .....	74
3.1.1	Summary .....	74
3.1.2	Introduction .....	75
3.1.3	The COYD Method .....	75
3.1.4	Use of COYD .....	76
3.1.5	Adapting COYD to special circumstances .....	77
3.1.5.1	Differences between years in the range of expression of a characteristic .....	77
3.1.5.2	Small numbers of varieties in trials: Long-Term COYD .....	77
3.1.5.3	Marked year-to-year changes in an individual variety's characteristic .....	78
3.1.6	Implementing COYD .....	78
3.1.7	References .....	78
3.1.8	COYD statistical methods .....	81
3.1.8.1	Analysis of variance .....	81
3.1.8.2	Modified joint regression analysis (MJRA) .....	81
3.1.8.3	Comparison of COYD with other criteria .....	82
3.1.9	COYD software .....	82
3.2	The Combined-Over-Years Uniformity Criterion (COYU) .....	89
3.2.1	Summary .....	89
3.2.2	Introduction .....	90
3.2.3	The COYU Criterion .....	90
3.2.4	Recommendations on COYU .....	91
3.2.5	Mathematical details .....	91
3.2.6	Early decisions for a three-year test .....	94
3.2.7	Example of COYU calculations .....	94
3.2.8	Implementing COYU .....	96
3.2.9	COYU Software .....	97
3.2.9.1	DUST Computer program .....	97
3.3	Standard probability levels used for COYD and COYU .....	100
4.	Parent Formula of Hybrid Varieties .....	105
4.1	Introduction .....	105
4.2	Requirements of the method .....	105
4.3	Assessing the originality of a new parent line .....	105
4.4	Verification of the formula .....	106
4.5	Uniformity and stability of parent lines .....	107
4.6	Description of the hybrid .....	107
5.	The GAIA methodology .....	108
5.1.	Weighting of characteristics .....	108
5.2.	Determining "Distinctness Plus" .....	109
5.3.	Computing GAIA phenotypic distance .....	109
5.4.	The GAIA software .....	110
5.5.	Using the GAIA methodology .....	113
5.6.	Example with qualitative, electrophoretic and quantitative characteristics ( <i>Zea mays</i> data) .....	114
5.7	GAIA screen copy. ....	120
5.8	Final remark .....	121
5.9	References .....	121
6.	Examining DUS in bulk samples .....	122
6.1	Introduction and abstract .....	122
6.2	Distinctness .....	122
6.3	Uniformity .....	124
6.3.1	Bulking within plot .....	124
6.3.2	Bulking across plots .....	125
6.3.3	Taking just one bulk sample per plot .....	125

## **INTRODUCTION**

The purpose of this document is to provide guidance on trial design and data analysis, and to provide information on certain techniques used for the examination of DUS. This document is structured as follows:

**PART I: DUS TRIAL DESIGN AND DATA ANALYSIS:** this part of the document provides guidance on trial design, data validation, and assumptions to be fulfilled for statistical analysis.

**PART II: TECHNIQUES USED IN DUS EXAMINATION:** this part of the document provides details on certain techniques referred to in TGP/9 Examining Distinctness, and TGP/10 Examining Uniformity.

An overview of the parts of the process of examining distinctness in which trial design and techniques covered in this document are relevant is provided in [the schematic overview of the process of examining distinctness provided in document TGP/9 “Examining Distinctness”, Section 1[*cross ref.*]].

## **PART I: DUS TRIAL DESIGN AND DATA ANALYSIS**

### **1. Reason for using statistics**

To be developed by the TWC

**[TWC chairperson: to consider whether this section is necessary]**

## 2. Trial Design

[TWC Chairperson: add trial design considerations when the trials are not for statistical analysis]

### 2.1 Experimental design practices

#### 2.1.1 Introduction

2.1.1.1 Document TGP/9 “Examining Distinctness” Section 3.6.1 Type of trial layout Growing Trial explains as follows:

“The organization of the trial layout is, in the first instance, determined by whether the trial will have replicated plots and whether it will be randomized, or whether it will be organized such that similar varieties are kept together in order to facilitate side-by-side visual comparison in the growing trial. ... Information concerning replicated and randomized trial designs is provided in TGP/8 [cross ref].”

2.1.1.2 Document TGP/9 “Examining Distinctness, Section 5.3 further explains that:

“The following table summarizes the common approaches to assessing distinctness based on the growing trial, taking into account the method of propagation, type of expression of the characteristic, method of observation and the type of record.

Method of propagation of the variety	Type of expression of characteristic		
	QL	PQ	QN
Vegetatively propagated, self-pollinated	<i>Notes (VG)</i>	<i>Notes (VG)</i> <i>Side-by-side (VG)</i>	<i>Notes (VG/MG/MS)</i> <i>Side-by-side (VG)</i> <i>Statistics (MG/MS)</i>
Cross-pollinated	<i>Notes (VG)</i> <i>Statistics (VS*)</i>	<i>Notes (VG)</i> <i>Side-by-side (VG)</i> <i>Statistics (VS*)</i>	<i>Statistics ([MG]/MS/VS)</i> <i>Side-by-side (VG)</i> <i>Notes (VG/MG/MS)</i>
Hybrids	<i>Notes (VG)</i> <i>Statistics (VS*)</i>	<i>Notes (VG)</i> <i>Side-by-side (VG)</i> <i>Statistics (VS*)</i>	**

\* records of individual plants only necessary if segregation is to be recorded

\*\* to be considered according to the type of hybrid (see section 4.3.3)”

2.1.1.3 DUS trials are experiments for the comparison of varieties and the observation of characteristics. This section addresses a number of issues concerning the basics of statistics and experimental design for the examination of distinctness, uniformity and stability. Some of the issues, e.g. variance components and sample size, require that the characteristics are

continuous, quantitative characteristics (see Part I: Section 2.3 [*cross ref.*] “Type of characteristics and their scale levels”).

2.1.1.4 The UPOV Test Guidelines, where available, provide a set of recommendations for conducting the DUS test: quantity of plant material, duration of the DUS test, list of characteristics, number of plants to be observed, etc.. The examiner conducting the tests should understand the objective of the DUS test and have good knowledge of the growing conditions for the species and the factors that can affect the expressions of the characteristics of the variety. It is also important that the requirements of experimental assumptions should be well recognized (see Part I: Section 2.2.4 [*cross ref.*]). The same procedure and protocol should be followed in all growing cycles of the test to minimize the interaction between varieties and environments.

2.1.1.4 Document TGP/9 Section 2 “Selecting Varieties for the Growing Trial” explains how the varieties to be included in the growing trial may be selected. The total number of varieties (i.e. varieties of common knowledge and candidate varieties) included in the trial will influence the experimental design.

## 2.1.2 Trials, experimental units and test of hypotheses

2.1.2.1 A plot is the experimental unit to which the varieties are allocated. A plot contains plants from the same variety. A block is a group of plots within which the varieties can be pair-wise allocated or randomized.

2.1.2.2 To decide whether a variety is distinct and uniform statistical tests may be required. If this is the case, for each of these two tests we have to consider two hypotheses as specified in the following table:

	<i>Distinctness</i>	<i>Uniformity</i>
Null hypothesis (H0)	two varieties are not distinct for the characteristic	a variety is uniform for the characteristic
Alternative Hypothesis (H1)	two varieties are distinct	a variety is not uniform

By using a test statistic (which is a formula of the observations) a decision has to be made to accept the null hypothesis H0 and thus to reject the alternative hypothesis H1 or vice versa. The decision to reject H0 occurs if the test statistic is greater than the chosen critical value, otherwise H0 is accepted. If H0 is rejected the test is called significant. The different types of error which can be made for distinctness are shown in the following table:

		Real situation	
		H0 true two varieties are not distinct	H1 true two varieties are distinct
Decision	H0 accept two varieties are not distinct	Correct decision	type II error ( $\beta$ )
	H0 reject two varieties are distinct	type I error ( $\alpha$ )	Correct decision

The same situation for uniformity is shown in the next table:

		Real situation	
		H0 true a variety is uniform	H1 true a variety is not uniform
Decision	H0 accept a variety is uniform	Correct decision	type II error ( $\beta$ )
	H0 reject a variety is not uniform	type I error ( $\alpha$ )	Correct decision

2.1.2.3 When doing tests there are always two types of error. They are called type I error and type II error. Let's take the test of whether two varieties are distinct. The type I error is the error that arises when we decide that the varieties are distinct, when, in fact, they are not distinct. The type II error is the error that arises when we decide that the varieties are not distinct, when, in fact, they are distinct (valid for distinctness). The risk of type I error can be controlled easily by taking a self chosen size  $\alpha$  of the test, whereas the risk of type II error is more difficult to control as it depends on the size of the real difference between the varieties, the random variability  $s$ , the number of replicates and the chosen  $\alpha$ .

2.1.2.4 One of the most important requirements of experimental units is independence. That means that observations within a plot are not influenced by the circumstances in other plots. For example, if tall varieties are planted next to short ones there could be a negative influence of the tall ones to the short ones and a positive influence in the other direction. In such a case, an additional row of plants can be planted on both sides of the plot in order to avoid this dependency. Another possibility to minimize this influence is to group similar varieties together. Where grouping is used it should be done according to the principles set out in document TGP/9 Section 3[cross ref.]

2.1.2.5 When the same variety is assigned to a number of different plots, the observations in the different plots may vary. The variation between these observations will be called the 'between-plot variability'. This variability is a mixture of different sources of variation: different plots, different plants, different times of observation, different errors of measurement and so on. It is not possible to distinguish between these sources of variation. When there are observations of more than one, say  $n$ , plants per plot it is possible to compute two variance components: the "within-plot" or "plant" component and the "plot" component.

### 2.1.3 Pair-wise comparisons of some varieties

**[TWC Chairperson: to consider moving after Section 2.1.5]**

2.1.3.1 When a close comparison is needed between a pair of varieties it may be good to grow them in neighbouring plots. A similar theory to that used in split-plot designs may be used for setting up a design where the comparisons between certain pairs of varieties are to be optimized. When setting up the design, the pairs of varieties are treated as the whole plot factor and the comparison between varieties within each pair is the sub-plot factor. As each whole plot consists of only two sub-plots, the comparisons within pairs will be (much) more precise than if a randomized block design was used.




2.1.3.2 If, for example, four pairs of varieties (A-B, C-D, E-F and G-H) have to be compared very precisely, then this can be done using the following design of 12 whole plots each having 2 subplots:

Pair 1 variety A	Pair 3 variety E	Pair 4 variety H
Pair 1 variety B	Pair 3 variety F	Pair 4 variety G
Pair 3 variety F	Pair 2 variety D	Pair 1 variety A
Pair 3 variety E	Pair 2 variety C	Pair 1 variety B
Pair 4 variety G	Pair 1 variety B	Pair 2 variety C
Pair 4 variety H	Pair 1 variety A	Pair 2 variety D
Pair 2 variety D	Pair 4 variety H	Pair 3 variety E
Pair 2 variety C	Pair 4 variety G	Pair 3 variety F

In this design each column represents a replicate. Each of these is then divided into four incomplete blocks (whole plots) each consisting of two (sub)plots. The four pairs of varieties are randomized to the incomplete blocks within each replicate and the order of varieties are randomized within each incomplete block. The comparison between varieties of the same pair is made more precise at the cost of the precision of the comparison between varieties of a different pair.

2.1.4 Completely randomized design and randomized complete block design

2.1.4.1 In designing an experiment it is important to choose an area of land that is as homogeneous as possible in order to minimize the variation between plots of the same variety. Assume that we have a field where it is known that the largest variability is in the ‘north-south’ direction, e.g. as in the following figure:



Let's take an example where four varieties are to be compared with each other in an experiment where each of the varieties is assigned to 4 different plots. It is important to randomize the varieties over the plots. If varieties are arranged systematically, not all varieties would necessarily be under the same conditions (see following figure).

Variety A	Variety A	Variety A	Variety A	Variety B	Variety B	Variety B	Variety B	Higher fertility row
Variety C	Variety C	Variety C	Variety C	Variety D	Variety D	Variety D	Variety D	
								Lower fertility row

For reasons of space only the allocation of the four varieties into two rows is presented

**[TWC Chairperson: to provide a complete illustration]**

If the fertility of the soil decreases from the north to the south of the field, the plants of variety A and B have grown on more fertile plots than the other varieties. The comparison of the varieties is influenced by a difference in fertility of the plots. Differences between varieties are said to be confounded with differences in fertility.

2.1.4.2 To avoid systematic errors it is advisable to randomize varieties across the site. A complete randomization of the four varieties over the sixteen plots could have resulted in the following layout:

Variety C	Variety A	Variety A	Variety B	Variety C	Variety D	Variety B	Variety C	Higher fertility row
Variety C	Variety A	Variety D	Variety A	Variety D	Variety B	Variety D	Variety B	
								Lower fertility row

For reasons of space only the allocation of the four varieties into two rows is presented

**[TWC Chairperson: to provide a complete illustration]**

However, looking at the design we find that variety C occurs three times in the top row (with high fertility) and only once in the second row (with lower fertility). For variety D we have the opposite situation. Because we know that there is a fertility gradient, this is still not a good design, but it is better than the first systematic design.

2.1.4.3 When we know that there are certain systematic sources of variation like the fertility gradient in the paragraphs before, we may take that information into account by making so-called blocks. The blocks should be formed so that the plots within each block are as homogeneous as possible. With the assumed gradients we may choose either two blocks each consisting of one row or we may choose four blocks – two blocks in each row with four plots each. In larger trials (more plots) the latter will most often be the best, as there will also be some variation within rows even though the largest gradient is between rows.

Block I				Block II				
Variety A	Variety C	Variety D	Variety B	Variety A	Variety C	Variety D	Variety B	Higher fertility row
Variety B	Variety C	Variety A	Variety D	Variety C	Variety A	Variety D	Variety B	Lower fertility row

Block III

Block IV

For reasons of space only the allocation of the four varieties into two rows is presented

**[TWC Chairperson: to provide a complete illustration]**

An alternative way of reducing the effect of any gradient between the columns is to use plots which extend over two rows, i.e. by using long and narrow plots:

Block I				Block II				Block III				Block IV			
Var	Var	Var	Var	Var	Var	Var	Var	Var	Var	Var	Var	Var	Var	Var	Var
A	C	D	B	A	C	D	B	B	C	A	D	C	A	D	B

For reasons of space only the allocation of the four varieties into two rows is presented

**[TWC Chairperson: to provide a complete illustration]**

In both designs above the ‘north-south’ variability will not affect the comparisons between varieties.

2.1.4.4 In a randomized complete block design the number of plots per block equals the number of varieties. All varieties are present once in each block and the order of the varieties within each block is randomized. The advantage of a randomized complete block design is that the standard deviation between plots (varieties) does not contain variation due to differences between blocks. The main reason for the random allocation is that it ensures that the results are unbiased and so represent the varieties being compared. An interesting feature of the randomization is that it makes the observations from individual plots ‘behave’ as independent observations (even though they may not be so). There is usually no extra cost associated with blocking, so it is recommended to arrange the plots in blocks.

**[TWC Chairperson: Highlighted text in paragraph 2.1.4.4 to be clarified]**

2.1.4.5 Blocking is introduced here on the basis of differences in fertility. Several other systematic sources of variation could have been used as the basis for blocking. Although it is not always clear how heterogeneous the field is, and therefore it is unknown how to arrange the blocks, it is usually a good idea to create blocks for other reasons. When there are different sowing machines, different observers, different observation days, such effects are included in the residual standard deviation if they are randomly assigned to the plots. However, these effects can be eliminated from the residual standard deviation if all the plots within each block have the same sowing machine, the same observer, the same observation day, and so on.

2.1.4.6 Management may influence the choice of the form of the plots. In some crops it may be easier to handle long and narrow plots than square plots. Long narrow plots are

usually considered to be more influenced by varieties in adjacent plots than square plots. The size of the plots should be chosen in such a way that the necessary number of plants for sampling is available. For some crops it may be necessary also to have guard plants (areas) in order to avoid large competition effects. However, overly large plots require more land and will often increase the random variability between plots. Grouping of the varieties according to e.g. height may also reduce the competition between adjacent plots. If nothing is known about the fertility of the area, then layouts with compact blocks (i.e. almost square blocks) will often be most appropriate because the larger the distance between two plots the more different they will usually be. In both designs above, the blocks can be placed as shown or they could be placed 'on top of each other' [TWC Chairperson: illustration to be provided]. This will usually not change the variability between plots considerably – unless one of the layouts, forces the crop expert to use more heterogeneous soil.

### 2.1.5 Randomized incomplete block designs

2.1.5.1 If the number of varieties becomes very large (>20-40), it may be impossible to construct complete blocks that would be sufficiently homogeneous. In that case it might be advantageous to form smaller blocks, each one containing only a fraction of the total number of varieties. Such designs are called incomplete block designs. Several types of incomplete block designs can be found in the literature [TWC Chairperson: to be specified]. One of the most familiar types for variety trials is a lattice design. The generalized lattice designs (also called  $\alpha$ -designs) are very flexible and can be constructed for any number of varieties and for a large range of block sizes and number of replicates. One of the features of generalized lattice designs is that some of the incomplete blocks can be (and usually are) collected to form a whole replicate. This means that such designs will be at least as good as randomized complete block designs, since the analysis can be performed using either a lattice model or a randomized complete block model. The lattice model should be preferred if conditions are fulfilled.

2.1.5.2 Incomplete blocks need to be constructed in such a way that it is possible to compare all varieties in an efficient way. An example of an  $\alpha$ -design is shown in the following figure:

Block	Sub-block	Variety			
3	5	6	5	15	19
	4	13	8	10	20
	3	2	3	4	7
	2	12	1	18	14
	1	17	11	16	9
Block	Sub-block	Variety			
2	5	4	16	6	1
	4	18	5	10	2
	3	14	7	17	8
	2	11	19	13	3
	1	15	9	20	12
1	5	4	20	5	17
	4	2	13	1	9
	3	3	6	12	8
	2	18	7	11	15
	1	16	10	14	19

In the example above, 20 varieties are to be grown in a trial with three replicates. In the design the 5 sub-blocks of each block form a complete replicate. Thus each replicate contains all varieties whereas any pair of varieties occurs either once or not at all in the same subblock.


2.1.5.3 The incomplete block design is most suitable for trials where grouping characteristics are not available. If grouping characteristics are available then some modification may be advantageous for trials with many varieties, such as using grouping characteristics to form separate trials rather than a single trial, see document TGP/9 Section 3.6.2.1 Grouping characteristics.

## 2.1.6 Trial elements

2.1.6.1 An experimental unit in a DUS trial is a plot containing plants of a variety. The observation of certain characteristics on individual plants is used to estimate the mean and the variability of the characteristic for a variety. A plot is the smallest subdivision of the trial and the unit on which the varieties and the soil and plant condition should be focussed. Therefore the following trial elements should be arranged accordingly:

- plot size
- shape of the plots
- alignment of the plots
- barrier rows and border strips and
- protective strips

2.1.6.2 The following figure may be helpful to give some explanations of the particular trial elements.



2.1.6.3 The Test Guidelines indicate the recommended method of observation and the type of record for the assessment of distinctness. Guidance on the method of observation, visual (V) or measurement (M); and on the type of record, single record for a group of plants or parts of plants (G) or records for a number of single individual plants or parts of plants (S) is provided in TGP/9 Section 4 Type of observation [cross ref.].

2.1.6.4 The highest requirements on the planning of the trial are based on characteristics on single individual plants (MS and VS). These characteristics determine the number of single plants and therefore the size of the plot. In some cases it is necessary to have border rows and strips to minimize the inter-plot interference and other special border effects.

2.1.6.5 The plot size depends upon the sample size. Furthermore, plot size and plot shape also depend on the soil conditions and on the sowing and harvesting machinery. The shape of the plot can be defined as the ratio of plot length divided by plot width. This ratio can be important to mitigate variation within the block (e.g. caused by soil variation).

2.1.6.6 Square plots have the smallest total length of the borders (circumference). From the theoretical point of view the square shape is optimal to minimize the interference of genotypes. Grouping of the varieties can have the same effect.

2.1.6.7 Narrow and long plots are preferred from the technological point of view. The best length to width ratio lies between 5:1 and 15:1 and depends on the plot size and the number of varieties. The larger the number of varieties in a block the narrower the plots - but not so narrow that the inter-plot competition becomes a problem. When statistical analysis is intended, the aim of DUS testing is to get averages of characteristics for each variety and to judge the within-variety variability by calculating the standard deviation. The averages will be used for determining the distinctness of the varieties; the standard deviations are the basis for examination of uniformity in the case of quantitative characteristics. For qualitative characteristics the number of off-types will be determined. (See document TG/1/3 "General Introduction, section 6.4.4.1.)

2.1.6.8 For assessment of distinctness unbiased and precise estimation of averages is necessary. The bias is difficult to calculate. Nevertheless it is common to reduce the bias by suitable precautions which are the exclusion of external influences by means of protective strips on the border of the trial. Additionally, it is often necessary to exclude border rows and strips of the plot from calculations of the average and the standard deviation. The rest of the plot without border rows and strips (effective plot size) are the basis of the unbiased and precise estimates.

2.1.6.9 The plants may be arranged in different ways in the trials:

- Rows of plants: This type of arrangement is used for many self-pollinated species, such as cereals. Most characteristics are assessed in an overall observation – usually using the notes stated in the Test Guidelines. In some cases it may be necessary to remove some plants from the plot in order to record some characteristics; and in that case the size of the plot should allow the removal of plants without prejudicing the observations which must be made up to the end of the growing cycle including the assessment of uniformity (see document TGP/7, ASW 6 [*cross ref.*]).
- Ear rows: This type of arrangement is frequently used for the assessment of uniformity in self-pollinated varieties.
- Spaced plants: This type of arrangement is used in many cross-pollinated and vegetatively propagated varieties.

## 2.1.7 Sample size

2.1.7.1 The Test Guidelines will usually define the sample size of one experiment. The final precision of a test based on the observations of one experiment depends for quantitative characteristics on at least three sources of variation:

- the variation between individual plants within a plot
- the variation between the plots within a block

- the variation caused by the environment, i.e. the variation in the expression of characteristics from year to year (or from location to location)

2.1.7.2 To estimate the optimal sample size when developing new Test Guidelines it is necessary to know the standard deviations, expected differences between the varieties which should be significant, the number of varieties and the number of blocks in the trial. Additionally, the crop expert has to determine the type I ( $\alpha$ ) and type II error ( $\beta$ ). In cooperation with a statistician the crop expert can compute the optimal sample size for some characteristics and then he can determine the optimal sample size for this trial for all characteristics. Especially for the assessment of uniformity, the type II error is sometimes more important than the type I error. In some cases the type II error could be greater than 50 % and becomes unacceptable.

#### 2.1.8 Analyses over years or cycles

2.1.8.1 The comparison between varieties is mostly based on observations from two to three years or cycles. Therefore, the number of replicates and the number of plants per plot in a single trial have an effect on the variability which is used in the statistical analyses (see Part II: Sections 2.1 and 2.2 [*cross ref.*] ). Before performing these analyses the means of the variety means and (log) standard deviations per year or cycle are calculated and then the analysis is performed on these means in the two-way variety-by-year or -cycle layout. The residual variation in these analyses is the variety-by-year or -cycle interaction.

2.1.8.2 The precision of the variety means in one experiment, when used for COY-D for example, is only used indirectly, because the standard deviation in that analysis is based on the interaction between the varieties and the years or cycles. If the differences between the varieties over the years or cycles are very large, the precision of the means per experiment are less important.



## 2.2 Validation of data and assumptions

### 2.2.1 Introduction

[TWC Chairperson: to be reviewed to clarify that aspects of this section may be relevant when statistics are not used]

In Part I: Section 2.1, [*cross ref.*] “Experimental Design Practices”, aspects of designing the experiments in which the data are recorded are discussed. In Part I: Section 2.3 [*cross ref.*], “Types of Characteristics and Their Scale Levels”, it is shown that the choice of which statistical methods to use depends on the type of characteristic, its scale level, and whether distinctness or uniformity is considered. The statistical methods are based on some theory and in order to ensure that the results can be trusted the assumptions behind the theory have to be met - at least approximately. The purpose of this section is to describe the assumptions behind the most common statistical methods used in DUS testing and to show how these assumptions may be validated. It is important to note that the COYD and COYU methods for quantitative characteristics are based on variety means per year for COYD, and variety means of the (logarithm of the) between-plants standard deviation per year for COYU. Some methods for checking the data are described in Part I: Section 2.2.2 [*cross ref.*] “Check on Data Quality” below. In Part I: Section 2.2.3, [*cross ref.*] “Assumptions”, the assumptions underlying the analysis of variance methods are given and in Part I: Section 2.2.4, [*cross ref.*] “Validation”, some methods for evaluating these assumptions are given. The assumptions and methods of validation are described here for the analyses of single experiments (randomized blocks). However, the principles are the same when analyzing data from several experiments over years. Instead of plot means, the analyses are then carried out on variety means per year (and blocks then become equivalent to years). The methods described here are intended for quantitative characteristics, but some of the methods may also be used for checking qualitative characteristics on the ordinal scale. Throughout this section data of ‘Leaf: Length’ (in mm) are used from an experiment laid out in 3 blocks of 26 plots with 20 plants per plot. Within each block, 26 different oilseed rape varieties were randomly assigned to each plot.

### 2.2.2 Check on data quality (before doing analyses)

2.2.2.1 In order to avoid mistakes in the interpretation of the results the data should always be inspected so that the data are logically consistent and not in conflict with prior information about the ranges likely to arise for the various characteristics. This inspection can be done manually (usually visually) or automatically.

2.2.2.2 Table 1 shows an extract of some recordings for 10 plants from a plot of field peas. For ‘Seed: shape’ the notes are visually scored on a scale with values 1, 2, 3, 4, 5 or 6. For ‘Stem: length’ the measurements are in cm and from past experience it is known that the length in most cases will be between 40 and 80 cm. The ‘Stipule: length’ is measured in mm and will in most cases be between 50 and 90 mm. The table shows 3 types of mistakes which occasionally occur when making manual recordings: for plant 4, ‘Seed: shape’ the recorded value, 7, is not among the allowed notes and must, therefore, be due to a mistake. It might be caused by a misreading a hand-written “1”. The ‘Stem: length’ of plant 6 is outside the expected range and could be caused by changing the order of the figures, so 96 has been keyed instead of 69. The ‘Stipule: length’ of 668 mm is clearly wrong. It might be caused by

accidentally repeating the figure 6 twice. In all cases a careful examination needs to be carried out in order to find out what the correct values should be.

Table 1 Extract of recording sheet for field peas

Plant no	Seed: shape (UPOV 1)	Stem: length (UPOV 12)	Stipule: length (UPOV 31)
1	1	43	80
2	2	53	79
3	1	50	72
4	7	43	668
5	2	69	72
6	1	96	72
7	1	51	70
8	2	64	63
9	1	44	62
10	2	49	62

2.2.2.3 Examination of frequency distributions of the characteristics to look for small groups of discrepant observations.

2.2.2.4 Examination of scatter plots of pairs of characteristics likely to be highly related. This may often detect discrepant observations very efficiently.

2.2.2.5 Other types of [plot] may also be used to validate the quality of the data. A so-called Box-plot is an efficient way to get an overview of the data. In a Box-plot a box is drawn for each group (plot or variety). In Figure 1, all 60 Leaf Lengths of each of the 26 varieties are taken together. (If there are large block differences a better Box-plot can be produced by taking the differences with respect to the plot mean). The box shows the range for the largest part of the individual observations (usually 75%). A horizontal line through the box and a symbol indicates the median and mean, respectively. At each end of the box, vertical lines are drawn to indicate the range of possible observations outside the box, but within a reasonable distance (usually 1.5 times the height of the box). Finally, more extreme observations are shown individually. In Figure 1, it is seen that one observation of variety 13 is clearly much larger than the remaining observations of that variety. Also it is seen that variety 16 has large leaf lengths and that about 4 observations are relatively far from the mean. Among other things that can be seen from the figure are the variability and the symmetry of the distribution. So it can be seen that the variability of variety 15 is relatively large and that the distribution is slightly skewed for this variety (as the mean and median are relatively far apart).




Figure 1. Box-plot for Leaf Length of 26 varieties of oil seed rape

2.2.2.6 When discrepant observations are found, it is important to try to find out why the observations are deviating. In some cases it may be possible to go back to the field and to check if the plant or plot is damaged by external factors (e.g. rabbits) or a measurement mistake has occurred. In the latter case a correction is possible. In other cases, it may be necessary to look in previous notes (or on other measurements from the same plant/plot) in order to find the reason for the discrepant observation. Generally observations should only be removed when there are good reasons.

## 2.2.3 Assumptions

### 2.2.3.1 Introduction

2.2.3.1.1 First of all it is very important to design experiments in a proper way. The most important assumptions of analysis of variance methods are:

- independent observations
- variance homogeneity
- additivity of block and variety effects for a randomized block design and additivity of year and variety effects for COYD
- normally distributed observations (residuals)

2.2.3.1.2 In addition, one could state that there should be no mistakes in the data. However, most mistakes (at least the largest) will usually also mean that the observations are not normally distributed and that they have different variances.

2.2.3.1.3 The assumptions mentioned here are most important when the statistical methods are used to test hypotheses. When statistical methods are used only to estimate effects (means), the assumptions are less important and the assumption of normal distributed observations is not necessary.

#### 2.2.3.2 Independent observations

This is a very important assumption. It means that no records may depend on other records in the same analysis (dependence between observations may be built into the model, but has not been built into COYD and COYU or the other methods included in TGP/8). Dependency may be caused by e.g. competition between neighboring plots, lack of randomisation or improper randomisation. More details on ensuring independence of observations may be found in Part I: Section 2.1 [*cross ref.*] “Experimental Design Practices.”

#### 2.2.3.3 Variance homogeneity

Variance homogeneity means that the variance of all observations should be identical apart from random variation. Typical deviations from the assumption of variance homogeneity fall most often into one of the following two groups:

The variance depends on the mean, e.g. the larger the mean value the larger the standard deviation is. In this case the data may often be transformed such that the variances on the transformed scale may be approximately homogeneous. Some typical transformations of characteristics are: the logarithmic transformation (where the standard deviation is approximately proportional to the mean), the square-root transformation (where the variance is approximately proportional to the mean, e.g. counts), and the angular transformation (where the variance is low at both ends of the scale and higher in between, typical for percentages).

The variance depends on for example, variety, year or block. If the variances depend on such variables in a way that is not connected to the mean value, it is not possible to obtain variance homogeneity by transformation. In such cases it might be necessary either to use more sophisticated statistical methods that can take unequal variances into account or to exclude the group of observations with deviant variances (if only a few observations have deviant variances). To illustrate the seriousness of variance heterogeneity: imagine a trial with 10 varieties where varieties A, B, C, D, E, F, G and H each have a variance of 5, whereas varieties I and J each have a variance of 10. The real probability of detecting differences between these varieties when, in fact, they have the same mean is shown in Table 2. In Table 2, the variety comparisons are based on the pooled variance as is normal in traditional ANOVA. If they are compared using the 1% level of significance, the probability that the two varieties with a variance of 10 become significantly different from each other is almost 5 times larger (4.6%) than it should be. On the other hand, the probability of significant differences between two varieties with a variance of 5 decreases to 0.5%, when it should be 1%. This means that it becomes too difficult to detect differences between two varieties with small variances and too easy to detect differences between varieties with large variances.

Table 2. Real probability of significant difference between two identical varieties in the case where variance homogeneity is assumed but not fulfilled (varieties A to H have a variance of 5 and varieties I and J have a variance of 10.)

Comparisons, variety names	Formal test of significance level	
	1%	5%
A and B	0.5%	3.2%
A and I	2.1%	8.0%
I and J	4.6%	12.9%

#### 2.2.3.4 Normal distributed observations

The residuals should be approximately normally distributed. The ideal normal distribution means that the distribution of the data is symmetric around the mean value and with the characteristic bell-shaped form (see Figure 2). If the residuals are not approximately normally distributed,

the actual level of significance may deviate from the nominal level. The deviation may be in both directions depending on the way the actual distribution of the residuals deviates from the normal distribution.

However, deviation from normality is usually not as serious as deviations from the previous two assumptions.




Figure 2. Histogram for normal distributed data with the ideal normal distribution shown as a curve

#### 2.2.3.5 Additivity of block and variety effects

2.2.3.5.1 The effects of blocks and varieties are assumed to be additive because the error term is the sum of random variation and the interaction between block and variety. This means that the effect of a given variety is the same in all blocks. This is demonstrated in Table 3 where plot means of artificial data (of Leaf Length in mm) are given for two small experiments with three blocks and four varieties. In experiment I, the effects of blocks and varieties are additive because the differences between any two varieties are the same in all blocks, e.g. the differences between variety A and B are 4 mm in all three blocks. In experiment II, the effects are not additive, e.g. the differences between variety A and B are 2, 2 and 8 mm in the three blocks.

Table 3. Artificial plot means of Leaf Length in mm from two experiments showing additive block and variety effects (left) and non-additive block and variety effects (right)

Experiment I				Experiment II			
Variety	Block			Variety	Block		
	1	2	3		1	2	3
A	240	242	239	A	240	242	239
B	244	246	243	B	242	244	247
C	245	247	244	C	246	244	243
D	241	243	240	D	241	242	241




Figure 3. Artificial plot means from two experiments showing additive block and variety effects (left) and non-additive block and variety effects (right) using same data as in table 2

2.2.3.5.2 In Figure 3 the same data are presented graphically. Plotting the means versus block numbers and joining the observations from the same varieties by straight lines produces the graphs. Plotting the means versus variety names and joining the observations from the same blocks could also have been used (and may be preferred especially if many varieties are to be shown in the same figure). The assumption on additivity is fulfilled if the lines for the varieties are parallel (apart from random variation). As there is just a single data value for each variety in each block, it is not possible to separate interaction effects and random variation. So in practice the situation is not as nice and clear as here because the effects may be masked by random variation.

## 2.2.4 Validation

**[TWC Chairperson: To review the content of this section for overlap with section 2.2.2]**

### 2.2.4.1 Introduction

2.2.4.1.1 The purpose of validation is partly to check that the data are without mistakes and that the assumptions underlying the statistical analyses are fulfilled.

2.2.4.1.2 There are different methods to use when validating the data. Some of these are:

- look through the data
- produce plots to verify the assumptions

- make formal statistical tests for the different types of assumptions. In the literature several methods to test for outliers, variance homogeneity, additivity and normality may be found. Such methods will not be mentioned here partly because many of these depend on assumptions that do not affect the validity of COYD and COYU seriously and partly because the power of such methods depends heavily on the sample size (this means that serious lack of assumptions may remain undetected in small datasets, whereas small and unimportant deviations may become statistically significant in large datasets)

#### 2.2.4.2 Looking through the data

In practice this method is only applicable when a few observations have to be checked. For large datasets this method takes too much time, is tedious and the risk of overlooking suspicious data increases as one goes through the data. In addition, it is very difficult to judge the distribution of the data and to judge the degree of variance homogeneity when using this method.

#### 2.2.4.3 Using Figures

2.2.4.3.1 Different kinds of figures can be prepared which are useful for the different aspects to be validated. Many of these consist of plotting the residuals in different ways. (The residuals are the differences between the observed values and the values predicted by the statistical model).

2.2.4.3.2 The plot of the residuals versus the predicted values may be used to judge the dependence of the variance on the mean. If there is no dependence, then the observations should fall approximately (without systematic deviation) in a horizontal band symmetric around zero (Figure 4). In cases where the variance increases with the mean, the observations will fall approximately in a funnel with the narrow end pointing to the left. Outlying observations, which may be mistakes, will be shown in such a figure as observations that clearly have escaped from the horizontal band formed by most other observations. In the example used in figure 4, no observations seem to be outliers (the value at the one bottom left corner where the residual is about -40 mm may at first glance look so, but several observations have positive values of the same numerical size). Here it is important to note that an outlier is not necessarily a mistake and also that a mistake will not necessarily show up as an outlier.




Figure 4. Plot of residuals versus plot predicted values for Leaf Length in 26 oil seed rape varieties in 3 blocks

2.2.4.3.3 The residuals can also be used to form a histogram, like Figure 2, from which the assumption about the distribution can be judged.

2.2.4.3.4 The range (maximum value minus minimum value) or standard deviation for each plot may be plotted versus some other variables such as the plot means, variety number or plot number. Such figures (Figure 5) may be useful to find varieties with an extremely large variation (all plots of the variety with a large value) or plots where the variation is extremely large (maybe caused by a single plant). It is clearly seen that the range for one of variety 13's plots is much higher than in the other two plots. Also the range in one of variety 3's plots seems to be relatively large.






Figure 5. Differences between minimum and maximum of 20 leaf lengths for 3 plots versus oil seed rape variety number

2.2.4.3.5 A figure with the plot means (or variety adjusted means) versus the plot number can be used to find out whether the characteristic depends on the location in the field (Figure 6). This, of course, requires that the plots are numbered such that the numbers indicate the relative location. In the example shown in Figure 6, there is a clear trend showing that the leaf length decreases slightly with plot number. However most of the trend over the area used for the trial will - in this case - be explained by differences between blocks (plot 1-26 is block 1, plot 27-52 is block 2 and plot 53-78 is block 3).




Figure 6. Plot means of 20 Leaf Lengths versus plot numbers

2.2.4.3.6 The plot means can also be used to form a figure where the additivity of block and variety effects can be visually checked at (see Figure 3).

2.2.4.3.7 Normal Probability Plots (Figure 7). This type of graph is used to evaluate to what extent the distribution of the variable follows the normal distribution. The selected variable will be plotted in a scatter plot against the values “expected from the normal distribution.” The standard normal probability plot is constructed as follows. First, the residuals (deviations from the predictions) are rank ordered. From these ranks the program computes the expected values from the normal distribution, hereafter called z-values. These z-values are plotted on the X-axis in the plot. If the observed residuals (plotted on the Y-axis) are normally distributed, then all values should fall onto a straight line. If the residuals are not normally distributed, then they will deviate from the line. Outliers may also become evident in this plot. If there is a general lack of fit, and the data seem to form a clear pattern (e.g. an S shape) around the line, then the variable may have to be transformed in some way.




Figure 7. Normal probability plot for the residuals of Leaf Length in 26 oil seed rape varieties in 3 blocks

## 2.3 Types of characteristics and their scale levels

[TWC Chairperson: TWC to consider whether this section helps crop experts to better understand the statistical basis for the examination of characteristics]

### 2.3.1 Introduction

2.3.1.1 The General Introduction makes the following recommendations with regard to the use of statistical methods in the assessment of distinctness:

“5.5 Interpretation of Observations for the Assessment of Distinctness with the Application of Statistical Methods

#### 5.5.1 General

5.5.1.1 For measured characteristics as well as for visually **assessed<sup>[\*]</sup>** characteristics statistical methods can be applied. Appropriate methods have to be chosen for the interpretation of observations. The data structure and the type of scale from a statistical point of view (nominal, ordinal, interval or ratio) is decisive for the choice of appropriate methods. The data structure depends on the method of **assessment<sup>[\*]</sup>** (visual **assessment<sup>[\*]</sup>** or measurements, observation of plots or single plants) which is influenced by the type of characteristic, the features of propagation of the variety, the experimental design and other factors. DUS examiners should be aware of certain basic rules of statistics and especially the fact that their use is linked to mathematical assumptions and the use of experimental design practices, such as randomization. Therefore, those assumptions should be verified before applying statistical methods.

Some statistical methods are quite robust, however, and can be used, with some caution, even if some assumptions are not fully met.

5.5.1.2 Document TGP/8, “Use of Statistical Procedures in DUS Testing,” provides guidance on some appropriate statistical procedures for DUS assessment and includes keys for the choice of methods in relation to the data structure.

[...]

#### 5.5.2 Visually Assessed<sup>[\*]</sup> Characteristics

Non-parametric statistics may be used when visually assessed<sup>[\*]</sup> characteristics have been recorded on a scale that does not fulfill the assumptions of the usual parametric statistics. The calculation of the mean value, for example, is only permitted if the Notes are taken on a graded scale which shows equal intervals throughout the scale. In the case of non-parametric procedures, the use of a scale that has been established on the basis of example varieties representative of the different states of the characteristics is recommended. The same variety should then always receive about the same Note and thereby facilitate the interpretation of data. More details on the handling of visually assessed<sup>[\*]</sup> characteristics are given in document TGP/9, “Examining Distinctness.”

([\*] the term “observed” would be more consistent with the use of the terms “observed” and “assessed” in TGP/9)

2.3.1.2 For the revision of UPOV Test Guidelines or for establishing new ones, and in order to understand the relations between the different steps of work of the crop experts during the DUS test, it is necessary to have an answer to the following questions:

1. What is a characteristic?
2. What is a process level?
3. What is a scale level of a characteristic?
4. What is the influence of the scale level on the :
  - planning of a trial,
  - recording of data,
  - determination of distinctness and uniformity and
  - description of varieties.

#### 2.3.2 Different levels to look at a characteristic

Characteristics can be considered in different levels of process (Table 1). The characteristics as expressed in the trial (type of expression) are considered as process level 1. The data taken from the trial for the assessment of distinctness, uniformity and stability are defined as process level 2. These data are transformed into states of expression for the purpose of variety description. The variety description is process level 3.

Table 1: Definition of different process levels to consider characteristics

Process level	Description of the process level
1	characteristics as expressed in trial
2	data for evaluation of characteristics
3	variety description

From the statistical point of view the information level decreases from process level 1 to 3. Statistical analysis is only applied in level 2.

Sometimes for crop experts it seems that there is no need to distinguish between different process levels. The process level 1, 2 and 3 could be identical. However, in general, this is not the case.

#### 2.3.2.1 Understanding the need for process levels:

2.3.2.1.1 The crop expert may know from UPOV Test Guidelines or his own experience that, for example, ‘Length of plant’ is a good characteristic for the examination of DUS. There are varieties which have longer plants than other varieties. Another characteristic could be ‘Variegation of leaf blade’. For some varieties, variegation is present and for others not. The crop expert has now two characteristics and he knows that ‘Plant length’ is a quantitative characteristic and ‘Variegation of leaf blade’ is a qualitative characteristic (definitions: see Part I: Section 2.3.3.2 [*cross ref.*] below). This stage of work can be described as **process level 1**.

2.3.2.1.2 The crop expert then has to plan the trial and to decide on the type of observation for the characteristics. For characteristic ‘Variegation of leaf blade’, the decision is clear. There are two possible expressions: ‘present’ or ‘absent’. The decision for characteristic ‘Plant length’ is not specific and depends on expected differences between the varieties and on the variation within the varieties. In many cases, the crop expert will decide to measure a number of plants (in cm) and to use special statistical procedures to examine distinctness and uniformity. But it could also be possible to assess the characteristic ‘Plant length’ visually by using expressions like ‘short’, ‘medium’ and ‘long’, if differences between varieties are large enough (for distinctness) and the variation within varieties is very small or absent in this characteristic. The continuous variation of a characteristic is assigned to appropriate states of expression which are recorded by notes (see TGP/9, Section 4)[*cross. ref.*]. The crucial element in this stage of work is the recording of data for further evaluations. It is described as **process level 2**.

2.3.2.1.3 At the end of the DUS test, the crop expert has to establish a description of the varieties using notes from 1 to 9 or parts of them. This phase can be described as **process level 3**. For ‘Variegation of leaf blade’ the crop expert can take the same states of expression (notes) he recorded in process level 2 and the three process levels appear to be the same. In cases where the crop expert decided to assess ‘Plant length’ visually, he can take the same states of expression (notes) he recorded in process level 2 and there is no obvious difference between process level 2 and 3. If the characteristic ‘Plant length’ is measured in cm, it is necessary to assign intervals of measurements to states of expressions like ‘short’, ‘medium’ and ‘long’ to establish a variety description. In this case, for statistical procedures, it is important to be clearly aware of the relevant level and to understand the differences between characteristics as expressed in the trial, data for evaluation of characteristics and the variety

description. This is absolutely necessary for choosing the most appropriate statistical procedures in cooperation with statisticians or by the crop expert.

### 2.3.3 Types of expression of characteristics

2.3.3.1 Characteristics can be classified according to their types of expression. The consideration of the type of expression of characteristics corresponds to process level 1. The following types of expression of characteristics are defined in the General Introduction to the Examination of Distinctness, Uniformity and Stability and the Development of Harmonized Descriptions of New Varieties of Plants, (document TG/1/3, the “General Introduction”, Chapter 4.4):

2.3.3.2 Qualitative characteristics” are those that are expressed in discontinuous states (e.g. sex of plant: dioecious female (1), dioecious male (2), monoecious unisexual (3), monoecious hermaphrodite (4)). These states are self-explanatory and independently meaningful. All states are necessary to describe the full range of the characteristic, and every form of expression can be described by a single state. The order of states is not important. As a rule, the characteristics are not influenced by environment.

2.3.3.3 “Quantitative characteristics” are those where the expression covers the full range of variation from one extreme to the other. The expression can be recorded on a one-dimensional, continuous or discrete, linear scale. The range of expressions is divided into a number of states for the purpose of description (e.g. length of stem: very short (1), short (3), medium (5), long (7), very long (9)). The division seeks to provide, as far as practical, an even distribution across the scale. The Test Guidelines do not specify the difference needed for distinctness. The states of expression should, however, be meaningful for DUS assessment.

2.3.3.4 In the case of “pseudo-qualitative characteristics” the range of expression is at least partly continuous, but varies in more than one dimension (e.g. shape: ovate (1), elliptic (2), circular (3), obovate (4)) and cannot be adequately described by just defining two ends of a linear range. In a similar way to qualitative (discontinuous) characteristics – hence the term “pseudo-qualitative” – each individual state of expression needs to be identified to adequately describe the range of the characteristic.

### 2.3.4 Types of scales of data

The possibility to use specific procedures for the assessment of distinctness, uniformity and stability depends on the scale level of the data which are recorded for a characteristic. The scale level of data depends on the type of expression of the characteristic and on the way of recording this expression. The type of scale may be quantitative or qualitative.

#### 2.3.4.1 Quantitatively scaled data (metric or ordinal scaled data)

Quantitatively scaled data are all data which are recorded by measuring or counting. Weighing is a special form of measuring. Quantitatively scaled data can have a continuous or a discrete distribution. Continuous data result from measurements. They can take every value out of the defined range. Discrete quantitative data result from counting.

Examples

Quantitatively scaled data	Example	Example number
- continuous	Plant length in cm.	1
- discrete	Number of stamens	2

For description of the states of expression, see Table 6.

The continuous quantitatively scaled data for the characteristic “Plant length” are measured on a continuous scale with defined units of assessment. A change of unit of measurement e.g. from cm into mm is only a question of precision and not a change of type of scale.

The discrete quantitatively scaled data of the characteristic “Number of stamens “ are assessed by counting (1, 2, 3, 4, and so on). The distances between the neighboring units of assessment are constant and for this example equal to 1. There are no real values between two neighboring units but it is possible to compute an average which falls between those units.

In biometrical terminology, quantitative scales are referred to as metric scales or cardinal scales. Quantitative scales can be subdivided into ratio scales and interval scales.

2.3.4.1.1 *Ratio scale*

**[TWC Chairperson:** To review if this paragraph is relevant for DUS testing]

A ratio scale is a quantitative scale with a defined absolute zero point. There is always a constant non-zero distance between two adjacent expressions. Ratio scaled data may be continuous or discrete.

*The absolute zero point:*

The definition of an absolute zero point makes it possible to define meaningful ratios. This is a requirement for the construction of index numbers (e.g. the ratio of length to width). An index is the combination of at least two characteristics. In the General Introduction, this is referred to as a combined characteristic (see document TG/1/3, Section 4.6.3).

It is also possible to calculate ratios between the expression of different varieties. For example, in the characteristic ‘Plant length’ assessed in cm, there is a lower limit for the expression which is ‘0 cm’ (zero). It is possible to calculate the ratio of length of plant of variety ‘A’ to length of plant of variety ‘B’ by division:

**[TWC Chairperson:** to review if this paragraph is relevant for DUS testing]

Length of plant of variety 'A' = 80 cm  
 Length of plant of variety 'B' = 40 cm  
 Ratio = Length of plant of variety 'A' / Length of plant of variety 'B'  
 = 80 cm / 40 cm  
 = 2.

So it is possible in this example to state that plant 'A' is double the length of plant 'B'. The existence of an absolute zero point ensures an unambiguous ratio.

The ratio scale is the highest classification of the scales (Table 2). That means that ratio scaled data include the highest information about the characteristic and it is possible to use many statistical procedures (Chapter 7 [*cross ref.*]).

The examples 1 and 2 (Table 6) are examples for characteristics with ratio scaled data.

#### 2.3.4.1.2 Interval scale

An Interval scale is a quantitative scale without a defined absolute zero point. There is always a constant non-zero distance between two adjacent expressions. Interval scaled data may be distributed continuously or discretely.

An example for a discrete interval scaled characteristic is 'Time of beginning of flowering' measured as date which is given as example 6 in Table 6. This characteristic is defined as the number of days from April 1. The definition is useful but arbitrary and April 1 is not a natural limit. It would also be possible to define the characteristic as the number of days from January 1.

It is not possible to calculate a meaningful ratio between two varieties which should be illustrated with the following example:

Variety 'A' begins to flower on May 30 and variety 'B' on April 30

Case I) Number of days from April 1 of variety 'A' = 60  
 Number of days from April 1 of variety 'B' = 30

$$\text{Ratio}_I = \frac{\text{Number of days from April 1 of variety 'A' } 60 \text{ days}}{\text{Number of days from April 1 of variety 'B' } 30 \text{ days}} = \frac{60}{30} = 2$$

Case II) Number of days from January 1 of variety 'A' = 150  
 Number of days from January 1 of variety 'B' = 120

$$\text{Ratio}_{II} = \frac{\text{Number of days from January 1 of variety 'A' } 150 \text{ days}}{\text{Number of days from January 1 of variety 'B' } 120 \text{ days}} = \frac{150}{120} = 1.25$$

$$\text{Ratio}_I = 2 > 1.25 = \text{Ratio}_{II}$$



It is impossible to state that the time of flowering of variety ‘A’ is twice that of variety ‘B’. The ratio depends on the choice of the zero point of the scale. This kind of scale is defined as an “Interval scale”: a quantitative scale without a defined absolute zero point.

The interval scale is lower classified than the ratio scale (Table 2). Fewer statistical procedures can be used with interval scaled data than with ratio scaled data (see Part I: Section 2.3.7 [*cross ref.*]). The interval scale is theoretically the minimum scale level to calculate arithmetic mean values.

#### 2.3.4.2 Qualitatively scaled data

Qualitatively scaled data are data which can be arranged in different discrete qualitative categories. Usually they result from visual assessment. Subgroups of qualitative scales are ordinal and nominal scales.

##### 2.3.4.2.1 Ordinal scale

**[TWC Chairperson: example for a non-quantitative characteristic to be provided]**

Ordinally scaled data are qualitative data of which discrete categories can be arranged in an ascending or descending order. They result from visually assessed quantitative characteristics.

Example:

Qualitative data	Example	Example number
- ordinal	Intensity of anthocyanin	3

For description of the states of expressions, see Table 6.

An ordinal scale consists of numbers which correspond to the states of expression of the characteristic (notes). The expressions vary from one extreme to the other and thus they have a clear logical order. It is not possible to change this order, but it is not important which numbers are used to denote the categories. In some cases ordinal data may reach the level of discrete interval scaled data or of discrete ratio scaled data (Chapter 6 [*cross ref.*]).

The distances between the discrete categories of an ordinal scale are not exactly known and not necessarily equal. Therefore, an ordinal scale does not fulfil the condition to calculate arithmetic mean values, which is the equality of intervals throughout the scale.

The ordinal scale is lower classified than the interval scale (Table 2). Less statistical procedures can be used for ordinal scale than for each of the higher classified scale data (see Part I: Section 3.7 [*cross ref.*]).

##### 2.3.4.2.2 Nominal scale

Nominal scaled qualitative data are qualitative data without any logical order of the discrete categories.

Examples:

Qualitative data	Example	Example number
- nominal	Sex of plant	4
- nominal with two states	Leaf blade: variegation	5

For description of the states of expressions, see Table 6.

A nominal scale consists of numbers which correspond to the states of expression of the characteristic, which are referred to in the Test Guidelines as notes. Although numbers are used for designation there is no inevitable order for the expressions and so it is possible to arrange them in any order.

Characteristics with only two categories (dichotomous characteristic) are a special form of nominal scales.

The nominal scale is the lowest classification of the scales (Table 2). Few statistical procedures are applicable for evaluations (Chapter 7 [*cross ref.*] ).

The different types of scales are summarized in the following table.

*Table 2: Types of scales and scale levels*

**[TWC Chairperson: To modify the table for consistency with the subsequent paragraphs]**

Type of scale		Description	Distribution	Data recording	Scale Level
quantitative (metric)	ratio	constant distances with absolute zero point	Continuous	Absolute Measurements	High
			Discrete	Counting	
	interval	constant distances without absolute zero point	Continuous	Relative measurements	↑
			Discrete	Date	
qualitative with underlying quantitative variable	ordinal	Ordered expressions with varying distances	Discrete	Visually assessed notes	↑
qualitative	nominal	No order, no distances	Discrete	Visually assessed notes	Low

From the statistical point of view a characteristic is only considered at the level of data which has been recorded, whether for analysis or for describing the expression of the characteristic. Therefore, characteristics with quantitative data are denoted as quantitative characteristics and characteristics with ordinal and nominal scaled data as qualitative characteristics.

### 2.3.5 Scale levels for variety description

The description of varieties is based on the states of expression (notes) which are given in the Test Guidelines for the specific crop. In the case of visual assessment, the notes from the Test Guidelines are usually used for recording the characteristic as well as for the assessment of DUS. The notes are distributed on a nominal or ordinal scale (see Part I: Section 2.3.4.2 [*cross ref.*]). For measured or counted characteristics, DUS assessment is based on the recorded values and the recorded values are transformed into states of expression only for the purpose of variety description.

### 2.3.6 Relation between types of expression of characteristics and scale levels of data

2.3.6.1 Records taken for the assessment of qualitative characteristics are distributed on a nominal scale, for example “Sex of plant”, “Leaf blade: variegation” (Table 6, examples 4 and 5).

2.3.6.2 For quantitative characteristics the scale level of data depends on the method of assessment. They can be recorded on a quantitative or ordinal scale. For example, “Length of plant” can be recorded by measurements resulting in ratio scaled continuous quantitative data. However, visual assessment on a 1 to 9 scale may also be appropriate. In this case, the recorded data are qualitatively scaled (ordinal scale) because the size of intervals between the midpoints of categories is not exactly the same.

Remark: In some cases visually assessed data on quantitative characteristics may be handled as measurements. The possibility to apply statistical methods for quantitative data depends on the precision of the assessment and the robustness of the statistical procedures. In the case of very precise visually assessed quantitative characteristics the usually ordinal data may reach the level of discrete interval scaled data or of discrete ratio scaled data.

2.3.6.3 A pseudo-qualitative type of characteristic is one in which the expression varies in more than one dimension. The different dimensions are combined in one scale. At least one dimension is quantitatively expressed. The other dimensions may be qualitatively expressed or quantitatively expressed. The scale as a whole has to be considered as a nominal scale (e.g. “Shape”, “Flower color”; Table 6, examples 7 and 8).

2.3.6.4 In the case of using the off-type procedure for the assessment of uniformity the recorded data are nominally scaled. The records fall into two qualitative classes: plants belonging to the variety (true-types) and plants not belonging to the variety (off-types). The type of scale is the same for qualitative, quantitative and pseudo-qualitative characteristics.

2.3.6.5 The relation between the type of characteristics (process level 1) and the type of scale of data recorded for the assessment of distinctness and uniformity is described in Table 3. A qualitative characteristic is recorded on a nominal scale for distinctness (state of expression) and for uniformity (true-types vs. off-types). Pseudo-qualitative characteristics are recorded on a combined scale for distinctness (state of expression) and on a nominal scale for uniformity (true-types vs. off-types). Quantitative characteristics are recorded on an ordinal, interval or ratio scale for the assessment of distinctness depending on the characteristic and the method of assessment. If the records are taken from single plants the same data may be used for the assessment of distinctness and uniformity. If distinctness is

assessed on the basis of a single record of a group of plants, uniformity has to be judged with the off-type procedure (nominal scale).

Table 3: Relation between type of characteristic and type of scale of assessed data

Procedure	Type of scale (level 2)	Distribution	Type of characteristic (level 1)		
			Quantitative	Pseudo-qualitative	Qualitative
Distinctness	ratio	Continuous	▪		
		Discrete	▪		
	interval	Continuous	▪		
		Discrete	▪		
	ordinal	Discrete	▪		
	combined	Discrete		▪	
nominal	Discrete			▪	
Uniformity	ratio	Continuous	▪		
		Discrete	▪		
	interval	Continuous	▪		
		Discrete	▪		
	ordinal	Discrete	▪		
	combined	Discrete	▪		
nominal	Discrete	▪	▪	▪	

2.3.7 Relation between method of observation of characteristics, scale levels of data and recommended statistical procedures

2.3.7.1 The scale level of data and the way of observation of characteristics are most important conditions for the application of different statistical procedures. There are four possible ways to observe characteristics (see document TGP/7 “Development of Test Guidelines”):

- MG: single record for a group of plants or parts of plants based on measurement(s)
- MS: records for a number of single, individual plants or parts of plants obtained by measurement
- VG: single record for a group of plants or parts of plants based on visual observation(s)
- VS: records for a number of single, individual plants or parts of plants obtained by visual observation.

2.3.7.2 The observation method depends primarily on the variation within and between varieties and affects the choice of the statistical method. All of the four observation methods may be relevant for the assessment of distinctness. For the assessment of uniformity observations must be done on individual plants. Consequently only MS or VS are appropriate. The indication of the method of observation of characteristics in the Test Guidelines refers only to the assessment of distinctness.

**[TWC Chairperson: To update these paragraphs in accordance with any changes to TGP/7 and TGP/9]**

2.3.7.3 Established statistical procedures can be used for the assessment of distinctness and uniformity considering the scale level and some further conditions such as the degree of freedom or unimodality (Tables 4 and 5).

2.3.7.4 The relation between the expression of characteristics and the scale levels of data for the assessment of distinctness and uniformity is summarized in Table 6.

Table 4: Statistical procedures for the assessment of distinctness

Type of scale	Distribution	Observation method	Procedure <sup>1)</sup> and further Conditions	Reference document
ratio	continuous	MS MG (VS) <sup>1)</sup>	COY-D Normal distribution, df $\geq 20$	TGP/9
	discrete		long term LSD Normal distribution, df $< 20$	
interval	continuous		2 out of 3 method (LSD 1%) Normal distribution, df $\geq 20$	
	discrete			
ordinal	discrete	VG	See explanation for QN characteristics in TGP/9 Sections 5.2.2 and 5.2.3,	TGP/9
		VS	See explanation for QN characteristics in TGP/9 Section 5.2.4	TWC/ 14/12
Combination of ordinal or ordinal and nominal scales	discrete	VG (VS) <sup>32)</sup>	See explanation for PQ characteristics in TGP/9 Sections 5.2.2 and 5.2.3	TGP/9
nominal	discrete	VG (VS) <sup>2)</sup>	See explanation for QL characteristics in TGP/9 Sections 5.2.2 and 5.2.3	TGP/9

1) see remark in Chapter 6 [cross ref.]

2) normally VG but VS would be possible

Table 5: Statistical procedures for the assessment of uniformity

Type of scale	Distribution	observation method	Procedure <sup>1)</sup> and Further Conditions	Reference document
ratio	continuous	MS	COY-U Normal distribution 2 out of 3 method ( $s_c^2 \leq 1.6s_s^2$ ) Normal distribution LSD for untransformed percentage of off-types	TGP/10
	discrete	MS		
interval	continuous	VS		
	discrete			
ordinal	discrete	VS	threshold model	TWC/ 14/12
Combination of ordinal or ordinal and nominal scales	discrete		There is no case where uniformity is assessed on combined scaled data	
nominal	discrete	VS	off-type procedure for dichotomous (binary) data	TGP/10

Table 6: Relation between expression of characteristics and scale levels of data for the assessment of distinctness and uniformity

Example	Name of characteristic	Distinctness			Uniformity		
		Unit of assessment	Description (states of expression)	Type of scale	Unit of assessment	Description (states of expression)	Type of scale
1	Length of plant	cm	assessment in cm without digits after decimal point	ratio scaled continuous quantitative data	cm	assessment in cm without digits after decimal point	ratio scaled continuous quantitative data
					True-type	Number of plants belonging to the variety	nominally scaled qualitative data
					Off-type	Number of off-types	
2	Number of stamens	counts	1, 2, 3, ... , 40,41, ...	ratio scaled discrete quantitative data	counts	1, 2, 3, ... , 40,41, ...	ratio scaled discrete quantitative data
3	Intensity of anthocyanin	1 2 3 4 5 6 7 8 9	very low very low to low low low to medium medium medium to high high high to very high very high	ordinally scaled qualitative data (with an underlying quantitative variable)	True-type	Number of plants belonging to the variety	nominally scaled qualitative data
					Off-type	Number of off-types	
4	Sex of plant	1 2 3 4	dioecious female dioecious male monoecious unisexual monoecious hermaphrodite	nominally scaled qualitative data	True-type	Number of plants belonging to the variety	nominally scaled qualitative data
					Off-type	Number of off-types	

Example	Name of characteristic	Distinctness			Uniformity		
		Unit of assessment	Description (states of expression)	Type of scale	Unit of assessment	Description (states of expression)	Type of scale
5	Leaf blade: variegation	1	absent	nominally scaled qualitative data	True-type	Number of plants belonging to the variety	nominally scaled qualitative data
		9	present		Off-type		
6	Time of beginning of flowering	date	e.g. May 21, 51 <sup>st</sup> day from April 1	interval scaled discrete quantitative data	date	e.g. May 21, 51 <sup>st</sup> day from April 1	interval scaled discrete quantitative data
					True-type	Number of plants belonging to the variety	nominally scaled qualitative data
7	Shape	1 2 3 4 5 6 7	deltate ovate elliptic obovate obdeltate circular oblate	combination of ordinal and nominal scaled discrete qualitative data	True-type	Number of plants belonging to the variety	nominally scaled qualitative data
					Off-type		
8	Flower color	1 2 3 4 5 6 7 8 9 10	dark red medium red light red white light blue medium blue dark blue red violet violet blue violet	combination of ordinal and nominal scaled discrete qualitative data	True-type	Number of plants belonging to the variety	nominally scaled qualitative data
					Off-type		



## PART II: TECHNIQUES USED IN DUS EXAMINATION

### 1. Methods for Assessing Uniformity on the Basis of Off-Types

#### 1.1 Fixed Population Standard

##### 1.1.1 Introduction

1.1.1.1 TGP/10 Section 4 provides guidance on the uniformity assessment on the basis of off-types. This section describes the method of assessing uniformity by comparing the number of off-types observed to a fixed population standard. In particular, the tables included in Section 3.1.12 provide the recommendation on the acceptable number of off-types in different sample sizes for specific population standards and acceptance probabilities within acceptable levels of errors.

1.1.1.2 This document also outlines procedures for when more than a single test (more than one year for instance) is used and explains the possibility of using sequential tests to minimize testing effort.

1.1.1.3 When testing for uniformity on the basis of a sample, there will always be some risk of making a wrong decision. The risks can be reduced by increasing the sample size but at a greater cost. The aim of the statistical procedure described here is to achieve an acceptable balance between risks.

1.1.1.4 The procedures described below require the user to define an acceptance standard (called the *population standard*) for the crop in question. The methods described then show how to determine the sample size and the maximum number of off-types allowed for various levels of risks.

1.1.1.5 The population standard is the maximum percentage of off-types that would be accepted if all individuals of the variety could be examined.

#### 1.1.2 Recommendations on the fixed population standard method of assessing uniformity by the number of off-types

1.1.2.1 This method is recommended for use in assessing the uniformity by number of off-types with a fixed population standard.

1.1.2.2 The sample size and acceptable number of off-types employed depend on the crop.

#### 1.1.3 Errors in testing for off-types

[TWF: to review whether such an explanation is appropriate in the light of sample sizes used in the TG. The document should reflect the positive experience of UPOV in the existing sample sizes]

[TWC Chairperson: to review the terms used and, in particular, terms such as “wrong decisions”]

1.1.3.1 As mentioned, there will be some risk of making wrong decisions. Two types of error exist:

(a) Declaring that the variety lacks uniformity when it in fact meets the standard for the crop. This is known as “type I error.”

(b) Declaring that the variety is uniform when it in fact does not meet the standard for the crop. This is known as “type II error.”

1.1.3.2 The types of error can be summarized in the following table:

True state of the variety	Decision made on variety	
	Acceptance as uniform	Rejection as non-uniform
uniform	correctly accepted	type I error
non-uniform	type II error	correctly rejected

1.1.3.3 The probability of correctly accepting a uniform variety is called the acceptance probability and is linked to the probability of type I error by the relation:

$$\text{“Acceptance probability”} + \text{“probability of type I error”} = 100\%$$

1.1.3.4 The probability of type II error depends on “how non-uniform” the candidate variety is. If it is much more non-uniform than the population standard then the probability of type II error will be small and there will be a small probability of accepting such a variety. If, on the other hand, the candidate variety is only slightly more non-uniform than the standard, there is a large probability of type II error. The probability of acceptance will approach the acceptance probability for a variety with a level of uniformity near to the population standard.

1.1.3.5 Because the probability of type II error is not fixed but depends on “how non-uniform” the candidate variety is, this probability can be calculated for different degrees of non-uniformity. This document gives probabilities of type II error for three degrees of non-uniformity: 2, 5 and 10 times the population standard.

1.1.3.6 In general, the probability of making errors will be decreased by increasing the sample size and increased by decreasing the sample size.

1.1.3.7 For a given sample size, the balance between the probabilities of making type I and type II errors may be altered by changing the number of off-types allowed.

1.1.3.8 If the number of off-types allowed is increased, the probability of type I error is decreased but the probability of type II error is increased. On the other hand, if the number of off-types allowed is decreased, the probability of type I errors is increased while the probability of type II errors is decreased.

1.1.3.9 By allowing a very high number of off-types it will be possible to make the probability of type I errors very low (or almost zero). However, the probability of making type II errors will now become (unacceptably) high. If only a very small number of off-types

is allowed, the result will be a small probability of type II errors and an (unacceptably) high probability of type I errors. This will be illustrated by examples.

#### 1.1.4 Examples

##### Example 1

1.1.4.1 From experience, a reasonable standard for the crop in question is found to be 1%. So the population standard is 1%. Assume that a single test with a maximum of 60 plants is used. From tables 4, 10 and 16 (chosen to give a range of target acceptance probabilities), the following schemes are found:

Scheme	Sample size	Target acceptance probability*	Maximum number of off-types
a	60	90%	2
b	53	90%	1
c	60	95%	2
d	60	99%	3

1.1.4.2 From the figures 4, 10 and 16, the following probabilities are obtained for the type I error and type II error for different percentages of off-types (denoted by  $P_2$ ,  $P_5$  and  $P_{10}$  for 2, 5 and 10 times the population standard).

Scheme	Sample size	Maximum number of off-types	Probabilities of error (%)			
			Type I	Type II		
				$P_2 = 2\%$	$P_5 = 5\%$	$P_{10} = 10\%$
a	60	2	2	88	42	5
b	53	1	10	71	25	3
c	60	2	2	88	42	5
d	60	3	0.3	97	65	14

1.1.4.3 The table lists four different schemes and they should be examined to see if one of them is appropriate to use. (Schemes a and c are identical since there is no scheme for a sample size of 60 with a probability of type I error between 5 and 10%). If it is decided to ensure that the probability of a type I error should be very small (scheme d) then the probability of the type II error becomes very large (97, 65 and 14%) for a variety with 2.5 and 10% of off-types, respectively. The best balance between the probabilities of making the two types of error seems to be obtained by allowing one off-type in a sample of 53 plants (scheme b).

\* See paragraph 54

Example 2

1.1.4.4 In this example, a crop is considered where the population standard is set to 2% and the number of plants available for examination is only 6.

1.1.4.5 Using the tables and the figures 3, 9 and 15, the following schemes a-d are found:

Scheme	Sample size	Acceptance probability	Maximum number of off-types	Probability of error (%)			
				Type I	Type II		
					P <sub>2</sub> = 4%	P <sub>5</sub> = 10%	P <sub>10</sub> = 20%
a	6	90	1	0.6	98	89	66
b	5	90	0	10	82	59	33
c	6	95	1	0.6	98	89	66
d	6	99	1	0.6	98	89	66
e	6		0	11	78	53	26

1.1.4.6 Scheme e of the table is found by applying the formulas (1) and (2) shown later in this document.

1.1.4.7 This example illustrates the difficulties encountered when the sample size is very low. The probability of erroneously accepting a non-uniform variety (a type II error) is large for all the possible situations. Even when all five plants must be uniform for a variety to be accepted (scheme b), the probability of accepting a variety with 20% of off-types is still 33%.

1.1.4.8 It should be noted that a scheme where all six plants must be uniform (scheme e) gives slightly smaller probabilities of type II errors, but now the probability of the type I error has increased to 11%.

1.1.4.9 However, scheme e may be considered the best option when only six plants are available in a single test for a crop where the population standard has been set to 2%.

Example 3

1.1.4.10 In this example we reconsider the situation in example 1 but assume that data are available for two years. So the population standard is 1% and the sample size is 120 plants (60 plants in each of two years).

1.1.4.11 The following schemes and probabilities are obtained from the tables and figures 4, 10 and 16:

Scheme	Sample size	Acceptance probability	Maximum number of off-types	Probability of error (%)			
				Type I	Type II		
					P <sub>2</sub> = 2%	P <sub>5</sub> = 5%	P <sub>10</sub> = 10%
a	120	90	3	3	78	15	<0.1
b	110	90	2	10	62	8	<0.1
c	120	95	3	3	78	15	<0.1
d	120	99	4	0.7	91	28	1

1.1.4.12 Here the best balance between the probabilities of making the two types of error is obtained by scheme c, i.e. to accept after two years a total of three off-types among the 120 plants examined.

1.1.4.13 Alternatively a two-stage testing procedure may be set up. Such a procedure can be found for this case by using formulae (3) and (4) later in this document.

1.1.4.14 The following schemes can be obtained:

Scheme	Sample size	Acceptance probability	Largest number for acceptance after year 1	Largest number before reject in year 1	Largest number to accept after 2 years
e	60	90	can never accept	2	3
f	60	95	can never accept	2	3
g	60	99	can never accept	3	4
h	58	90	1	2	2

1.1.4.15 Using the formulas (3), (4) and (5) the following probabilities of errors are obtained:

Scheme	Probability of error (%)				Probability of testing in a second year
	Type I	Type II			
		P <sub>2</sub> = 2%	P <sub>5</sub> = 5%	P <sub>10</sub> = 10%	
e	4	75	13	0.1	100
f	4	75	13	0.1	100
g	1	90	27	0.5	100
h	10	62	9	0.3	36

1.1.4.16 Schemes e and f both result in a probability of 4% for rejecting a uniform variety (type I error) and a probability of 13% for accepting a variety with 5% off-types (type II error). The decision is:

- Never accept the variety after 1 year
- More than 2 off-types in year 1: reject the variety and stop testing
- Between and including 0 and 2 off types in year 1: do a second year test
- At most 3 off-types after 2 years: accept the variety
- More than 3 off-types after 2 years: reject the variety

1.1.4.17 Alternatively, scheme h may be chosen but scheme g seems to have a too large probability of type II errors compared with the probability of type I error.

1.1.4.18 Scheme h has the advantage of often allowing a final decision to be taken after the first test (year) but, as a consequence, there is a higher probability of a type I error.

*[TWC: paragraphs 1.1.5.17 and 1.1.5.18 to be elaborated further]*

#### Example 4

1.1.4.19 In this example, we assume that the population standard is 3% and that we have 8 plants available in each of two years.

1.1.4.20 From the tables and figures 2, 8 and 14, we have:

Scheme	Sample size	Acceptance probability	Maximum number of off-types	Probability of error (%)			
				Type I	Type II		
					P <sub>2</sub> = 6%	P <sub>5</sub> = 15%	P <sub>10</sub> = 30%
a	16	90	1	8	78	28	3
b	16	95	2	1	93	56	10
c	16	99	3	0.1	99	79	25

1.1.4.21 Here the best balance between the probabilities of making the two types of error is obtained by scheme a.

### 1.1.5 Introduction to the tables and figures

1.1.5.1 In the TABLES AND FIGURES section (Part II: Section 3.1.12 [*cross ref.*]), there are 21 table and figure pairs corresponding to different combinations of population standard and acceptance probability. These are design to be applied to a single off-type test. An overview of the tables and the figures are given in table A.

1.1.5.2 Each table shows the maximum numbers of off-types (k) with the corresponding ranges in sample sizes (n) for the given population standard and acceptance probability. For example, in table 1 (population standard 5%, acceptance probability ≥ 90%), for a maximum set at 2 off-types, the corresponding sample size (n) is in the range from 11 to 22. Likewise,

if the maximum number of off-types (k) is 10, the corresponding sample size (n) to be used should be in the range 126 to 141.

1.1.5.3 For small sample sizes, the same information is shown graphically in the corresponding figures (figures (1 to 21). These show the actual risk of rejecting a uniform variety and the probability of accepting a variety with a true proportion of off-types 2 times (2P), 5 times (5P) and 10 times (10P) greater than the population standard. (To ease the reading of the figure, lines connect the risks for the individual sample sizes, although the probability can only be calculated for each individual sample size).

Table A. Overview of table and figure 1 to 18.

Population standard %	Acceptance probability %	See table and figure no.
10	>90	19
10	>95	20
10	>99	21
5	>90	1
5	>95	7
5	>99	13
3	>90	2
3	>95	8
3	>99	14
2	>90	3
2	>95	9
2	>99	15
1	>90	4
1	>95	10
1	>99	16
0.5	>90	5
0.5	>95	11
0.5	>99	17
0.1	>90	6
0.1	>95	12
0.1	>99	18

1.1.5.4 When using the tables the following procedure is suggested:

**[TWC Chairperson: to be revised in accordance with the use of the tables set out in TGP/10 and with established practice]**

(a) Choose the relevant population standard.

(b) Write down the different relevant decision schemes (combinations of sample size and maximum number of off-types), with the probabilities of type I and type II errors read from the figures.

(c) Choose the decision scheme with the best balance between the probabilities of errors.

1.1.5.5 The use of the tables and figures is illustrated in the example section.

#### 1.1.6 Detailed description of the method for one single test

The mathematical calculations are based on the binomial distribution and it is common to use the following terms:

(a) The percentage of off-types to be accepted in a particular case is called the “population standard” and symbolized by the letter P.

(b) The “acceptance probability” is the probability of accepting a variety with P% of off-types. However, because the number of off-types is discrete, the actual probability of accepting a uniform variety varies with sample size but will always be greater than or equal to the “acceptance probability.” The acceptance probability is usually denoted by  $100 - \alpha$ , where  $\alpha$  is the percent probability of rejecting a variety with P% of off-types (i.e. type I error probability). In practice, many varieties will have less than P% off-types and hence the type I error will in fact be less than  $\alpha$  for such varieties.

(c) The number of plants examined in a random sample is called the sample size P and  $P_q$  are expressed here as proportions, i.e. percents divided by 100.

and denoted by n.

(d) The maximum number of off-types tolerated in a random sample of size n is denoted by k.

(e) The probability of accepting a variety with more than P% off-types, say  $P_q\%$  of off-types, is denoted by the letter  $\beta$  or by  $\beta_q$ .

(f) The mathematical formulae for calculating the probabilities are:

$$\alpha = 100 - 100 \sum_{i=0}^k \binom{n}{i} P^i (1-P)^{n-i} \quad (1)$$

$$\beta_q = 100 \sum_{i=0}^k \binom{n}{i} P_q^i (1-P_q)^{n-i} \quad (2)$$

P and  $P_q$  are expressed here as proportions, i.e. percents divided by 100.



### 1.1.7 More than one single test (year)

1.1.7.1 Often a candidate variety is grown in two (or three years). The question then arises of how to combine the uniformity information from the individual years. Two methods will be described:

- (a) Make the decision after two (or three) years based on the total number of plants examined and the total number of off-types recorded. (A combined test).
- (b) Use the result of the first year to see if the data suggests a clear decision (reject or accept). If the decision is not clear then proceed with the second year and decide after the second year. (A two-stage test).

1.1.7.2 However, there are some alternatives (e.g. a decision may be made in each year and a final decision may be reached by rejecting the candidate variety if it shows too many off-types in both (or two out of three years)). Also there are complications when more than one single year test is done. It is therefore suggested that a statistician should be consulted when two (or more) year tests have to be used.

### 1.1.8 Detailed description of the methods for more than one single test

#### 1.1.8.1 Combined Test

The sample size in test  $i$  is  $n_i$ . So after the last test we have the total sample size  $n = \sum n_i$ . A decision scheme is set in exactly the same way as if this total sample size had been obtained in a single test. Thus, the total number of off-types recorded through the tests is compared with the maximum number of off-types allowed by the chosen decision scheme.

#### 1.1.8.2 Two-stage Test

1.1.8.2.1 The method for a two-year test may be described as follows: In the first year take a sample of size  $n$ . Reject the candidate variety if more than  $r_1$  off-types are recorded and accept the candidate variety if less than  $a_1$  off-types are recorded. Otherwise, proceed to the second year and take a sample of size  $n$  (as in the first year) and reject the candidate variety if the total number of off-types recorded in the two years' test is greater than  $r$ . Otherwise, accept the candidate variety. The final risks and the expected sample size in such a procedure may be calculated as follows:

$$\begin{aligned}
 \alpha &= P(K_1 > r_1) + P(K_1 + K_2 > r \mid K_1) \\
 &= P(K_1 > r_1) + P(K_2 > r - K_1 \mid K_1) \\
 &= \sum_{i=r_1+1}^n \binom{n}{i} P^i (1-P)^{n-i} + \sum_{i=\alpha_1}^{r_1} \binom{n}{i} P^i (1-P)^{n-i} \sum_{j=r-i+1}^n \binom{n}{i} P^j (1-P)^{n-j} \quad (3)
 \end{aligned}$$

$$\begin{aligned}
 \beta_q &= P(K_1 < \alpha_1) + P(K_1 + K_2 \leq r \mid K_1) \\
 &= P(K_1 < \alpha_1) + P(K_2 \leq r - K_1 \mid K_1) \\
 &= \sum_{i=0}^{\alpha_1-1} \binom{n}{i} P_q^i (1-P_q)^{n-i} + \sum_{i=\alpha_1}^{r_1} \binom{n}{i} P_q^i (1-P_q)^{n-i} \sum_{j=0}^{r-i} \binom{n}{i} P_q^j (1-P_q)^{n-j} \quad (4)
 \end{aligned}$$

$$n_e = n \left( 1 + \sum_{i=\alpha_1}^{r_1} \binom{n}{i} P^i (1-P)^{n-i} \right) \quad (5)$$

where

$P$  = population standard

$\alpha$  = probability of actual type I error for  $P$

$\beta_q$  = probability of actual type II error for  $q P$

$n_e$  = expected sample size

$r_1, \alpha_1$  and  $r$  are decision-parameters

$P_q = q$  times population standard =  $q P$

$K_1$  and  $K_2$  are the numbers of off-types found in years 1 and 2 respectively.

The decision parameters,  $\alpha_1, r_1$  and  $r$ , may be chosen according to the following criteria:

- (a)  $\alpha$  must be less than  $\alpha_0$ , where  $\alpha_0$  is the maximum type I error, i.e.  $\alpha_0$  is 100 minus the required acceptance probability
- (b)  $\beta_q$  (for  $q=5$ ) should be as small as possible but not smaller than  $\alpha_0$
- (c) if  $\beta_q$  (for  $q=5$ )  $< \alpha_0 n_e$  should be as small as possible.

1.1.8.2.2 However, other strategies are available. No tables/figures are produced here as there may be several different decision schemes that satisfy a certain set of risks. It is suggested that a statistician should be consulted if a 2-stage test (or any other sequential tests) is required.

### 1.1.8.3 Sequential tests

The two-stage test mentioned above is a type of sequential test where the result of the first stage determines whether the test needs to be continued for a second stage. Other types of sequential tests may also be applicable. It may be relevant to consider such tests when the practical work allows analyses of off-types to be carried out at certain stages of the examination. The decision schemes for such methods can be set up in many different ways

and it is suggested that a statistician should be consulted when sequential methods are to be used.

#### 1.1.9 Note on type I and type II errors

1.1.9.1 We cannot in general obtain type I-errors that are nice pre-selected values because the number of off-types is discrete. The scheme a of example 2 with 6 plants above showed that we could not obtain an  $\alpha$  of 10% - our actual  $\alpha$  became 0.6%. Changing the sample size will result in varying  $\alpha$  and  $\beta$  values. Figure 3 - as an example - shows that  $\alpha$  gets closer to its nominal values at certain sample sizes and that this is also the sample size where  $\beta$  is relatively small.

1.1.9.2 Larger sample sizes are generally beneficial. With same acceptance probability, a larger sample will tend to have proportionally less probability of type II errors. Small sample sizes result in high probabilities of accepting non-uniform varieties. The sample size should therefore be chosen to give an acceptably low level of type II errors. However small increases in the sample size may not always be advantageous. For instance, a sample size of five gives  $\alpha = 10\%$  and  $\beta_2 = 82\%$  whereas a sample size of six gives  $\alpha = 0.6\%$  and  $\beta_2 = 98\%$ . It appears that the sample sizes, which give  $\alpha$ -values in close agreement with the acceptance probability are the largest in the range of sample sizes with a specified maximum number of off-types. Thus, the largest sample sizes in the range of sample sizes with a given maximum number of off-types should be used.

#### 1.1.10 Definition of statistical terms and symbols

The statistical terms and symbols used have the following definitions:

*Population standard.* The percentage of off-types to be accepted if all the individuals of a variety could be examined. The population standard is fixed for the crop in question and is based on experience.

*Acceptance probability.* The probability of accepting a uniform variety with P% of off-types. Here P is population standard. However, note that the actual probability of accepting a uniform variety will always be greater than or equal to the acceptance probability in the heading of the table and figures. The probability of accepting a uniform variety and the probability of a type I error sum to 100%. For example, if the type I error probability is 4%, then the probability of accepting a uniform variety is  $100 - 4 = 96\%$ , see e.g. figure 1 for  $n=50$ ). The type I error is indicated on the graph in the figures by the sawtooth peaks between 0 and the upper limit of type I error (for instance 10 on figure 1). The decision schemes are defined so that the actual probability of accepting a uniform variety is always greater than or equal to the acceptance probability in the heading of the table.

*Type I error:* The error of rejecting a uniform variety.

*Type II error:* The error of accepting a variety that is too non-uniform.

P Population standard

$P_q$  The assumed true percentage of off-types in a non-uniform variety.  $P_q = q P$ .

In the present document  $q$  is equal to 2, 5 or 10. These are only 3 examples to help the visualization of type II errors. The actual percentage of off-types in a variety may take any value. For instance we may examine different varieties which in fact may have respectively 1.6%, 3.8%, 0.2%, ... of off-types.

- n Sample size
- k Maximum number of off-types allowed
- $\alpha$  Probability of type I error
- $\beta$  Probability of type II error

3.1.11 Tables and figures

Table and figure 1:

Population Standard = 5%  
Acceptance Probability  $\geq 90\%$   
n=sample size, k=maximum number of off-types

	n	k
1	to 2	0
3	to 10	1
11	to 22	2
23	to 35	3
36	to 49	4
50	to 63	5
64	to 78	6
79	to 94	7
95	to 109	8
110	to 125	9
126	to 141	10
142	to 158	11
159	to 174	12
175	to 191	13
192	to 207	14
208	to 224	15
225	to 241	16
242	to 258	17
259	to 275	18
276	to 292	19
293	to 310	20
311	to 327	21
328	to 344	22
345	to 362	23
363	to 379	24
380	to 397	25
398	to 414	26
415	to 432	27
433	to 449	28
450	to 467	29
468	to 485	30
486	to 503	31
504	to 520	32
521	to 538	33
539	to 556	34
557	to 574	35
575	to 592	36
593	to 610	37
611	to 628	38
629	to 646	39
647	to 664	40
665	to 682	41
683	to 700	42
701	to 718	43
719	to 736	44
737	to 754	45
755	to 772	46
773	to 791	47
792	to 809	48
810	to 827	49
828	to 845	50
846	to 864	51
865	to 882	52
883	to 900	53
901	to 918	54
919	to 937	55
938	to 955	56
956	to 973	57
974	to 992	58
993	to 1010	59




Table and figure 2:

Population Standard = 3%  
Acceptance Probability  $\geq 90\%$   
n=sample size, k=maximum number of off-types

n	k
1 to 3	0
4 to 17	1
18 to 37	2
38 to 58	3
59 to 81	4
82 to 105	5
106 to 130	6
131 to 156	7
157 to 182	8
183 to 208	9
209 to 235	10
236 to 262	11
263 to 289	12
290 to 317	13
318 to 345	14
346 to 373	15
374 to 401	16
402 to 429	17
430 to 457	18
458 to 486	19
487 to 515	20
516 to 543	21
544 to 572	22
573 to 601	23
602 to 630	24
631 to 659	25
660 to 689	26
690 to 718	27
719 to 747	28
748 to 777	29
778 to 806	30
807 to 836	31
837 to 865	32
866 to 895	33
896 to 925	34
926 to 955	35
956 to 984	36
985 to 1014	37
1015 to 1044	38
1045 to 1074	39
1075 to 1104	40
1105 to 1134	41
1135 to 1164	42
1165 to 1195	43
1196 to 1225	44
1226 to 1255	45
1256 to 1285	46
1286 to 1315	47
1316 to 1346	48
1347 to 1376	49
1377 to 1406	50
1407 to 1437	51
1438 to 1467	52
1468 to 1498	53
1499 to 1528	54




Table and figure 3:

Population Standard = 2%  
Acceptance Probability  $\geq 90\%$   
n=sample size, k=maximum number of off-types

	n	k
1	to 5	0
6	to 26	1
27	to 55	2
56	to 87	3
88	to 122	4
123	to 158	5
159	to 195	6
196	to 233	7
234	to 272	8
273	to 312	9
313	to 352	10
353	to 393	11
394	to 433	12
434	to 475	13
476	to 516	14
517	to 558	15
559	to 600	16
601	to 643	17
644	to 685	18
686	to 728	19
729	to 771	20
772	to 814	21
815	to 857	22
858	to 901	23
902	to 944	24
945	to 988	25
989	to 1032	26
1033	to 1076	27
1077	to 1120	28
1121	to 1164	29
1165	to 1208	30
1209	to 1252	31
1253	to 1297	32
1298	to 1341	33
1342	to 1386	34
1387	to 1431	35
1432	to 1475	36
1476	to 1520	37
1521	to 1565	38
1566	to 1610	39
1611	to 1655	40
1656	to 1700	41
1701	to 1745	42
1746	to 1790	43
1791	to 1835	44
1836	to 1881	45
1882	to 1926	46
1927	to 1971	47
1972	to 2000	48




Table and figure 4:

Population Standard = 1%  
Acceptance Probability  $\geq 90\%$   
n=sample size, k=maximum number of off-types

	n		k
1	to	10	0
11	to	53	1
54	to	110	2
111	to	175	3
176	to	244	4
245	to	316	5
317	to	390	6
391	to	466	7
467	to	544	8
545	to	623	9
624	to	703	10
704	to	784	11
785	to	866	12
867	to	948	13
949	to	1031	14
1032	to	1115	15
1116	to	1199	16
1200	to	1284	17
1285	to	1369	18
1370	to	1454	19
1455	to	1540	20
1541	to	1626	21
1627	to	1713	22
1714	to	1799	23
1800	to	1887	24
1888	to	1974	25
1975	to	2061	26
2062	to	2149	27
2150	to	2237	28
2238	to	2325	29
2326	to	2414	30
2415	to	2502	31
2503	to	2591	32
2592	to	2680	33
2681	to	2769	34
2770	to	2858	35
2859	to	2948	36
2949	to	3000	37






Table and figure 5:

Population Standard = .5%  
Acceptance Probability  $\geq 90\%$   
n=sample size, k=maximum number of off-types

n	k
1 to 21	0
22 to 106	1
107 to 220	2
221 to 349	3
350 to 487	4
488 to 631	5
632 to 780	6
781 to 932	7
933 to 1087	8
1088 to 1245	9
1246 to 1405	10
1406 to 1567	11
1568 to 1730	12
1731 to 1895	13
1896 to 2061	14
2062 to 2228	15
2229 to 2397	16
2398 to 2566	17
2567 to 2736	18
2737 to 2907	19
2908 to 3000	20




Table and figure 6:

Population Standard = .1%  
Acceptance Probability  $\geq 90\%$   
n=sample size, k=maximum number of off-types

	n	k
1	to 105	0
106	to 532	1
533	to 1102	2
1103	to 1745	3
1746	to 2433	4
2434	to 3000	5




Table and figure 7:

Population Standard = 5%  
Acceptance Probability  $\geq 95\%$   
 $n$ =sample size,  $k$ =maximum number of off-types

	$n$		$k$
1	to	1	0
2	to	7	1
8	to	16	2
17	to	28	3
29	to	40	4
41	to	53	5
54	to	67	6
68	to	81	7
82	to	95	8
96	to	110	9
111	to	125	10
126	to	140	11
141	to	155	12
156	to	171	13
172	to	187	14
188	to	203	15
204	to	219	16
220	to	235	17
236	to	251	18
252	to	268	19
269	to	284	20
285	to	300	21
301	to	317	22
318	to	334	23
335	to	351	24
352	to	367	25
368	to	384	26
385	to	401	27
402	to	418	28
419	to	435	29
436	to	452	30
453	to	469	31
470	to	487	32
488	to	504	33
505	to	521	34
522	to	538	35
539	to	556	36
557	to	573	37
574	to	590	38
591	to	608	39
609	to	625	40
626	to	643	41
644	to	660	42
661	to	678	43
679	to	696	44
697	to	713	45
714	to	731	46
732	to	748	47
749	to	766	48
767	to	784	49
785	to	802	50
803	to	819	51
820	to	837	52
838	to	855	53
856	to	873	54
874	to	891	55
892	to	909	56
910	to	926	57
927	to	944	58
945	to	962	59
963	to	980	60
981	to	998	61




Table and figure 8:

Population Standard = 3%  
Acceptance Probability  $\geq 95\%$   
n=sample size, k=maximum number of off-types

n	k
1	0
2	1
13	2
28	3
47	4
67	5
89	6
111	7
135	8
159	9
183	10
208	11
233	12
259	13
285	14
311	15
338	16
364	17
391	18
418	19
445	20
473	21
500	22
528	23
555	24
583	25
611	26
639	27
667	28
696	29
724	30
752	31
781	32
810	33
838	34
867	35
896	36
925	37
953	38
982	39
1011	40
1041	41
1070	42
1099	43
1128	44
1157	45
1187	46
1216	47
1245	48
1275	49
1304	50
1334	51
1363	52
1393	53
1423	54
1452	55
1482	56
1512	57
1542	58
1571	59
1601	60
1631	61
1661	62
1691	63
1721	64
1751	65
1781	66
1811	67
1841	68
1871	69




Table and figure 9:

Population Standard = 2%  
Acceptance Probability  $\geq 95\%$   
 $n$ =sample size,  $k$ =maximum number of off-types

	$n$	$k$
1	to 2	0
3	to 18	1
19	to 41	2
42	to 69	3
70	to 99	4
100	to 131	5
132	to 165	6
166	to 200	7
201	to 236	8
237	to 273	9
274	to 310	10
311	to 348	11
349	to 386	12
387	to 425	13
426	to 464	14
465	to 504	15
505	to 544	16
545	to 584	17
585	to 624	18
625	to 665	19
666	to 706	20
707	to 747	21
748	to 789	22
790	to 830	23
831	to 872	24
873	to 914	25
915	to 956	26
957	to 998	27
999	to 1040	28
1041	to 1083	29
1084	to 1126	30
1127	to 1168	31
1169	to 1211	32
1212	to 1254	33
1255	to 1297	34
1298	to 1340	35
1341	to 1383	36
1384	to 1427	37
1428	to 1470	38
1471	to 1514	39
1515	to 1557	40
1558	to 1601	41
1602	to 1645	42
1646	to 1689	43
1690	to 1732	44
1733	to 1776	45
1777	to 1820	46
1821	to 1864	47
1865	to 1909	48
1910	to 1953	49
1954	to 1997	50
1998	to 2000	51




Table and figure 10:

Population Standard = 1%  
Acceptance Probability  $\geq 95\%$   
n=sample size, k=maximum number of off-types

n	k
1	0
6	1
36	2
83	3
138	4
199	5
263	6
330	7
400	8
472	9
545	10
619	11
695	12
772	13
849	14
928	15
1007	16
1086	17
1167	18
1247	19
1329	20
1411	21
1493	22
1576	23
1659	24
1742	25
1826	26
1910	27
1994	28
2079	29
2164	30
2249	31
2334	32
2420	33
2506	34
2592	35
2678	36
2764	37
2851	38
2938	39




Table and figure 11:

Population Standard = .5%  
Acceptance Probability  $\geq 95\%$   
n=sample size, k=maximum number of off-types

	n	k
1	to 10	0
11	to 71	1
72	to 164	2
165	to 274	3
275	to 395	4
396	to 523	5
524	to 658	6
659	to 797	7
798	to 940	8
941	to 1086	9
1087	to 1235	10
1236	to 1386	11
1387	to 1540	12
1541	to 1695	13
1696	to 1851	14
1852	to 2009	15
2010	to 2169	16
2170	to 2329	17
2330	to 2491	18
2492	to 2653	19
2654	to 2817	20
2818	to 2981	21
2982	to 3000	22




Table and figure 12: Population Standard = .1%  
Acceptance Probability  $\geq 95\%$   
n=sample size, k=maximum number off-types

	n		k
	1	to	51
	52	to	355
	356	to	818
	819	to	1367
	1368	to	1971
	1972	to	2614
	2615	to	3000
			6






Table and figure 13:

Population Standard = 5%  
Acceptance Probability  $\geq 99\%$   
 $n$ =sample size,  $k$ =maximum number of off-types

	$n$	$k$
1	to 3	1
4	to 9	2
10	to 17	3
18	to 26	4
27	to 37	5
38	to 48	6
49	to 60	7
61	to 72	8
73	to 85	9
86	to 98	10
99	to 111	11
112	to 124	12
125	to 138	13
139	to 152	14
153	to 167	15
168	to 181	16
182	to 196	17
197	to 210	18
211	to 225	19
226	to 240	20
241	to 255	21
256	to 270	22
271	to 286	23
287	to 301	24
302	to 317	25
318	to 332	26
333	to 348	27
349	to 364	28
365	to 380	29
381	to 395	30
396	to 411	31
412	to 427	32
428	to 444	33
445	to 460	34
461	to 476	35
477	to 492	36
493	to 508	37
509	to 525	38
526	to 541	39
542	to 558	40
559	to 574	41
575	to 591	42
592	to 607	43
608	to 624	44
625	to 640	45
641	to 657	46
658	to 674	47
675	to 690	48
691	to 707	49
708	to 724	50
725	to 741	51
742	to 758	52
759	to 775	53
776	to 792	54
793	to 809	55
810	to 826	56
827	to 843	57
844	to 860	58
861	to 877	59
878	to 894	60
895	to 911	61
912	to 928	62
929	to 945	63
946	to 962	64
963	to 979	65
980	to 997	66
998	to 1014	67




Table and figure 14:

Population Standard = 3%

Acceptance Probability  $\geq 99\%$

n=sample size, k=maximum number of off-types

n	k
1 to 5	1
6 to 15	2
16 to 28	3
29 to 44	4
45 to 61	5
62 to 79	6
80 to 98	7
99 to 119	8
120 to 140	9
141 to 161	10
162 to 183	11
184 to 206	12
207 to 229	13
230 to 252	14
253 to 276	15
277 to 300	16
301 to 324	17
325 to 348	18
349 to 373	19
374 to 398	20
399 to 423	21
424 to 448	22
449 to 474	23
475 to 499	24
500 to 525	25
526 to 551	26
552 to 577	27
578 to 603	28
604 to 629	29
630 to 656	30
657 to 682	31
683 to 709	32
710 to 736	33
737 to 763	34
764 to 789	35
790 to 816	36
817 to 844	37
845 to 871	38
872 to 898	39
899 to 925	40
926 to 953	41
954 to 980	42
981 to 1008	43
1009 to 1035	44
1036 to 1063	45
1064 to 1091	46
1092 to 1119	47
1120 to 1146	48
1147 to 1174	49
1175 to 1202	50
1203 to 1230	51
1231 to 1258	52
1259 to 1286	53
1287 to 1315	54
1316 to 1343	55
1344 to 1371	56
1372 to 1399	57
1400 to 1428	58
1429 to 1456	59
1457 to 1484	60
1485 to 1513	61




Table and figure 15:

Population Standard = 2%  
Acceptance Probability  $\geq 99\%$   
n=sample size, k=maximum number of off-types

n	k
1 to 7	1
8 to 22	2
23 to 42	3
43 to 65	4
66 to 90	5
91 to 118	6
119 to 147	7
148 to 177	8
178 to 208	9
209 to 241	10
242 to 274	11
275 to 307	12
308 to 342	13
343 to 377	14
378 to 412	15
413 to 448	16
449 to 484	17
485 to 521	18
522 to 558	19
559 to 595	20
596 to 632	21
633 to 670	22
671 to 708	23
709 to 747	24
748 to 785	25
786 to 824	26
825 to 863	27
864 to 902	28
903 to 942	29
943 to 981	30
982 to 1021	31
1022 to 1061	32
1062 to 1101	33
1102 to 1141	34
1142 to 1182	35
1183 to 1222	36
1223 to 1263	37
1264 to 1303	38
1304 to 1344	39
1345 to 1385	40
1386 to 1426	41
1427 to 1467	42
1468 to 1509	43
1510 to 1550	44
1551 to 1591	45
1592 to 1633	46
1634 to 1675	47
1676 to 1716	48
1717 to 1758	49
1759 to 1800	50
1801 to 1842	51
1843 to 1884	52
1885 to 1926	53
1927 to 1968	54
1969 to 2000	55




Table and figure 16:

Population Standard = 1%  
Acceptance Probability  $\geq 99\%$   
n=sample size, k=maximum number of off-types

n	k
1	0
2	1
16	2
45	3
84	4
130	5
181	6
235	7
293	8
354	9
416	10
480	11
546	12
613	13
682	14
751	15
822	16
894	17
966	18
1039	19
1113	20
1187	21
1262	22
1338	23
1414	24
1490	25
1567	26
1645	27
1723	28
1801	29
1880	30
1959	31
2038	32
2118	33
2198	34
2278	35
2359	36
2440	37
2521	38
2602	39
2684	40
2765	41
2847	42
2930	43




Table and figure 17:

Population Standard = .5%  
Acceptance Probability  $\geq 99\%$   
n=sample size, k=maximum number of off-types

n	k
1	0
3	1
31	2
88	3
166	4
258	5
359	6
468	7
584	8
704	9
829	10
957	11
1089	12
1223	13
1360	14
1499	15
1640	16
1783	17
1927	18
2073	19
2221	20
2370	21
2520	22
2671	23
2823	24
2976	25




Table and figure 18:

Population Standard = .1%  
Acceptance Probability  $\geq 99\%$   
n=sample size, k=maximum number of off-types

n	to	k
1	to 10	0
11	to 148	1
149	to 436	2
437	to 824	3
825	to 1280	4
1281	to 1786	5
1787	to 2332	6
2333	to 2908	7
2909	to 3000	8




Table and figure 19:

Population Standard = 10%  
Acceptance Probability  $\geq 90\%$   
n=sample size, k=maximum number of off-types

n	k
1	0
2	1
6	2
12	3
19	4
26	5
33	6
41	7
48	8
56	9
64	10
72	11
80	12
89	13
97	14
105	15
114	16
122	17
131	18
139	19
148	20
157	21
165	22
174	23
183	24
192	25
200	26





Table and figure 20:

Population Standard = 10%  
Acceptance Probability  $\geq 95\%$   
n=sample size, k=maximum number of off-types


n	k
1	3
4	8
9	14
15	20
21	27
28	34
35	41
42	48
49	56
57	63
64	71
72	79
80	86
87	94
95	102
103	110
111	119
120	127
128	135
136	143
144	152
153	160
161	168
169	177
178	185
186	194
195	200





**Table and figure 21 :** Population Standard = 10%  
Acceptance Probability  $\geq 99\%$   
n=sample size, k=maximum number of off-types

n	to	k
1	to	2
3	to	5
6	to	9
10	to	14
15	to	19
20	to	25
26	to	31
32	to	37
38	to	43
44	to	50
51	to	57
58	to	64
65	to	71
72	to	78
79	to	85
86	to	92
93	to	99
100	to	107
108	to	114
115	to	122
123	to	130
131	to	137
138	to	145
146	to	153
154	to	161
162	to	168
169	to	176
177	to	184
185	to	192
193	to	200



## 2. LSD

to be provided by the TWC (see extract below from TGP/9/1 Draft 5)

[5.2.4.10 The General Introduction clarifies the situation with regard to measured, quantitative characteristics for vegetatively propagated and self-pollinated varieties as follows:

### “5.5.3 Measured Characteristics

The following paragraphs provide guidance on the typical methods for examining distinctness according to the particular features of propagation of the variety:

[...]

#### 5.5.3.1 Self-Pollinated and Vegetatively Propagated Varieties

UPOV has endorsed several statistical methods for the handling of measured quantitative characteristics. One method established for self-pollinated and vegetatively propagated varieties is that varieties can be considered clearly distinguishable if the difference between two varieties equals or exceeds the Least Significant Difference (LSD) at a specified probability level with the same sign over an appropriate period, even if they are described by the same state of expression. This is a relatively simple method but is considered appropriate for self-pollinated and vegetatively propagated varieties because the level of variation within such varieties is relatively low. Further details are provided in document TGP/9, “Examining Distinctness.””

5.2.4.11 Information on the Least Significant Difference (LSD) method is provided in TGP/8 [*cross ref.*].

[TWC Chairperson: to find users and crops in which this approach is used]

## 3. The Combined Over-Years Criterion for Distinctness and Uniformity

### 3.1 The Combined Over-Years Distinctness Criterion (COYD)

#### 3.1.1 Summary

3.1.1.1 Document TGP/9, Section 5.2.4.14 explains that “To assess distinctness for varieties on the basis of a quantitative characteristic it is possible to calculate a minimum distance between varieties such that, when the distance calculated between a pair of varieties is greater than this minimum distance, they may be considered as “distinct” in respect of that characteristic. Amongst the possible ways of establishing minimum distances is the method known as the Combined-Over-Years Distinctness (COYD). The COYD analysis takes into account variations between years. Its main use is for cross-pollinated, including synthetic, varieties but, if desired, it can also be used for self-pollinated and vegetatively propagated varieties in certain circumstances. This method requires the size of the differences to be sufficiently consistent over the years and takes into account the variation between years.” [*cross ref.*].

3.1.1.2 The COYD method involves:

- for each characteristic, taking the variety means from the two or three years of trials for candidates and established varieties and producing over-year means for the varieties;
- calculate a least significant difference (LSD), based on variety-by-years variation, for comparing variety means.
- if the over-years mean difference between two varieties is greater than or equal to the LSD then the varieties are said to be distinct in respect of that characteristic.

3.1.1.3 The main advantages of the COYD method are:

- it combines information from several seasons into a single criterion (the “COYD criterion”) in a simple and straightforward way;
- it ensures that judgements about distinctness will be reproducible in other seasons; in other words, the same genetic material should give similar results, within reasonable limits, from season-to-season;
- the risks of making a wrong judgement about distinctness are constant for all characteristics.

### 3.1.2 Introduction

3.1.2.1 The following sections describe:

- the principles underlying the COYD method;
- UPOV recommendations on the application of COYD to individual species;
- details of ways in which the procedure can be adapted to deal with special circumstances. This includes when there are small numbers of varieties in trial;
- the computer software which is available to apply the procedure.

### 3.1.3 The COYD Method

3.1.3.1 The COYD method aims to establish for each characteristic a minimum difference, or distance, which, if achieved by two varieties in trials over a period of two or three years, would indicate that those varieties are distinct with a specified degree of confidence.

3.1.3.2 The method uses variation in variety expression of a characteristic from year-to-year to establish the minimum distance. Thus, characteristics which show consistency in variety ranking between years will have smaller minimum distances than those with marked changes in ranking.

3.1.3.3 Calculation of the COYD criterion involves analysing the variety-by-year table of means for each characteristic to get an estimate of the varieties-by-years variation, which is used in the next step: to calculate an LSD. Usually data for all candidate and established varieties which appeared in trials over the two or three test years are included in the table, the analysis is by analysis of variance, the varieties-by-years mean square is used as the estimate of the varieties-by-years variation, and the resulting LSD is known as the COYD LSD. However, where there are small numbers of varieties in trial, the approach is different.

3.1.3.4 Where there are small numbers of varieties in trial, the table used to calculate of the COYD criterion is expanded with means from other varieties and earlier years, a different method of analysis is used to get a varieties-by-years mean square to estimate the varieties-by-years variation, and the resulting LSD is known as the Long-Term LSD. This is discussed later.

3.1.3.5 Equation [1]

$$\text{LSD}_p = t_p \times \sqrt{2} \times \text{SE}(\bar{x})$$

where  $\text{SE}(\bar{x})$  is the standard error of a variety's over-year mean calculated as:

$$\text{SE}(\bar{x}) = \sqrt{\frac{\text{varieties - by - years mean square}}{\text{number of test years}}}$$

and  $t_p$  is the value in Student's t table appropriate for a two-tailed test with probability  $p$  and with degrees of freedom associated with the variety-by-years mean square. The probability level  $p$  that is appropriate for individual species is discussed under UPOV RECOMMENDATIONS ON COYD below.

3.1.3.6 An example of the application of COYD to a small data set is given in Figure 1. Statistical details of the method are in Part I: Section 2.1.8 [*cross ref.*]. Further information about the COYD criterion can be found in Patterson and Weatherup (1984).

### 3.1.4 Use of COYD

3.1.4.1 COYD is an appropriate method for assessing the distinctness of varieties where:

- the characteristic is quantitative;
- there are some differences between plants (or plots) of a variety.
- observations are made on a plant (or plot) basis over two or more years;

3.1.4.2 A pair of varieties is considered to be distinct if their over-years means differ by at least the COYD LSD in one or more characteristics.

3.1.4.3 The UPOV recommended probability level  $p$  for the  $t_p$  value used to calculate the COYD LSD differs depending on the crop and for some crops depends on whether the test is over two or three years. The testing schemes that usually arise in distinctness testing are described in [.....] [*cross ref.*].

### 3.1.5 Adapting COYD to special circumstances

#### 3.1.5.1 Differences between years in the range of expression of a characteristic.

Occasionally, marked differences between years in the range of expression of a characteristic can occur. For example, in a late spring, the heading dates of grass varieties can converge. To take account of this effect it is possible to fit extra terms, one for each year, in the analysis of variance. Each term represents the linear regression of the observations for the year against the variety means over all years. The method is known as modified joint regression analysis (MJRA) and is recommended in situations where there is a statistically significant ( $p \leq 1\%$ ) contribution from the regression terms in the analysis of variance. Statistical details, and a computer program to implement the procedure, are described in Part II Sections 2.1.8 and 2.1.9 [*cross ref.*].

#### 3.1.5.2 Small numbers of varieties in trials: Long-Term COYD

3.1.5.2.1 It is recommended that there should be at least 20 degrees of freedom for the varieties-by-years mean square in the COYD analysis of variance. This is in order to ensure that the varieties-by-years mean square is based on sufficient data to be a reliable estimate of the varieties-by-years variation for the LSD. Twenty degrees of freedom corresponds to 11 varieties common in three years of trials, or 21 varieties common in two years. Trials with fewer varieties in common over years are considered to have small numbers of varieties in trial.

3.1.5.2.2 In such trials the variety-by-year tables of means can be expanded to include means for earlier years, and if necessary, other established varieties. As not all varieties are present in all years, the resulting tables of variety-by-year means are not balanced. Consequently, each table is analysed by the least squares method of fitted constants (FITCON) or by REML, which produces an alternative varieties-by-years mean square as a long-term estimate of variety-by-years variation. This estimate has more degrees of freedom as it is based on more years and varieties.

$$\text{degrees of freedom} = \left( \begin{array}{c} \text{No. values in expanded} \\ \text{variety - by - year table} \end{array} \right) - (\text{No. varieties}) - (\text{No. years}) + 1$$

3.1.5.2.3 The alternative varieties-by-years mean square is used in equation [1] above to calculate an LSD. This LSD is known as a “Long-Term LSD” to distinguish it from COYD LSD based on just the test years and varieties. The Long-Term LSD is used in the same way as the COYD LSD is used to assess the distinctness of varieties by comparing their over-year (the test years) means. The act of comparing the means of varieties using a “Long-Term LSD” is known as “Long-Term COYD”.

3.1.5.2.4 Long-Term COYD should only be applied to those characteristics lacking the recommended minimum degrees of freedom. However, when there is evidence that a characteristic’s LSD fluctuates markedly across years, it may be necessary to base the LSD for that characteristic on the current two or three-years of data, even though it has few degrees of freedom.

3.1.5.2.5 Figure 2 gives an example of the application of Long-Term COYD to the Italian ryegrass characteristic “Growth habit in spring”. A flow diagram of the stages and DUST

modules used to produce Long-Term LSD's and perform Long-Term COYD is given in Figure B2 in Part II: Section 2.1.9. [*cross ref.*]

### 3.1.5.3 Marked year-to-year changes in an individual variety's characteristic

Occasionally, a pair of varieties may be declared distinct on the basis of a t-test which is significant solely due to a very large difference between the varieties in a single year. To monitor such situations a check statistic is calculated, called  $F_3$ , which is the variety-by-years mean square for the particular variety pair expressed as a ratio of the overall variety-by-years mean square. This statistic should be compared with F-distribution tables with 1 and  $g$ , or 2 and  $g$ , degrees of freedom, for tests with two or three years of data respectively where  $g$  is the degrees of freedom for the variety-by-years mean square. If the calculated  $F_3$  value exceeds the tabulated F value at the 1% level then an explanation for the unusual result should be sought before making a decision on distinctness.

### 3.1.6 Implementing COYD

The COYD method can be applied using the DUST package for the statistical analysis of DUS data, which is available from Dr. Sally Watson, Biometrics Division, Department of Agriculture for Northern Ireland (DANI), Newforge Lane, Belfast BT9 5PX, United Kingdom. Sample outputs are given in Part II: Section 2.1.9. [*cross ref.*]

### 3.1.7 References

- DIGBY, P.G.N. (1979). Modified joint regression analysis for incomplete variety x environment data. J. Agric. Sci. Camb. 93, 81-86.
- PATTERSON, H.D. & WEATHERUP, S.T.C. (1984). Statistical criteria for distinctness between varieties of herbage crops. J. Agric. Sci. Camb. 102, 59-68.
- TALBOT, M. (1990). Statistical aspects of minimum distances between varieties. UPOV TWC Paper TWC/VIII/9, UPOV, Geneva.

**Figure 1: Illustrating the calculation of the COYD criterion**

**Characteristic: Days to ear emergence in perennial ryegrass varieties**

Varieties	Years			Over Year Means	<i>Difference (Varieties compared to C2)</i>	
	1	2	3			
<i>Reference</i>		Means				
R1	38	41	35	38	35	<i>D</i>
R2	63	68	61	64	9	<i>D</i>
R3	69	71	64	68	5	<i>D</i>
R4	71	75	67	71	2	
R5	69	78	69	72	1	
R6	74	77	71	74	-1	
R7	76	79	70	75	-2	
R8	75	80	73	76	-3	
R9	78	81	75	78	-5	<i>D</i>
R10	79	80	75	78	-5	<i>D</i>
R11	76	85	79	80	-7	<i>D</i>
<i>Candidate</i>						
C1	52	56	48	52	21	<i>D</i>
C2	72	79	68	73	0	-
C3	85	88	85	86	-13	<i>D</i>

**ANALYSIS OF VARIANCE**

Source	df	Mean square
Years	2	174.93
Variety	13	452.59
Variety-by-years	26	2.54

$$LSD_p = t_p * \sqrt{2} * SE(\bar{X})$$

$$LSD_{0.01} = 2.779 * 1.414 * \sqrt{(2.54/3)} = 3.6$$

Where  $t_p$  is taken from Student's t table with  $p = 0.01$  (two-tailed) and 26 degrees of freedom.

To assess the distinctness of a candidate, the difference in the means between the candidate and all other varieties is computed. In practice a column of differences is calculated for each candidate. In this case, varieties with mean differences greater than, or equal to, 3.6 are regarded as distinct (marked *D* above).

**Figure 2: Illustrating the application of Long-Term COYD**

**Characteristic: Growth habit in spring in italian ryegrass varieties**

Varieties	1	2	Years			Mean over test years	<i>Difference (Varieties compared to C2)</i>	
			3*	4*	5*			
<i>Reference</i>			Means					
R1	43	42	41	44				
R2		39	45					
R3	43	38	41	45	40	42	6	<i>D</i>
R4	44	40	42	48	44	44.7	3.3	<i>D</i>
R5	46	43	48	49	45	47.3	0.7	
R6	51	48	52	53	51	52	-4	<i>D</i>
<i>Candidate</i>								
C1			43	45	44	44	4	<i>D</i>
C2			49	50	45	48	0	
C3			48	53	47	49.3	-1.3	

\* indicates a test year

The aim is to assess the distinctness of the candidate varieties C1, C2 & C3 grown in the test years 3, 4 & 5.

The trial has a small number of varieties in trial because there are just seven varieties in common over the test years 3, 4 & 5 (data marked by a black border).

FITCON analysis of the variety-by-years table of means expanded to nine varieties in five years gives: varieties-by-years mean square = 1.924, on 22 degrees of freedom

$$\text{Long-term LSD}_p = t_p * \sqrt{2} * \text{SE}(\bar{X})$$

$$\text{Long-term LSD}_{0.01} = 2.819 * 1.414 * \sqrt{(1.924/3)} = 3.19$$

Where  $t_p$  is taken from Student's t table with  $p = 0.01$  (two-tailed) and 22 degrees of freedom

To assess the distinctness of a candidate, the difference in the means between the candidate and all other varieties is computed. In practice a column of differences is calculated for each candidate. In the case of variety C2, varieties with mean differences greater than, or equal to 3.19 are regarded as distinct (marked *D* above).



### 3.1.8 COYD statistical methods

#### 3.1.8.1 Analysis of variance

The standard errors used in the COYD criterion are based on an analysis of variance of the variety-by-years table of a characteristic's means. For  $m$  years and  $n$  varieties this analysis of variance breaks down the available degrees of freedom as follows:

Source	Df
Years	$m-1$
Varieties	$n-1$
Varieties-by-years	$(m-1)(n-1)$

#### 3.1.8.2 Modified joint regression analysis (MJRA)

3.1.8.2.1 As noted above, the COYD criterion bases the SE of a variety mean on the varieties-by-years variation as estimated by the varieties-by-years mean square. Systematic variation can sometimes be identified as well as non-systematic variation. This systematic effect causes the occurrence of different slopes of the regression lines relating variety means in individual years to the average variety means over all years. Such an effect can be noted for the heading date characteristic in a year with a late spring: the range of heading dates can be compressed compared with the normal. This leads to a reduction in the slope of the regression line for variety means in that year relative to average variety means. Non-systematic variation is represented by the variation about these regression lines. Where only non-systematic varieties-by-years variation occurs, the slope of the regression lines have the constant value 1.0 in all years. However, when systematic variation is present, slopes differing from 1.0 occur but with an average of 1.0. When MJRA is used, the SE of a variety mean is based on the non-systematic part of the varieties-by-year variation.

3.1.8.2.2 The difference between the total varieties-by-years variation and the varieties-by-years variation adjusted by MJRA is illustrated in Figure B1, where variety means in each of three years are plotted against average variety means over all years. The variation about three parallel lines fitted to the data, one for each year, provides the total varieties-by-years variation as used in the COYD criterion described above. These regression lines have the common slope 1.0. This variation may be reduced by fitting separate regression lines to the data, one for each year. The resultant residual variation about the individual regression lines provides the MJRA-adjusted varieties-by-years mean square, on which the SE for a variety mean may be based. It can be seen that the MJRA adjustment is only effective where the slopes of the variety regression lines differ between years, such as can occur in heading dates.

3.1.8.2.3 The use of this technique in assessing distinctness has been included as an option in the computer program which applies the COYD criterion in the DUST package. It is recommended that it is only applied where the slopes of the variety regression lines are significantly different between years at the 1% significance level. This level can be specified in the computer program.

3.1.8.2.4 To calculate the adjusted variety means and regression line slopes the following model is assumed.

$$y_{ij} = u_j + b_j v_i + e_{ij}$$

where  $y_{ij}$  is the value for the  $i^{\text{th}}$  variety in the  $j^{\text{th}}$  year.

$u_j$  is the mean of year  $j$  ( $j = 1, \dots, m$ )

$b_j$  is the regression slope for year  $j$

$v_i$  is the effect of variety  $i$  ( $i = 1, \dots, n$ )

$e_{ij}$  is an error term.

3.1.8.2.5 From equations (6) and (7) of Digby (1979), with the meaning of years and varieties reversed, the following equations relating these terms are derived for the situation where data are complete:

$$\sum_{i=1}^n v_i y_{ij} = b_j \sum_{i=1}^n v_i^2$$

$$\sum_{j=1}^m b_j y_{ij} = v_i \sum_{j=1}^m b_j^2$$

3.1.8.2.6 These equations are solved iteratively. All  $b_j$  values are taken to be 1.0 as a starting point in order to provide values for the  $v_i$ 's. The MJRA residual sum of squares is then calculated as:

$$\sum_{j=1}^m \sum_{i=1}^n (y_{ij} - u_j - b_j v_i)^2$$

3.1.8.2.7 This sum of squares is used to calculate the MJRA-adjusted varieties-by-years mean square on  $(m-1)(n-1) - m + 1$  degrees of freedom.

### 3.1.8.3 Comparison of COYD with other criteria

3.1.8.3.1 It can be shown that, for a three-year test, the COYD criterion applied at the 1% probability level is of approximately the same stringency as the 2x1% criterion for a characteristic where the square root of the ratio of the variety-by-years mean square to the variety-by-replicates-within-trials mean square ( $\lambda$ ) has a value of 1.7. The COYD criterion applied at the 1% level is less stringent than the 2x1% criterion if  $\lambda < 1.7$ , and more stringent if  $\lambda > 1.7$ .

## 3.1.9 COYD software

3.1.9.1 An example of the output from the computer program in the DUST package which applies the COYD criterion is given in Tables B 1 to 3. It is taken from a perennial

ryegrass (diploid) trial involving 40 reference varieties (R1 to R40) and 9 candidate varieties (C1 to C9) in 6 replicates on which 8 characteristics were measured over the years 1988, 1989 and 1990.

3.1.9.2 Each of the 8 characteristics is analysed by analysis of variance. As this analysis is of the variety-by-year-by-replicate data, the mean squares are 6 (= number of replicates) times the size of the mean squares of the analysis of variance of the variety-by-year data referred to in the main body of this paper. The results are given in Table B 1. Apart from the over-year variety means there are also presented:

YEAR MS:	the mean square term for years
VARIETY MS:	the mean square term for varieties
VAR.YEAR MS:	the mean square for varieties-by-years interaction
F1 RATIO:	ratio of VARIETY MS to VAR.YEAR MS (a measure of the discriminating power of the characteristic - large values indicate high discriminating power)
VAR.REP MS:	average of the variety-by-replicate mean squares from each year
LAMBDA VALUE ( $\lambda$ ):	square root of the ratio of VAR.YEAR MS to VAR.REP MS
BETWEEN SE:	standard error of variety means over trials on a plot basis i.e. the square root of the VAR.YEAR MS divided by 18 (3 years x 6 replicates)
WITHIN SE:	the standard error of variety means within a trial on a plot basis i.e. the square root of the VAR.REP MS divided by 18
DF:	the degrees of freedom for varieties-by-years
MJRA SLOPE:	the slope of the regression of a single year's variety means on the means over the three years
REGR F VALUE:	the mean square due to MJRA regression as a ratio of the mean square about regression
REGR PROB:	the statistical significance of the REGR F VALUE
TEST:	indicates whether MJRA adjustment was applied (REG) or not (COY).

3.1.9.3 Each candidate variety is compared with every other candidate and reference variety. The mean differences between pairs of varieties are compared with the LSD for the characteristic. The results for the variety pair R1 and C1 are given in Table B 2. The individual within year t-values are listed to provide information on the separate years. Varieties R1 and C1 are considered distinct since, for at least one characteristic, a mean difference is COYD significant at the 1% level. If the  $F_3$  ratio for characteristic 8 had been significant at the 1% level rather than the 5% level, the data for characteristic 8 would have been investigated, and because the differences in the three years are not all in the same direction, the COYD significance for characteristic 8 would not have counted towards distinctness.

3.1.9.4 The outcome in terms of the tests for distinctness of each candidate variety from all other varieties is given in Table B 3, where D indicates "distinct" and ND denotes "not distinct."

**Table B 1: An example of the output from the COYD program showing variety means and analysis of variance of characteristics**

PRG (DIPLOID) EARLY N.I. UPOV 1988-90

	VARIETY MEANS OVER YEARS								
	5	60	8	10	11	14	15	24	
	SP.HT	NSPHT	DEEE	H.EE	WEE	LFL	WFL	LEAR	
1 R	45.27	34.60	67.87	45.20	70.05	20.39	6.85	24.54	
2 R2	42.63	31.84	73.85	41.96	74.98	19.68	6.67	24.44	
3 R3	41.57	27.40	38.47	27.14	57.60	17.12	6.85	22.57	
4 R4	33.35	21.80	77.78	30.77	78.04	18.25	6.40	21.09	
5 R5	37.81	25.86	50.14	27.24	62.64	16.41	6.41	16.97	
6 R6	33.90	21.07	78.73	32.84	79.15	19.44	6.46	21.79	
7 R7	41.30	31.37	73.19	41.35	71.87	20.98	6.92	24.31	
8 R8	24.48	19.94	74.83	32.10	62.38	15.22	6.36	19.46	
9 R9	46.68	36.69	63.99	44.84	68.62	18.11	7.02	22.58	
10 R10	25.60	20.96	75.64	32.31	57.20	14.68	5.51	20.13	
11 R11	41.70	30.31	74.60	40.17	76.15	19.45	6.79	22.72	
12 R12	28.95	21.56	66.12	27.96	59.56	14.83	5.53	20.55	
13 R13	40.67	29.47	70.63	36.81	74.12	19.97	7.04	24.05	
14 R14	26.68	20.53	75.84	34.14	63.29	15.21	6.37	20.37	
15 R15	26.78	20.18	75.54	30.39	66.41	16.34	6.01	20.94	
16 R16	42.44	27.01	59.03	30.39	72.71	17.29	6.47	22.48	
17 R17	27.94	21.58	76.13	32.53	68.37	16.72	6.11	22.03	
18 R18	41.34	30.85	69.80	37.28	69.52	20.68	7.09	25.40	
19 R19	33.54	23.43	73.65	30.35	75.54	18.97	6.37	22.43	
20 R20	44.14	34.48	68.74	42.60	64.17	18.63	6.56	22.02	
21 R21	27.77	21.53	80.52	31.59	69.41	16.81	5.81	22.35	
22 R22	38.90	27.83	75.68	43.25	75.08	19.63	7.46	23.99	
23 R23	42.43	31.80	72.40	42.07	74.77	20.99	6.78	23.57	
24 R24	38.50	27.73	73.19	37.12	75.76	19.28	6.91	22.77	
25 R25	43.84	29.60	68.82	39.79	74.83	20.63	7.08	22.65	
26 R26	49.48	36.53	63.45	42.01	70.46	22.14	7.84	25.91	
27 R27	25.61	19.25	78.78	29.81	56.81	15.81	5.07	18.94	
28 R28	26.70	20.31	79.41	32.75	66.54	16.92	6.00	21.91	
29 R29	27.90	20.94	72.66	29.85	67.14	16.85	6.28	21.79	
30 R30	43.07	30.34	70.53	40.51	73.23	19.49	7.28	23.70	
31 R31	38.18	25.47	74.23	36.88	80.23	20.40	7.09	25.21	
32 R32	35.15	27.56	71.49	37.26	63.10	18.18	6.80	23.13	
33 R33	42.71	31.09	67.58	39.14	70.36	19.85	7.12	23.35	
34 R34	23.14	18.05	72.09	24.29	59.37	13.98	5.63	18.91	
35 R35	32.75	25.41	77.22	38.90	67.07	17.16	6.42	21.49	
36 R36	41.71	31.94	77.98	44.33	73.00	19.72	7.09	23.45	
37 R37	44.06	32.99	74.38	45.77	71.59	20.88	7.40	24.06	
38 R38	42.65	32.97	74.76	44.42	74.13	20.29	7.38	24.32	
39 R39	28.79	22.41	76.83	35.91	64.52	16.85	6.34	22.24	
40 R40	44.31	31.38	72.24	43.83	74.73	21.53	7.60	25.46	
41 C1	42.42	31.68	64.03	40.22	67.02	20.73	6.90	26.16	
42 C2	41.77	32.35	86.11	46.03	75.35	20.40	6.96	22.99	
43 C3	41.94	31.09	82.04	43.17	74.04	19.06	6.26	23.44	
44 C4	39.03	28.71	78.63	45.97	70.49	21.27	6.67	23.37	
45 C5	43.97	30.95	72.99	39.14	77.89	19.88	6.68	25.44	
46 C6	37.56	27.14	83.29	39.16	81.18	19.47	6.97	25.25	
47 C7	38.41	28.58	83.90	42.53	76.44	19.28	6.00	23.47	
48 C8	40.08	27.25	83.50	43.33	80.16	22.77	7.92	26.81	
49 C9	46.77	34.87	51.89	37.68	61.16	19.25	6.92	24.82	
YEAR MS	1279.09	3398.82	3026.80	2278.15	8449.20	672.15	3.36	51.32	
VARIETY MS	909.21	476.72	1376.10	635.27	762.41	80.21	6.44	74.17	
VAR.YEAR MS	23.16	18.86	14.12	23.16	46.58	4.76	0.28	2.73	
F1 RATIO	39.26	25.27	97.43	27.43	16.37	16.84	22.83	27.16	
VAR.REP MS	8.83	8.19	4.59	11.95	23.23	1.52	0.15	1.70	
LAMBDA VALUE	1.62	1.52	1.75	1.39	1.42	1.77	1.37	1.27	
BETWEEN SE	1.13	1.02	0.89	1.13	1.61	0.51	0.13	0.39	
WITHIN SE	0.70	0.67	0.50	0.81	1.14	0.29	0.09	0.31	
DF	96	94	96	96	96	96	96	96	
MJRA SLOPE 88	0.90	0.86	0.99	0.91	0.99	1.09	0.97	0.95	
MJRA SLOPE 89	1.05	1.08	1.01	0.99	1.06	0.97	1.02	0.98	
MJRA SLOPE 90	1.05	1.06	1.00	1.10	0.95	0.94	1.01	1.07	
REGR F VAL	4.66	6.17	0.06	4.48	0.76	1.62	0.29	1.91	
REGR PROB	1.17	0.30	93.82	1.39	47.08	20.27	74.68	15.38	
TEST	COY	REG	COY	COY	COY	COY	COY	COY	

**Table B 2: An example of the output from the COYD program showing a comparison of varieties R1 and C1**

PRG (DIPLOID) EARLY N.I. UPOV 1988-90

41 C1 VERSUS 1 R1

\*\*\* USING REGR WHERE SIG \*\*\*

(T VALUES + VE IF 41 C1 > 1 R1)

	SIG LEVELS				COYD			T VALUES				F3
	YEARS				T	PROB%	SIG	YEARS			TSCORE	
	88	89	90					88	89	90		
5 SP.HGHT	-	-	-1	ND	-1.78	7.88	NS	-1.05	-1.34	-2.64	-2.64	0.23 NS
60 NATSPHT	-	-1	-	ND	-2.02	4.61	*	-1.58	-2.61	-1.17	-2.61	0.22 NS
8 DATEEE	-1	-1	+	D	-3.06	0.29	**	-4.14	-6.33	0.80	-6.74	3.99 *
10 HGHT.EE	-1	-1	-5	D	-3.11	0.25	**	-2.79	-2.69	-2.06	-7.55	0.06 NS
11 WIDTHEE	-	-	-	ND	-1.33	18.58	NS	-1.47	-1.80	-0.21	0.00	0.32 NS
14 LGTHFL	+	+	-	ND	0.47	63.61	NS	0.17	1.83	-0.67	0.00	0.56 NS
15 WIDTHFL	+	-	+	ND	0.27	78.83	NS	0.31	-0.41	0.67	0.00	0.17 NS
24 EARLGTH	5	1	+	ND	2.93	0.42	**	2.10	3.33	1.01	5.43	0.84 NS

Notes

1. The three “COYD” columns headed, T PROB% and SIG give the COYD t value, its significance probability and significance level. The t value is the test statistic formed by dividing the mean difference between two varieties by the standard error of that difference. The t value can be tested for significance by comparing it with appropriate values from Students t-table. Calculating and testing a t value in this manner is equivalent to deriving an LSD and checking to see if the mean difference between the two varieties is greater than the LSD.
2. The two right-hand “F3” columns give the F<sub>3</sub> variance ratio statistic and its significance level. The F<sub>3</sub> statistic is defined in Part II, Section 2.1.5.2 [cross ref.].
3. The sections in boxes refer to earlier distinctness criteria. The three “T VALUES, YEARS” columns headed 88, 89 and 90 are the individual within year t-test values (the Student’s two-tailed t test of the variety means with standard errors estimated using the plot residual mean square), and the three “SIG LEVELS, YEARS” columns headed 88, 89 and 90 give their direction and significance levels. The column containing D and ND gives the distinctness status of the two varieties by the 2 x 1% criterion described in Section 5.2.4.18 of document TGP/8 [cross ref.]. The column headed T SCORE gives the obsolete T Score statistic and should be ignored.

**Table B 3: An example of the output from the COYD program showing the distinctness status of the candidate varieties**


PRG (DIPLOID) EARLY N.I. UPOV 1988-90

SUMMARY FOR COYD AT 1.0% LEVEL


\*\*\* USING REGR ADJ WHEN SIG \*\*\*

CANDIDATE VARIETIES	C1	C2	C3	C4	C5	C6	C7	C8	C9
1 R1	D	D	D	D	D	D	D	D	D
2 R2	D	D	D	D	ND	D	D	D	D
3 R3	D	D	D	D	D	D	D	D	D
4 R4	D	D	D	D	D	D	D	D	D
5 R5	D	D	D	D	D	D	D	D	D
6 R6	D	D	D	D	D	D	D	D	D
7 R7	D	D	D	D	D	D	D	D	D
8 R8	D	D	D	D	D	D	D	D	D
9 R9	D	D	D	D	D	D	D	D	D
10 R10	D	D	D	D	D	D	D	D	D
11 R11	D	D	D	D	D	D	D	D	D
12 R1	D	D	D	D	D	D	D	D	D
13 R13	D	D	D	D	ND	D	D	D	D
14 R14	D	D	D	D	D	D	D	D	D
15 R15	D	D	D	D	D	D	D	D	D
16 R16	D	D	D	D	D	D	D	D	D
17 R17	D	D	D	D	D	D	D	D	D
18 R18	D	D	D	D	D	D	D	D	D
19 R19	D	D	D	D	D	D	D	D	D
20 R20	D	D	D	D	D	D	D	D	D
21 R21	D	D	D	D	D	D	D	D	D
22 R22	D	D	D	D	D	D	D	D	D
23 R23	D	D	D	D	D	D	D	D	D
24 R24	D	D	D	D	D	D	D	D	D
25 R25	D	D	D	D	D	D	D	D	D
26 R26	D	D	D	D	D	D	D	D	D
27 R27	D	D	D	D	D	D	D	D	D
28 R28	D	D	D	D	D	D	D	D	D
29 R29	D	D	D	D	D	D	D	D	D
30 R30	D	D	D	D	D	D	D	D	D
31 R31	D	D	D	D	D	D	D	D	D
32 R32	D	D	D	D	D	D	D	D	D
33 R33	D	D	D	D	D	D	D	D	D
34 R34	D	D	D	D	D	D	D	D	D
35 R35	D	D	D	D	D	D	D	D	D
36 R36	D	D	D	ND	D	D	D	D	D
37 R37	D	D	D	D	D	D	D	D	D
38 R38	D	D	D	D	D	D	D	D	D
39 R39	D	D	D	D	D	D	D	D	D
40 R40	D	D	D	D	D	D	D	D	D
41 C1	-	D	D	D	D	D	D	D	D
42 C2	D	-	D	D	D	D	D	D	D
43 C3	D	D	-	D	D	D	ND	D	D
44 C4	D	D	D	-	D	D	D	D	D
45 C5	D	D	D	D	-	D	D	D	D
46 C6	D	D	D	D	D	-	D	D	D
47 C7	D	D	ND	D	D	D	-	D	D
48 C8	D	D	D	D	D	D	D	-	D
49 C9	D	D	D	D	D	D	D	D	-
NO OF ND VARS	0	0	1	1	2	0	1	0	0
DISTINCTNESS	D	D	ND	ND	ND	D	ND	D	D
CANDIDATE VAR	C1	C2	C3	C4	C5	C6	C7	C8	C9

Figure B1. Heading date yearly variety means against over-year variety means



**Figure B2. Flow Diagram of the stages and DUST modules used to produce long-term LSD's and perform long-term COYD**





## 3.2 The Combined-Over-Years Uniformity Criterion (COYU)

### 3.2.1 Summary

3.2.1.1 TGP/10 explains that when the off-type approach for the assessment of uniformity is not appropriate for the assessment of uniformity, the standard deviation approach can be used. It further states the following with respect to determination of the acceptable level of variation.

#### 5.2 Determining the acceptable level of variation

5.2.1.1 The comparison between a candidate variety and comparable varieties is carried out on the basis of standard deviations, calculated from individual plant observations. UPOV has proposed several statistical methods for dealing with uniformity in measured quantitative characteristics. One method, which takes into account variations between years, is the Combined Over Years Uniformity (COYU) method. The comparison between a candidate variety and comparable varieties is carried out on the basis of standard deviations, calculated from individual plant observations. This COYU procedure calculates a tolerance limit on the basis of comparable varieties already known i.e. uniformity is assessed using a relative tolerance limit based on varieties within the same trial with comparable expression of characteristics.

3.2.1.2 Uniformity is often related to the expression of a characteristic. For example, in some species, varieties with larger plants tend to be less uniform in size than those with smaller plants. If the same standard is applied to all varieties then it is possible that some may have to meet very strict criteria while others face standards that are easy to satisfy. COYU addresses this problem by adjusting for any relationship that exists between uniformity, as measured by the plant-to-plant SD, and the expression of the characteristic, as measured by the variety mean, before setting a standard.

#### **[TWC Chairperson: Explanation of “reference” varieties to be developed]**

3.2.1.3 The technique involves ranking reference and candidate varieties by the mean value of the characteristic. Each variety's SD is taken and the mean SD of the most similar varieties is subtracted. This procedure gives, for each variety, a measure of its uniformity expressed relative to that of comparable varieties.

3.2.1.4 The results for each year are combined in a variety-by-years table of adjusted SDs and analysis of variance is applied. The mean adjusted SD for the candidate is compared with the mean for the reference varieties using a standard t-test.

3.2.1.5 COYU, in effect, compares the uniformity of a candidate with that of the reference varieties most similar in relation to the characteristic being assessed. The main advantages of COYU are that all varieties can be compared on the same basis and that information from several years of testing may be combined into a single criterion.

### 3.2.2 Introduction

3.2.2.1 Uniformity is sometimes assessed by measuring individual characteristics and calculating the standard deviation (SD) of the measurements on individual plants within a plot. The SDs are averaged over all replicates to provide a single measure of uniformity for each variety in a trial.

3.2.2.2 This section outlines a procedure known as the combined-over-years uniformity (COYU) criterion. COYU assesses the uniformity of a variety relative to reference varieties based on SDs from trials over several years. A feature of the method is that it takes account of possible relationships between the expression of a characteristic and uniformity.

3.2.2.3 This section describes:

- The principles underlying the COYU method.
- UPOV recommendations on the application of COYU to individual species.
- Mathematical details of the method with an example of its application.
- The computer software that is available to apply the procedure.

### 3.2.3 The COYU Criterion

3.2.3.1 The application of the COYU criterion involves a number of steps as listed below. These are applied to each characteristic in turn. Details are given under Part II: Section 2.2.4 [*cross ref.*] below.

- Calculation of within-plot SDs for each variety in each year.
- Transformation of SDs by adding 1 and converting to natural logarithms.
- Estimation of the relationship between the SD and mean in each year. The method used is based on moving averages of the log SDs of reference varieties ordered by their means.
- Adjustments of log SDs of candidate and reference varieties based on the estimated relationships between SD and mean in each year.
- Averaging of adjusted log SDs over years.
- Calculation of the maximum allowable SD (the uniformity criterion). This uses an estimate of the variability in the uniformity of reference varieties derived from analysis of variance of the variety-by-year table of adjusted log SDs.
- Comparison of the adjusted log SDs of candidate varieties with the maximum allowable SD.

3.2.3.2 The advantages of the COYU criterion are:

- It provides a method for assessing uniformity that is largely independent of the varieties that are under test.
- The method combines information from several trials to form a single criterion for uniformity.

- Decisions based on the method are likely to be stable over time.
- The statistical model on which it is based reflects the main sources of variation that influence uniformity.
- Standards are based on the uniformity of reference varieties.

### 3.2.4 Recommendations on COYU

#### 3.2.4.1 COYU is recommended for use in assessing the uniformity of varieties

- For quantitative characteristics.
- When observations are made on a plant basis over two or more years.
- When there are some differences between plants of a variety, representing quantitative variation rather than presence of off-types.

3.2.4.2 A variety is considered to be uniform for a characteristic if its mean adjusted log SD does not exceed the uniformity criterion.

3.2.4.3 The probability level “p” used to determine the uniformity criterion depends on the crop. Recommended probability levels are given in [.....] [*cross ref.*]

3.2.4.4 The uniformity test may be made over two or three years. If the test is normally applied over three years, it is possible to choose to make an early acceptance or rejection of a variety using an appropriate selection of probability values.

3.2.4.5 It is recommended that there should be at least 20 degrees of freedom for the estimate of variance for the reference varieties formed in the COYU analysis. This corresponds to 11 reference varieties for a COYU test based on two years of trials and 8 reference varieties for three years. In some situations, there may not be enough reference varieties to give the recommended minimum degrees of freedom. Advice is being developed for such cases.

### 3.2.5 Mathematical details

Step 1: Derivation of the within-plot standard deviation

3.2.5.1 Within-plot standard deviations for each variety in each year are calculated by averaging the plot between-plant standard deviations,  $SD_j$ , over replicates:

$$SD_j = \sqrt{\frac{\sum_{i=1}^n (y_{ij} - y_j)^2}{(n-1)}}$$

$$SD = \frac{\sum_{j=1}^r SD_j}{r}$$

where  $y_{ij}$  is the observation on the  $i^{\text{th}}$  plant in the  $j^{\text{th}}$  plot,  $y_j$  is the mean of the observations from the  $j^{\text{th}}$  plot,  $n$  is the number of plants measured in each plot and  $r$  is the number of replicates.

Step 2: Transformation of the SDs


3.2.5.2 Transformation of SDs by adding 1 and converting to natural logarithms. The purpose of this transformation is to make the SDs more amenable to statistical analysis.

Step 3: Estimation of the relationship between the SD and mean in each year

3.2.5.3 For each year separately, the form of the average relationship between SD and characteristic mean is estimated for the reference varieties. The method of estimation is a 9-point moving average. The log SDs (the Y variate) and the means (the X variate) for each variety are first ranked according to the values of the mean. For each point  $(X_i, Y_i)$  take the trend value  $T_i$  to be the mean of the values  $Y_{i-4}, Y_{i-3}, \dots, Y_{i+4}$  where  $i$  represents the rank of the X value and  $Y_i$  is the corresponding Y value. For X values ranked 1<sup>st</sup> and 2<sup>nd</sup> the trend value is taken to be the mean of the first three values. In the case of the X value ranked 3<sup>rd</sup> the mean of the first five values are taken and for the X value ranked 4<sup>th</sup> the mean of the first seven values are used. A similar procedure operates for the four highest-ranked X values.

3.2.5.4 A simple example in Figure 1 illustrates this procedure for 16 varieties. The points marked "O" in Figure 1a represent the log SDs and the corresponding means of 16 varieties. The points marked "X" are the 9-point moving-averages, which are calculated by taking, for each variety, the average of the log SDs of the variety and the four varieties on either side. At the extremities the moving average is based on the mean of 3, 5, or 7 values.

**Figure 1: Association between SD and mean – days to ear emergence in cocksfoot varieties** (symbol O is for observed SD, symbol X is for moving average SD)



Step 4: Adjustment of transformed SD values based on estimated SD-mean relationship


3.2.5.5 Once the trend values for the reference varieties have been determined, the trend values for candidates are estimated using linear interpolation between the trend values of the nearest two reference varieties as defined by their means for the characteristic. Thus if the trend values for the two reference varieties on either side of the candidate are  $T_i$  and  $T_{i+1}$  and the observed value for the candidate is  $X_c$ , where  $X_i \leq X_c \leq X_{i+1}$ , then the trend value  $T_c$  for the candidate is given by

$$T_c = \frac{(X_c - X_i)T_{i+1} + (X_{i+1} - X_c)T_i}{X_{i+1} - X_i}$$

3.2.5.6 To adjust the SDs for their relationship with the characteristic mean the estimated trend values are subtracted from the transformed SDs and the grand mean is added back.

3.2.5.7 The results for the simple example with 16 varieties are illustrated in Figure 2.

**Figure 2: Adjusting for association between SD and mean – days to ear emergence in cocksfoot varieties** (symbol A is for adjusted SD)



Step 5: Calculation of the uniformity criterion

3.2.5.8 An estimate of the variability in the uniformity of the reference varieties is derived by applying a one-way analysis of variance to the adjusted log SDs, i.e. with years as the classifying factor. The variability (V) is estimated from the residual term in this analysis of variance.

3.2.5.9 The maximum allowable standard deviation (the uniformity criterion), based on k years of trials, is

$$UC_p = SD_r + t_p \sqrt{V \left( \frac{1}{k} + \frac{1}{Rk} \right)}$$

where  $SD_r$  is the mean of adjusted log SDs for the reference varieties,  $V$  is the variance of the adjusted log SDs after removing year effects,  $t_p$  is the one-tailed t-value for probability  $p$  with degrees of freedom as for  $V$ ,  $k$  is the number of years and  $R$  is the number of reference varieties.

### 3.2.6 Early decisions for a three-year test

3.2.6.1 Decisions on uniformity may be made after two or three years depending on the crop. If COYU is normally applied over three years, it is possible to make an early acceptance or rejection of a candidate variety using an appropriate selection of probability values.

3.2.6.2 The probability level for early rejection of a candidate variety after two years should be the same as that for the full three-year test. For example, if the three-year COYU test is applied using a probability level of 0.2%, a candidate variety can be rejected after two years if its uniformity exceeds the COYU criterion with probability level 0.2%.

3.2.6.3 The probability level for early acceptance of a candidate variety after two years should be larger than that for the full three-year test. As an example, if the three-year COYU test is applied using a probability level of 0.2%, a candidate variety can be accepted after two years if its uniformity does not exceed the COYU criterion with probability level 2%.

3.2.6.4 Some varieties may fail to be rejected or accepted after two years. In the example set out in paragraphs 26 and 27, a variety might have a uniformity that exceeds the COYU criterion with probability level 2% but not the criterion with probability level 0.2%. In this case, such varieties should be re-assessed after three years.

### 3.2.7 Example of COYU calculations

3.2.7.1 An example of the application of COYU is given here to illustrate the calculations involved. The example consists of days to ear emergence scores for perennial ryegrass over three years for 11 reference varieties (R1 to R11) and one candidate (C1). The data is tabulated in Table 1.

**Table 1: Example data-set – days to ear emergence in perennial ryegrass**

Variety	Character Means			Within Plot SD			Log (SD+1)		
	Year 1	Year 2	Year 3	Year 1	Year 2	Year 3	Year 1	Year 2	Year 3
R1	38	41	35	8.5	8.8	9.4	2.25	2.28	2.34
R2	63	68	61	8.1	7.6	6.7	2.21	2.15	2.04
R3	69	71	64	9.9	7.6	5.9	2.39	2.15	1.93
R4	71	75	67	10.2	6.6	6.5	2.42	2.03	2.01
R5	69	78	69	11.2	7.5	5.9	2.50	2.14	1.93
R6	74	77	71	9.8	5.4	7.4	2.38	1.86	2.13
R7	76	79	70	10.7	7.6	4.8	2.46	2.15	1.76
R8	75	80	73	10.9	4.1	5.7	2.48	1.63	1.90
R9	78	81	75	11.6	7.4	9.1	2.53	2.13	2.31
R10	79	80	75	9.4	7.6	8.5	2.34	2.15	2.25
R11	76	85	79	9.2	4.8	7.4	2.32	1.76	2.13
C1	52	56	48	8.2	8.4	8.1	2.22	2.24	2.21

3.2.7.2 The calculations for adjusting the SDs in year 1 are given in Table 2. The trend value for candidate C1 is obtained by interpolation between values for varieties R1 and R2, since the characteristic mean for C1 (i.e. 52) lies between the means for R1 and R2 (i.e. 38 and 63). That is

$$T_c = \frac{(X_c - X_i)T_{i+1} + (X_{i+1} - X_c)T_i}{X_{i+1} - X_i} = \frac{(52 - 38) \times 2.28 + (63 - 52) \times 2.21}{63 - 38} = 2.28$$

**Table 2: Example data-set – calculating adjusted log(SD+1) for year 1**

Variety	Ranked mean (X)	Log (SD+1) (Y)	Trend Value T	Adj. Log (SD+1)
R1	38	2.25	$(2.25 + 2.21 + 2.39)/3 = 2.28$	$2.25 - 2.28 + 2.39 = 2.36$
R2	63	2.21	$(2.25 + 2.21 + 2.39)/3 = 2.28$	$2.21 - 2.28 + 2.39 = 2.32$
R3	69	2.39	$(2.25 + \dots + 2.42)/5 = 2.35$	$2.39 - 2.35 + 2.39 = 2.42$
R5	69	2.50	$(2.25 + \dots + 2.48)/7 = 2.38$	$2.50 - 2.38 + 2.39 = 2.52$
R4	71	2.42	$(2.25 + \dots + 2.32)/9 = 2.38$	$2.42 - 2.38 + 2.39 = 2.43$
R6	74	2.38	$(2.21 + \dots + 2.53)/9 = 2.41$	$2.38 - 2.41 + 2.39 = 2.36$
R8	75	2.48	$(2.39 + \dots + 2.34)/9 = 2.42$	$2.48 - 2.42 + 2.39 = 2.44$
R7	76	2.46	$(2.42 + \dots + 2.34)/7 = 2.42$	$2.46 - 2.42 + 2.39 = 2.43$
R11	76	2.32	$(2.48 + \dots + 2.34)/5 = 2.43$	$2.32 - 2.43 + 2.39 = 2.28$
R9	78	2.53	$(2.32 + 2.53 + 2.34)/3 = 2.40$	$2.53 - 2.40 + 2.39 = 2.52$
R10	79	2.34	$(2.32 + 2.53 + 2.34)/3 = 2.40$	$2.34 - 2.40 + 2.39 = 2.33$
Mean	70	2.39		
C1	52	2.22	2.28	$2.22 - 2.28 + 2.39 = 2.32$

3.2.7.3 The results of adjusting for all three years are shown in Table 3.

**Table 3: Example data-set – adjusted log(SD+1) for all three years with over-year means**

Variety	Over-Year Means		Adj. Log (SD+1)		
	Char. mean	Adj. Log (SD+1)	Year 1	Year 2	Year 3
R1	38	2.26	2.36	2.13	2.30
R2	64	2.10	2.32	2.00	2.00
R3	68	2.16	2.42	2.10	1.95
R4	71	2.15	2.43	1.96	2.06
R5	72	2.20	2.52	2.14	1.96
R6	74	2.12	2.36	1.84	2.16
R7	75	2.14	2.43	2.19	1.80
R8	76	2.02	2.44	1.70	1.91
R9	78	2.30	2.52	2.16	2.24
R10	78	2.22	2.33	2.23	2.09
R11	80	2.01	2.28	1.78	1.96
Mean	70	2.15	2.40	2.02	2.04
C1	52	2.19	2.32	2.08	2.17

3.2.7.4 The analysis of variance table for the adjusted log SDs is given in Table 4 (based on reference varieties only). The variability in the uniformity of reference varieties is estimated from this ( $V=0.0202$ ).

**Table 4: Example data set – analysis of variance table for adjusted log (SD+1)**

Source	Degrees of freedom	Sums of squares	Mean squares
Year	2	1.0196	0.5098
Varieties within years (=residual)	30	0.6060	<b>0.0202</b>
Total	32	1.6256	

3.2.7.5 The uniformity criterion for a probability level of 0.2% is calculated thus:

$$UC_p = SD_r + t_p \sqrt{V \left( \frac{1}{k} + \frac{1}{Rk} \right)} = 2.15 + 3.118 \times \sqrt{0.0202 \times \left( \frac{1}{3} + \frac{1}{3 \times 11} \right)} = 2.42$$

where  $t_p$  is taken from Student's t table with  $p=0.002$  (one-tailed) and 30 degrees of freedom.

3.2.7.6 Varieties with mean adjusted log (SD + 1) less than, or equal to, 2.42 can be regarded as uniform for this characteristic. The candidate variety C1 satisfies this criterion.

### 3.2.8 Implementing COYU

The COYU criterion can be applied using the DUST software package for the statistical analysis of DUS data. This is available from the Dr. Sally Watson, Biometrics Division, Department of Agriculture for Northern Ireland, Newforge Lane, Belfast BT9 5PX, UK .



### 3.2.9 COYU Software

#### 3.2.9.1 DUST Computer program

3.2.9.1.1 The main output from the DUST COYU program is illustrated in Table A1. This summarises the results of analyses of within-plot SDs for 49 perennial ryegrass varieties assessed over a three-year period. Supplementary output is given in Table A2 where details of the analysis of a single characteristic, date of ear emergence, are presented. Note that the analysis of variance table given has an additional source of variation; the variance,  $V$ , of the adjusted log SDs is calculated by combining the variation for the variety and residual sources.

3.2.9.1.2 In Table A1, the adjusted SD for each variety is expressed as a percent of the mean SD for all reference varieties. A figure of 100 indicates a variety of average uniformity; a variety with a value less than 100 shows good uniformity; a variety with a value much greater than 100 suggests poor uniformity in that characteristic. Lack of uniformity in one characteristic is often supported by evidence of poor uniformity in related characteristics.

3.2.9.1.3 The symbols “\*” and “+” to the right of percentages identify varieties whose SDs exceed the COYU criterion after 3 and 2 years respectively. The symbol “:” indicates that after two years uniformity is not yet acceptable and the variety should be considered for testing for a further year. Note that for this example a probability level of 0.2% is used for the three-year test. For early decisions at two years, probability levels of 2% and 0.2% are used to accept and reject varieties respectively. All of the candidates had acceptable uniformity for the 8 characters using the COYU criterion.

3.2.9.1.4 The numbers to the right of percentages refer to the number of years that a within-year uniformity criterion is exceeded. This criterion has now been superseded by COYU.

3.2.9.1.5 The program will operate with a complete set of data or will accept some missing values, e.g. when a variety is not present in a year.

**Table A1: Example of summary output from COYU program**

\*\*\*\* OVER-YEARS UNIFORMITY ANALYSIS SUMMARY \*\*\*\*

WITHIN-PLOT STANDARD DEVIATIONS AS % MEAN OF  
REFERENCE VARIETY SDS

	CHARACTERISTIC NUMBER					
	5	60	8	10	11	
R1	100	100	95 1	100	97	97
R2	105	106	98	99	104	101
R3	97	103	92 1	103	96	98
R4	102	99	118 2	105	101	101
R5	102	99	116 3	95	104	110
R6	103	102	101	99	97	104
R7	100	95	118 2	102 1	98	99
R8	97	98	84	95	97	93
R9	97	105	87	99	101	99
R10	104	100	96	105 1	96	102
R11	99	96	112	99	101	98
R12	100	97	99 1	103	105	106
R13	95	96	101	100	96	101
R14	105	103	90	97	101	97
R15	102	100 1	89	105	105 1	101
R16	99	98	92 1	98	102	98
R17	97	101	98	101	101	95
R18	99	97	96	96	102	99
R19	103	101	105	102	100	98
R20	104	99	93	91	100	102
R21	97	94	103	97	100	102
R22	101	110*1	112	107 1	103 1	101
R23	94	101	107	99	104	97
R24	99	97	95	99	100	103
R25	104 1	103	93 1	99	101	96
R26	98	97	111 2	96	102 1	106
R27	102	99	106 1	99	103	107
R28	101	106	90	95	101	101
R29	101	105	83	102	94	93
R30	99	96	97	99	95	100
R31	99	102	107	107 1	102	99
R32	98	93	111 2	102	98	103
R33	104	102 1	107 1	103	100	97
R34	95	94	82	95	97	96
R35	100	102	95	100	99	94
R36	99	98	111 1	99	100	103
R37	100	107 1	107	101	100	107
R38	95	97	102	107 1	97	101
R39	99	99	90	98	101	100
R40	104	102	112 1	100	101	97
C1	100 1	106	113 2	104 1	106 1	106
C2	103	101	98	97	101	109
C3	97	93	118 2	98	99	109
C4	102	101	106	103	99	101
C5	100	104	99	103	100	107
C6	101	102	103	100	103	107
C7	96	98	106	97	102	103
C8	101	105 1	116 2	103	103	93
C9	99	99	90 2	91	97	98

CHARACTERISTIC

5	SPRING	60	NATURAL SPRIN
8	DATE OF EAR	10	HEIGHT AT EAR
11	WIDTH AT EAR	14	LENGTH OF FLA
15	WIDTH OF FLAG	24	EAR LENGTH

SYMBOLS

\* - SD EXCEEDS OVER-YEARS CRITERION AFTER  
 + - SD EXCEEDS OVER-YEARS CRITERION AFTER  
 : - SD NOT YET ACCEPTABLE AFTER 2 YEARS V  
 1,2,3 - THE NUMBER OF OCCASIONS THE WITHIN-YE

TGP/8/1 Draft 3  
page 99

\*\*\*\* UNIFORMITY ANALYSIS OF BETWEEN-PLANT STANDARD DEVIATIONS (SD) \*\*\*\*

VARIETY	OVER-YEARS			INDIVIDUAL YEARS								
	CHAR.	ADJ.	UNADJ	CHAR. MEAN			LOG (SD+1)			ADJ LOG (SD+1)		
	MEAN	LOG SD	LOG SD	88	89	90	88	89	90	88	89	90
REFERENCE												
R3	38.47	1.823	2.179	39.07	41.21	35.12	2.02	2.18	2.34X	1.73	1.78	1.96
R5	50.14	2.315	2.671	48.19	53.69	48.54	2.52X	2.74X	2.76X	2.23	2.33	2.39
R16	59.03	1.833	2.179	57.25	63.33	56.50	2.28X	2.24	2.01	1.96	1.73	1.81
R26	63.44	2.206	2.460	61.00	66.53	62.81	2.50X	2.75X	2.13	2.18	2.33	2.11
R9	63.99	1.739	1.994	62.92	68.32	60.72	2.21	2.03	1.74	1.96	1.64	1.62
R12	66.12	1.964	2.086	67.89	65.35	65.12	2.07	2.58X	1.60	1.97	2.14	1.78
R33	67.58	2.124	2.254	66.66	71.54	64.53	2.55X	2.26	1.95	2.32	1.92	2.12
R1	67.87	1.880	1.989	69.07	70.64	63.90	1.60	2.45X	1.93	1.60	2.08	1.96
R20	68.74	1.853	1.893	67.17	74.31	64.74	2.05	1.95	1.68	1.92	1.75	1.89
R25	68.82	1.853	1.905	68.28	72.38	65.81	1.83	2.39X	1.49	1.75	2.09	1.72
R18	69.80	1.899	1.853	68.61	75.22	65.58	1.88	1.84	1.84	1.82	1.80	2.08
R30	70.53	1.919	1.864	70.36	75.08	66.15	2.04	1.84	1.71	2.00	1.78	1.98
R13	70.63	2.005	2.000	70.23	75.00	66.66	1.97	2.03	2.01	1.91	1.86	2.24
R32	71.49	2.197	2.238	70.03	74.98	69.44	2.32X	2.45X	1.94	2.31	2.27	2.01
R34	72.09	1.630	1.545	71.32	77.35	67.59	1.57	1.49	1.58	1.54	1.58	1.78
R40	72.24	2.222	2.178	72.71	75.07	68.95	2.25X	2.26	2.03	2.29	2.16	2.22
R23	72.40	2.122	2.058	69.72	78.39	69.10	2.11	2.14	1.93	2.16	2.14	2.06
R29	72.66	1.657	1.580	73.13	75.80	69.04	1.46	1.63	1.65	1.47	1.69	1.81
R7	73.19	2.341	2.342	72.23	75.80	71.52	2.62X	2.30X	2.10	2.61	2.30	2.11
R24	73.19	1.888	1.796	74.00	76.37	69.20	1.62	1.84	1.93	1.71	1.91	2.04
R19	73.65	2.083	2.049	73.32	76.06	71.57	1.96	2.05	2.14	1.96	2.13	2.16
R2	73.85	1.946	1.897	72.98	78.16	70.42	1.76	1.96	1.97	1.79	2.02	2.03
R31	74.23	2.119	2.012	73.73	78.23	70.71	2.05	1.86	2.13	2.25	1.94	2.17
R37	74.38	2.132	2.020	74.87	76.95	71.32	1.97	2.04	2.04	2.23	2.11	2.06
R11	74.60	2.224	2.150	73.87	78.07	71.87	2.21	2.08	2.16	2.36	2.10	2.21
R38	74.76	2.029	1.916	76.11	78.24	69.93	1.84	2.15	1.75	1.98	2.24	1.87
R8	74.83	1.677	1.593	74.27	78.77	71.45	1.62	1.55	1.61	1.75	1.64	1.64
R15	75.54	1.760	1.682	75.72	78.68	72.22	1.53	1.79	1.73	1.64	1.84	1.80
R10	75.64	1.915	1.847	73.47	79.24	74.23	1.87	1.66	2.00	1.99	1.78	1.98
R22	75.68	2.228	2.133	74.57	79.17	73.32	2.18	2.21	2.01	2.40	2.26	2.03
R14	75.84	1.797	1.688	74.53	79.56	73.43	1.54	1.63	1.90	1.70	1.76	1.93
R17	76.13	1.942	1.832	75.34	79.09	73.96	1.65	2.04	1.81	1.90	2.10	1.83
R39	76.83	1.781	1.676	75.49	80.50	74.50	1.56	1.51	1.96	1.72	1.70	1.92
R35	77.22	1.886	1.773	76.67	80.85	74.15	1.73	1.67	1.92	1.88	1.85	1.93
R4	77.78	2.349	2.268	76.80	81.22	75.33	2.36X	2.13	2.31X	2.52	2.33	2.20
R36	77.98	2.209	2.173	78.97	79.85	75.11	2.13	2.15	2.25X	2.24	2.21	2.18
R6	78.73	2.009	1.935	77.53	82.88	75.78	2.00	1.75	2.06	2.03	2.09	1.91
R27	78.78	2.116	2.098	77.61	80.03	78.69	1.80	2.25	2.24X	1.87	2.39	2.09
R28	79.41	1.785	1.722	78.28	81.99	77.97	1.68	1.43	2.05	1.79	1.67	1.89
R21	80.52	2.045	1.950	77.43	85.02	79.11	1.98	1.75	2.13	2.07	2.09	1.98
CANDIDATE												
C1	64.03	2.252	2.438	63.85	63.33	64.92	2.49X	2.81X	2.02	2.25	2.29	2.21
C2	86.11	1.940	1.837	84.83	88.63	84.85	1.79	1.71	2.01	1.90	2.05	1.87
C3	82.04	2.349	2.248	82.26	87.45	76.40	2.37X	2.03	2.35X	2.48	2.37	2.20
C4	78.63	2.104	2.033	78.01	82.17	75.72	2.05	2.01	2.04	2.15	2.27	1.90
C5	72.99	1.973	1.869	71.98	79.40	67.59	1.95	1.78	1.88	1.93	1.90	2.08
C6	83.29	2.050	1.947	84.10	85.57	80.21	2.05	1.69	2.10	2.16	2.03	1.96
C7	83.90	2.100	1.997	84.12	87.99	79.60	1.93	1.95	2.11	2.04	2.29	1.97
C8	83.50	2.304	2.201	82.43	85.98	82.08	2.27X	2.00	2.34X	2.38	2.33	2.20
C9	51.89	1.788	2.157	52.35	55.77	47.56	1.83	2.34X	2.31X	1.52	1.91	1.93
MEAN OF REFERENCE	71.47	1.988		70.78	74.97	68.65	1.97	2.03	1.96	1.99	1.99	1.99

UNIFORMITY CRITERION	PROB. LEVEL	
3-YEAR REJECTION	2.383	0.002
2-YEAR REJECTION	2.471	0.002
2-YEAR ACCEPTANCE	2.329	0.020

\*\*\*\* ANALYSIS OF VARIANCE OF ADJUSTED LOG (SD+1) \*\*\* \*

	DF	MS	F RATIO
YEARS	2	0.06239	
VARIETIES	39	0.11440	5.1
RESIDUAL	78	0.02226	
TOTAL	119	0.05313	

SYMBOLS

- \* - SD EXCEEDS OVER-YEARS UNIFORMITY CRITERION AFTER 3 YEARS.
- + - SD EXCEEDS OVER-YEARS UNIFORMITY CRITERION AFTER 2 YEARS.
- : - SD NOT YET ACCEPTABLE ON OVER-YEARS CRITERION AFTER 2 YEARS.
- X - SD EXCEEDS 1.265 TIMES MEAN OF REFERENCE VARIETIES

### 3.3 Standard probability levels used for COYD and COYU

The following four cases are those which, in general, represent the different situations which may arise where COYD and COYU are used in DUS testing:

Scheme A. Test is conducted over 2 independent growing cycles and decisions made after 2 growing cycles (a growing cycle could be a year and is further on denoted by cycle)

Scheme B. Test is conducted over 3 independent growing cycles and decisions made after 3 cycles

Scheme C. Test is conducted over 3 independent growing cycles and decisions made after 3 cycles, but a variety may be accepted after 2 cycles

Scheme D. Test is conducted over 3 independent growing cycles and decisions made after 3 cycles, but a variety may be accepted or rejected after 2 cycles

The stages at which the decisions are made in Cases A to D are illustrated in figures 1 to 4 respectively. These also illustrate the various standard probability levels ( $p_{d2}$ ,  $p_{nd2}$ ,  $p_{d3}$ ,  $p_{u2}$ ,  $p_{nu2}$  and  $p_{u3}$ ) which are needed to calculate the COYD and COYU criteria depending on the case. These are defined as follows:


<b>Probability Level</b>	<b>Used to decide whether a variety is :-</b>
$p_{d2}$	distinct after 2 cycles
$p_{nd2}$	non-distinct in a characteristic after 2 cycles
$p_{d3}$	distinct after 3 cycles
$p_{u2}$	uniform in a characteristic after 2 cycles
$p_{nu2}$	non-uniform after 2 cycles
$p_{u3}$	uniform in a characteristic after 3 cycles

In figures 1 to 4 the COYD criterion calculated using say the probability level  $p_{d2}$  is denoted by  $LSD_{p_{d2}}$  etc., and the COYU criterion calculated using say the probability level  $p_{u2}$  is denoted by  $UC_{p_{u2}}$  etc. The term “diff” represents the difference between the means of a candidate variety and another variety for a characteristic, while “U” represents the mean adjusted  $\log(SD+1)$  of a variety for a characteristic.

Table 1 summarises the various standard probability levels needed to calculate the COYD and COYU criteria in each of Cases A to D. For example, in Case B only two probability levels are needed ( $p_{d3}$  and  $p_{u3}$ ), whereas Case C requires four ( $p_{d2}$ ,  $p_{d3}$ ,  $p_{u2}$  and  $p_{u3}$ ).

CASE	COYD			COYU		
	$p_{d2}$	$p_{nd2}$	$p_{d3}$	$p_{u2}$	$p_{nu2}$	$p_{u3}$
A						
B						
C						
D						

Figure 1. COYD and COYU decisions and standard probability levels ( $p_i$ ) in Case A



NOTE:-


“diff” is the difference between the means of the candidate variety and another variety for the characteristic.

LSD<sub>p</sub> is the COYD criterion calculated at probability level p.

“U” is the mean adjusted log(SD+1) of the candidate variety for the characteristic.

UC<sub>p</sub> is the COYU criterion calculated at probability level p.

Figure 2. COYD and COYU decisions and standard probability levels ( $p_i$ ) in Case B



NOTE:-


“diff” is the difference between the means of the candidate variety and another variety for the characteristic.

LSD<sub>p</sub> is the COYD criterion calculated at probability level p.

“U” is the mean adjusted log(SD+1) of the candidate variety for the characteristic.

UC<sub>p</sub> is the COYU criterion calculated at probability level p.

Figure 3. COYD and COYU decisions and standard probability levels ( $p_i$ ) in Case C



NOTE:-


“diff” is the difference between the means of the candidate variety and another variety for the characteristic.

LSD $p$  is the COYD criterion calculated at probability level  $p$ .

“U” is the mean adjusted  $\log(\text{SD}+1)$  of the candidate variety for the characteristic.

UC $p$  is the COYU criterion calculated at probability level  $p$ .

Figure 4. COYD and COYU decisions and standard probability levels ( $p_i$ ) in Case D



NOTE:-

“diff” is the difference between the means of the candidate variety and another variety for the characteristic.

LSDp is the COYD criterion calculated at probability level p.

“U” is the mean adjusted  $\log(\text{SD}+1)$  of the candidate variety for the characteristic

UCp is the COYU criterion calculated at probability level p.



## **4. Parent Formula of Hybrid Varieties**

### 4.1 Introduction

The use of the parental formula requires that the difference between parent lines is sufficient to ensure that the hybrid obtained from those parents is distinct. The method is based on the following steps:

- (i) description of parent lines according to the Test Guidelines;
- (ii) checking the originality of those parent lines in comparison with the variety collection, based on the table of characteristics in the Test Guidelines, in order to identify similar parent lines;
- (iii) checking the originality of the hybrid formula in relation to the hybrids in the variety collection, taking into account the most similar parent lines; and
- (iv) assessment of distinctness at the hybrid level for varieties with a similar formula.

### 4.2 Requirements of the method

The application of the method requires:

- (i) a declaration of the formula and submission of plant material of the parent lines of hybrid varieties;
- (ii) inclusion in the variety collection of the parent lines used as parents in the hybrid varieties of the variety collection (for guidance on the constitution of a variety collection see document TGP/4 section 1) and a list of the formulae of the hybrid varieties;
- (iii) application of the method to all varieties in the variety collection. This condition is important to obtain the full benefit; and
- (iv) a rigorous approach to assess the originality of any new parent line in order to be confident on the distinctness of the hybrid variety based on that parent line.

### 4.3 Assessing the originality of a new parent line

4.3.1 The originality of a parental line is assessed using the characteristics included in the relevant Test Guidelines.

4.3.2 The difference between parent lines must be sufficient to be sure that hybrids produced using different parent lines will be distinct. For example:

Characteristic 1: a characteristic having two states of expression (absent/present), which are determined by two alleles of a single gene, with one dominant allele (+) for the expression “present” and one recessive allele (-) for the expression “absent”.


Three parent lines:

- A: with the recessive allele (-) with expression “absent”
- B: with the dominant allele (+) with expression “present”
- C: with the dominant allele (+) with expression “present”

Crossing the above-mentioned parent lines to obtain the following F1 hybrids:

- (A x C): having expression “present” for Characteristic 1
- (B x C): having expression “present” for Characteristic 1

The following diagram shows the ways the two different crossings result in the same expression of Characteristic 1 (i.e. “present” in both hybrids), although parent line A(-) and parent line B(+) have different expressions.



4.3.3 Although the parent lines A and B are clearly different for characteristic 1, the two hybrid varieties A x C and B x C have the same expression. Thus, a difference between A and B for Characteristic 1 is not sufficient.

4.3.4 With a more complex genetic control involving several genes, not precisely described, the interaction between the different alleles of each gene and between genes might also lead to similar expression at the level of the hybrid varieties. In such cases, a larger difference is appropriate to establish distinctness between two parent lines.

4.3.5 Determining the difference required is mainly based on a good knowledge of the species, of the characteristics and, when available, on their genetic control.

#### 4.4 Verification of the formula

4.4.1 The aim of verifying the formula is to check if the candidate hybrid variety has been produced by crossing the parent lines declared and submitted by the applicant.

4.4.2 Different characteristics can be used to perform this check when the genetic pattern of each parent can be identified in the hybrid. Generally, characteristics based on polymorphism of enzymes or of some storage proteins can be used.

4.4.3 If no suitable characteristics are available, the only possibility is to cross the parent lines using the plant material submitted by the applicant and to compare the hybrid variety seedlots (the sample submitted by the applicant and the sample harvested after the cross).

4.5 Uniformity and stability of parent lines

4.5.1 The uniformity and stability of the parent lines should be assessed according to the appropriate recommendations for the variety concerned. The uniformity and stability of the parent lines are important for the stability of the hybrid. Another requirement for the stability of the hybrid is the use of the same formula for each cycle of the hybrid seed production.

4.5.2 A check of the uniformity on the hybrid should also be done, even if distinctness of the hybrid has been established on the basis of the parent lines.

4.6 Description of the hybrid

4.6.1 A description of the hybrid variety should be established, even where the distinctness of the hybrid has been established on the basis of the parent formula.

## 5. The GAIA methodology

[TWC: to delete references to multivariate analysis and refer to phenotypic distance. To harmonize the wording with Section 2.3 Types of characteristics and their scale level.]

The GAIA method has been developed by experts from France to calculate a phenotypic distance between two varieties. The principle is to compute a phenotypic distance between a pair of varieties, which is a sum of distances for individual characteristics. For each species, this system must be calibrated to determine the weighting given to the difference in each characteristic used and the threshold for the phenotypic distance used to eliminate varieties from the growing trial.

### 5.1. Weighting of characteristics

5.1.1 Weighting is defined as the contribution in a given characteristic to the total distance between a pair of varieties. For each species, this system must be calibrated to determine the weight which can be given to each difference and to evaluate the reliability of each characteristic in a given environment and for the genetic variability concerned. For that reason the role of the crop expert is essential.

5.1.2 Weighting depends on the size of the difference and on the individual characteristic. The weightings are defined by crop experts on the basis of their expertise in the crop and on a “try-and-check” learning process and stored in the GAIA database. Experts can give zero weighting to small differences, thus, even if two varieties have different observed values in many characteristics, the overall distance might be zero. The same weighting is attributed to any pair of varieties whose absolute differences between observed values are the same for a given characteristic.

5.1.3 The weighting should be simple and consistent. The following three rules are given:

- (i) the distances for the characteristic should be integer values, i.e. 0, 1, 2, 3, etc. where 3 is considered to be about 3 times greater than 1;
- (ii) if for a characteristic a given difference “expressed as an absolute value” is considered as a double distance for character *a* compared to character *b*, the distance value for this difference should be double that in character *a* than it is in character *b*;
- (iii) define the values by “try-and-check” (see Diagram 1.)

5.1.4 The following simple example shows the computation of the distance between two varieties on the basis of a qualitative characteristic:

Example: taking the characteristic “Shape of ear”, observed on a 1 to 3 scale, the crop experts have attributed weighting to differences which they consider significant:

Shape of ear:

- 1 = conical
- 2 = conico-cylindrical
- 3 = cylindrical

Comparison between difference in notes and weighting		
	Different in notes	Weighting
conical (1) vs. conical (1)	0	0
conical (1) vs. conico-cylindrical (2)	1	2
conical (1) vs. cylindrical (3)	3	6
conico-cylindrical (2) vs. conico-cylindrical (2)	0	0
conico-cylindrical (2) vs. cylindrical (3)	1	2
cylindrical (3) vs. cylindrical (3)	0	0

When the crop experts compare a variety ‘i’ with conical ear (note 1) to a variety ‘j’ with cylindrical ear (note 3), they attribute a weighting of 6 etc. The weightings are summarized in the form of a weighting matrix:

Weighting matrix ‘i’				
		Variety i		
		1	2	3
Variety ‘j’	1	0	2	6
	2		0	2
	3			0

When the crop experts compare a variety i with conical ear (note 1) to a variety j with cylindrical ear (note 3), they attribute a weighting of 6.

## 5.2. Determining “Distinctness Plus”

The threshold for the phenotypic distance used to eliminate varieties from the growing trial is called “Distinctness Plus” and is determined by the crop expert at a level which is higher than the difference needed to establish distinctness. This ensures that all pairs of varieties having a distance equal or greater than the threshold (Distinctness Plus) would be distinct in the growing trial. The Distinctness Plus threshold must be based on experience gained with the varieties of common knowledge and must minimize the possibility of not including in the growing trial a pair of varieties which should be further compared in the field.

## 5.3. Computing GAIA phenotypic distance

5.3.1 The principle is to compute a phenotypic distance between two varieties, which is the total distance between a pair of varieties as the result of the addition of the weightings of all characteristics (see section 2 of this Annex). Thus, the GAIA phenotypic distance is:

$$dist(i, j) = \sum_{k=1, nchar} W_k(i, j)$$

where:

$dist(i, j)$  is the computed distance between variety  $i$  and variety  $j$ .

$k$  is the  $k^{th}$  characteristic, from the  $nchar$  characteristics selected for computation.

$W_k(i, j)$  is the weighting of characteristics  $k$ , which is a function of the difference observed between variety  $i$  and variety  $j$  for that characteristic  $k$ .

$OV_{ki}$  is the observed value on characteristic  $k$  for variety  $i$ .

$$W_k(i, j) = f(|OV_{ki} - OV_{kj}|)$$

5.3.2 Varieties are compared in pairs in different combinations of pair-wise comparisons, e.g.:

- compare two varieties,
- compare a given variety to all varieties in the variety collection,
- compare all candidate varieties to all the other candidate varieties and the varieties in the variety collection,
- compare all possible combinations.

#### 5.4. The GAIA software


5.4.1 The GAIA software allows the computation of the phenotypic distance using qualitative, quantitative or electrophoretic characteristics, which can be used alone or in combination. The user can decide on the type of data and the way it is used:

- (i) select all the available characteristics, or different subsets of characteristics.
- (ii) define different weighting values:
  - experts can choose different values as the weighting/distance for a characteristic (1, 2, 5, etc.);
  - some crops have more characteristics than others;
  - the crop expert can use all available information, or a subset of characteristics only.
- (iii) the way the Distinctness Plus threshold is used:
  - a low Distinctness Plus threshold, which helps to find the more difficult cases (to identify similar varieties- very often used by crop experts);
  - intermediate Distinctness Plus threshold (different levels according to the needs);
  - a large Distinctness Plus threshold when there is a need to have a comparison which uses all the available characteristics;
  - a Distinctness Plus threshold greater than the maximum distance possible on all characteristics to see all available raw data and the weightings for each characteristic

5.4.2 The software provides a comprehensive report for each pair-wise comparison. It computes an overall distance, but also provides all the individual absolute values and the distance contribution of each characteristic (see section 6 of this Annex).

5.4.3 In order to minimize computation time, as soon as the threshold is achieved for a comparison between two given varieties, the software proceeds to the next pair of varieties. Remaining characteristics and their raw values will not be shown in the summary output, and will not contribute to the distance.

**Diagram 1: “Try-and-check” process to define and revise the weightings for a crop**





5.4.4 Section 6 of this Annex provides a screen copy of a display tree which shows how the expert can navigate and visualise the results of computations.

5.4.5 GAIA software has been developed with WINDEV-7.5. The general information (species, characteristics, weighting, etc.), the data collected on the varieties and the results of computations are stored in an integrated database. Import and export facilities allow for other information systems to be used in connection with the GAIA software. ODBC allows access to the GAIA database and to other databases simultaneously.

5.4.6 For qualitative characteristics, 1 or 2 notes per variety can be used. In general, two notes are present when there are two trial locations. For electrophoresis data, only one description can be entered per variety. For quantitative characteristics at least 2 values (different trials, repeats, etc.) are necessary and the user selects which to use in the computation.

5.4.7 GAIA is most suitable for self-pollinated and vegetatively propagated varieties, but can also be used for other types of varieties.

## 5.5. Using the GAIA methodology

The GAIA methodology can be used:

- (i) to eliminate from subsequent growing cycles all pairs of varieties reaching or surpassing the GAIA distance threshold;
- (ii) to focus on close varieties, having a GAIA distance lower than the threshold, for the next growing cycle(s).

### *5.5.1 Using phenotypic distance in the first growing cycle*

5.5.1.1 A crop that has a large variety collection and uses only quantitative characteristics on a 1 to 9 scale; the GAIA methodology allows the selection of varieties to be included in the growing trial. This can be used to plan the first growing cycle trials as well as the subsequent growing cycles.

5.5.1.2 In crops with relatively few candidates and a small variety collection, which enables the crop expert to sow all candidates (e.g. an agricultural crop), and the appropriate reference varieties, in two or three successive growing cycles. The same varieties are sown in growing cycles 1, 2 and 3, in a randomized layout. The software will help to identify the pairs with a small distance, to enable the expert to focus his attention on these particular cases when visiting the field.

### *5.5.2 Using phenotypic distance after the first growing trial*

5.5.2.1 After one growing cycle (e.g. in the examination of an ornamental crop), the absolute data and distance computations are an objective way to confirm the opinion or the decision of the expert. There might be cases where pairs of varieties have a small distance, but nevertheless the expert has clear evidence of distinctness. If more growing cycles are necessary before a decision is taken, the software helps to identify on which cases the expert will need to focus.

5.5.2.2 In cases where there are many candidate and reference varieties and there is a wide variability in the species (e.g. a vegetable crop); on the one hand there are already obvious differences after only one cycle, but on the other hand some varieties are very similar. In order to be more efficient in their checks, the crop experts wish to grow “similar” varieties close to each other. The raw results and distances will help to select the “similar” varieties and decide on the layout of the trial for the next growing cycle.


5.5.2.3 In crops in which there are many similar varieties, for which it is a common practice to make side-by-side comparisons, GAIA can be used to identify the similar varieties after the first cycle, in particular, when the number of varieties in a trial increases, making it less easy to identify all the problem situations. The software can help to “not miss” the less obvious cases.

5.5.2.4 In vegetatively propagated ornamental varieties, the examination lasts for one or two growing cycles: after the first growing cycle, some reference varieties in the trial are obviously different from all candidates, and their inclusion in the second growing cycle is not necessary. When the number of varieties is large, the raw data and distance(s) can help the expert to detect reference varieties for which the second growing cycle is unnecessary.

## 5.6. Example with qualitative, electrophoretic and quantitative characteristics (*Zea mays* data)

### 5.6.1 Introduction

5.6.1.1 The software can use qualitative, quantitative and/or electrophoretic data. These types of data can be used alone or in combination, as shown in Diagram 2.



5.6.1.2 In this example, it is assumed that the crop expert has decided to use a Distinctness Plus threshold  $S_{dist}$  of 10 (see section 2 of this Annex).

### 5.6.2 Qualitative Analysis

5.6.2.1 For each characteristic, weightings according to differences between levels of expression are pre-defined in a matrix of distances.

5.6.2.2 “Shape of ear”: observed on a 1 to 3 scale, the crop experts have attributed weightings greater than zero to differences which they consider significant:

1 = conical  
2 = conico-cylindrical  
3 = cylindrical

		Variety ‘i’		
		1	2	3
Variety ‘j’	1	0	2	6
	2		0	2
	3			0

When the crop experts compare a variety ‘i’ with conical ear (note 1) to a variety ‘j’ with cylindrical ear (note 3), they attribute a weighting of 6.

5.6.2.3 “Length of husks”, observed on a 1 to 9 scale, the crop experts have defined the following weighting matrix:

1 = very short  
2 = very short to short  
3 = short  
4 = short to medium  
5 = medium  
6 = medium to long  
7 = long  
8 = long to very long  
9 = very long

		Variety ‘i’								
		1	2	3	4	5	6	7	8	9
Variety ‘j’	1	0	0	0	2	2	2	2	2	2
	2		0	0	0	2	2	2	2	2
	3			0	0	0	2	2	2	2
	4				0	0	0	2	2	2
	5					0	0	0	2	2
	6						0	0	0	2
	7							0	0	0
	8								0	0
	9									0

The weighting between a variety ‘i’ with very short husks (note 1) and a variety ‘j’ with short husks (note 3) is 0. Experts consider a difference of 3 notes is necessary in order to recognise a non-zero distance between two varieties. Even if the difference in notes is greater than 3, the experts do not increase the distance more than 2.

5.6.2.4 The reason for using a lower weighting for some characteristics compared to others can be that they are less “reliable” or “consistent” (e.g. more subject to the effect of the environment); and/or they are considered to indicate a lower distance between varieties.

5.6.2.5 The matrix for a qualitative analysis for 5 characteristics for varieties A and B:

	Ear shape	Husk length	Type of grain	Number of rows of grain	Ear diameter	
Notes for variety A (1 to 9 scale)	1	1	4	6	5	
Notes for variety B (1 to 9 scale)	3	3	4	4	6	
Difference observed	2	2	0	2	1	
<i>Weighting according to the crop expert</i>	6	0	0	2	0	$D_{qual} = 8$

In this example  $D_{qual} = 8 < S_{dist}$  varieties A and B are declared “GAIA NON-distinct” and can be passed on to electrophoretic analysis.

### 5.6.3 Electrophoretic analysis

5.6.3.1 The electrophoretic characteristic is a homozygous allele in the UPOV Test Guidelines (see Diagram 3). The software does not allow the use of heterozygous alleles.




Diagram 3: The Isocitrate Dehydrogenase (IDH) enzyme has two genes (Idh1 and Idh2) located on two different chromosomes. Each of them has two alleles which are observed as 1 (presence) or 0 (absence).

5.6.3.2 Electrophoretic characteristics are noted as 0 or 1 (absence or presence). The decision rule, used to give a weighting to two varieties, is the addition of the weighting number of differences observed and the weighting number of chromosomes related to these differences (see example below):

	Chromosome 8		Chromosome 6	
	Idh1 4	Idh1 6	Idh2 4	Idh2 6
Variety A	0	1	1	0
Variety B	0	1	0	1
Difference	0	0	1	1

5.3.3 In this example, varieties A and B are described for 4 electrophoretic characteristics:

Idh1 4, Idh1 6, Idh2 4 and Idh2 6. The software looks at differences and gives the phenotypic distance using the following computation:

$$D_{elec} = 2 \times 0.25 + 1 \times 1 = 1.5$$

The diagram shows the calculation  $D_{elec} = 2 \times 0.25 + 1 \times 1 = 1.5$ . Four boxes with arrows point to the numbers in the equation:

- A box containing "2 is the number of differences observed" points to the number 2.
- A box containing "0.25 is the weighting attributed by experts to the number of differences" points to the number 0.25.
- A box containing "1 is the number of chromosome on which differences are observed" points to the number 1.
- A box containing "1 is the weighting associated by experts to chromosome." points to the number 1.

5.6.3.4 This formula, which might be difficult to understand, was established by the crop experts in collaboration with biochemical experts. Both the *number of differences* and the *number of chromosomes on which differences are observed* are used. Thus, less importance is attached to differences when these occur on the same chromosome, than when they occur on different chromosomes.

5.6.3.5 After qualitative and electrophoretic analysis, the phenotypic distance between varieties A and B is equal to:

$$D = D_{qual} + D_{elec} = 8 + 1.5 = 9.5$$

The phenotypic distance is *lower than*  $S_{dist}$ , *therefore varieties A and B are considered "GAIA NON-distinct"*.

5.6.3.6 It is not possible to establish distinctness solely on the basis of electrophoretic analysis. It is necessary to have a minimal phenotypic distance in qualitative analysis in order to take into account the electrophoresis results. This minimal phenotypic distance must also be defined by crop experts.

#### 5.6.4 Quantitative Analysis

5.6.4.1 For each quantitative characteristic, the comparison of two varieties is made by looking for consistent differences in at least two different experimental units. Experimental units are defined by the user depending on data present in the database. It can, for example, be the data from two geographic locations of the first growing cycle, or 2 or 3 replications in the case of a single geographical location.

5.6.4.2 For a comparison to be made, the two varieties must be present in the same experimental units. The differences observed must be greater than one of the two threshold values (or minimal distances), fixed by the crop experts.

- $D_{\min-\inf}$  is the lower value from which a weighting is attributed,
- $D_{\min-\sup}$  is the higher minimal distance. These values could be chosen arbitrarily or calculated (15% and 20% of the mean for the trial, or LSD at 1% and 5%, etc.)

5.6.4.3 For each minimal distance a weighting is attributed:

- $D_{\min-\inf}$  a weighting  $P_{\min}$  is attributed;
- $D_{\min-\sup}$  a weighting  $P_{\max}$  is attributed;
- the observed difference is lower than  $D_{\min-\inf}$  a zero weighting is associated.

5.6.4.4 Varieties A and B have been measured for characteristics “Width of blade” and “Length of plant” in two trials.

5.6.4.5 For each trial, and each characteristic, the crop experts have decided to define ( $D_{\min-\inf}$ ) and  $D_{\min-\sup}$  by calculating respectively the 15% and 20% of the mean for the trial:

	Width of blade		Length of plant	
	Trial 1	Trial 2	Trial 1	Trial 2
$D_{\min-\inf} = 15\%$ of the mean	1.2 cm	1.4 cm	28 cm	24 cm
$D_{\min-\sup} = 20\%$ of the mean	1.6 cm	1.9 cm	37 cm	32 cm

5.6.4.6 For each characteristic, the crop experts have attributed the following weighting:

A weighting  $P_{\min} = 3$  is attributed when the difference is greater than  $D_{\min-\inf}$ .

A weighting  $P_{\max} = 6$  is attributed when the difference is greater than  $D_{\min-\sup}$ .

	Width of blade		Length of plant		
	Trial 1	Trial 2	Trial 1	Trial 2	
Variety A	9.9 cm	9.8 cm	176 cm	190 cm	
Variety B	9.6 cm	8.7cm	140 cm	152 cm	
Difference	0.3 cm	1.1 cm	36 cm	38 cm	
Weighting according to the crop expert	0	0	3	6	$D_{\text{quan}} = ?$

5.6.4.7 In this example, for the characteristic “Width of blade”, the differences observed are lower than  $D_{\min-\inf}$ , so no weighting is associated. On the other hand, for the characteristic “Length of plant” one difference is greater than the  $D_{\min-\inf}$  value and the other is greater than the  $D_{\min-\sup}$  value. These two differences are attributed different weightings.

5.6.4.8 The user must decide which weighting will be used for the analysis:

- minimalist option: the weighting chosen is that attributed to the lowest difference;
- maximalist option: the weighting chosen is that attributed to the highest difference;
- mean option: the weighting chosen is the mean of the others.

5.6.4.9 In this example, the crop experts have decided to choose the lowest of the two weightings, so the phenotypic distance based on quantitative characteristics is  $D_{\text{quan}} = 3$ .

5.6.4.10 In summary, at the end of all analysis, the phenotypic distance between varieties A and B is:

$$D = D_{\text{qual}} + D_{\text{elec}} + D_{\text{quan}} = 8 + 1.5 + 3 = 12.5 > S_{\text{dist}}$$

5.6.4.11 The phenotypic distance is greater than the distinction threshold  $S_{\text{dist}}$ , fixed by the crop experts at 10, so varieties A and B are declared “GAIA-distinct”.

5.6.4.12 In this example, the use of electrophoresis data “confirms” a distance between the two varieties; but on the basis of qualitative and quantitative data alone, the threshold is exceeded ( $8 + 3 = 11$  is greater than 10).

5.6.4.13 If the threshold had been set at 6, the difference on the characteristic ear shape would have been sufficient, as variety A is conical and variety B is cylindrical, which is already a clear difference.

1 = conical  
2 = conico-cylindrical  
3 = cylindrical


Variety i			
	1	2	3
1	0	2	6
2		0	2
3			0

### 5.6.5 *Quantitative and qualitative analysis on the same characteristics*

5.6.5.1 For some crops, it is common practice to produce notes on a 1 to 9 scale for quantitative characteristics. Sometimes the transformation process is very simple, sometimes it is a complex process where all available data are used, but with a special manipulation of example varieties to adjust the raw values to the notes on the scale.

5.6.5.2 GAIA can include both as two separate characteristics: the original quantitative scale; and the “transformed into qualitative notes” scale. They are associated in the description of the characteristics. Using the knowledge of this association, when quantitative and qualitative characteristics are both present, only one characteristic is kept, in order to avoid the information being used twice.

5.7 GAIA screen copy.



The upper part shows 3 different computations which have been kept in the database.

The display tree on the left shows results for a [qualitative + electrophoresis at threshold of 6] computation.

*Distinct cultivars [3]* demonstrates that 3 varieties were found distinct from all others. There was a total of 52 (49 + 3) cultivars in the computation.

The display tree is used to navigate through all possible pairs.

The user can expand or reduce the branches of the tree according to his needs.

*NON-distinct cultivars [49]*. Forty-nine cultivars were found “not distinct from all others” with a threshold of 6.

The first variety, *Variety 107*, has only 3 close varieties, whereas the second, *Variety 112*, has 9 close varieties, the third, *Variety 113*, 4 close varieties, etc.

The raw data for *Variety 112* and *Variety 26* are visible for the 6 qualitative characteristics observed on both varieties.

*Variety 112 [1][9]* indicates variety 112 is in the first year of examination [1]; and has 9 close varieties according to the threshold of 6 [9].



[*dist=3.5*]Variety 26 [2] indicates variety 26 has a GAIA distance of 3.5 from variety 112, which is in second year of examination.

The third column is the weighting according to the pre-defined matrices. The notes for both varieties are displayed for the two available locations (Std stands for “studied” which are the candidate varieties).

In this screen copy the varieties have been numbered for sake of confidentiality, the crop experts can name the varieties according to their need (lot or application number, name, etc.).

## 5.8 Final remark

The above example was described in order to explain how GAIA uses different types of characteristics in a practical case. The efficiency of the use of GAIA depends on the species.

## 5.9 References

Dagnelie Pierre. (1981). *Principes d'expérimentation*

Dagnelie Pierre. (1998). *Statistique théorique et appliquée volume 2 inférence statistique à une et deux dimensions*. Bibliothèque des universités Statistique

Digby, P.G.N. (1979). Modified joint regression analysis for incomplete variety x environment data. *J. Agric. Sci. Camb.* 93, 81-86.

Kala, R. (2002). *Statystyka dla przyrodników*, Agric. Univ. of Poznan

Mead, R., Curnow, R. N. and Hasted, R. M. (1993). *Statistical Methods in Agriculture and Experimental Biology*. Chapman & Hall, London.

Patterson, H. D. and Thompson, R. (1971). *Recovery of interblock information when block sizes are unequal*. *Biometrika*, 58, 545-554.

Patterson, H.D. & Weatherup, S.T.C. (1984). Statistical criteria for distinctness between varieties of herbage crops. *J. Agric. Sci. Camb.* 102, 59-68.

Sokal, R. R. and Rohlf, F. J. (1995). *Biometry*, W. H. Freeman Company

Talbot, M. (1990). Statistical aspects of minimum distances between varieties. UPOV TWC Paper TWC/VIII/9, UPOV, Geneva.

Yates, F. (1933). *The principles of orthogonality and confounding in replicated experiments*. *Journal of Agricultural Science, Cambridge*, 23, 108-45.

## 6. Examining DUS in bulk samples

### 6.1 Introduction and abstract

In some crops samples are bulked before certain characteristics are examined. The term “bulk sampling” is used here for the process of merging some or all individual plants before recording a characteristic. There are different degrees of bulking ranging from: 1) merging pairs of plants, 2) merging 3 or 4 up to all plants within a plot up to 3) merging all plants within a variety. The degree of bulking may play an important role in the efficiency of the tests. Bulking is usually only applied where the measurement of the characteristic is very expensive or very difficult to obtain for individual plants. Some examples are seed weight in cereals and peas and beans, and erucic acid content in rapeseed. This section describes some of the consequences of bulk sampling. It is shown that the test of distinctness (using COYD, see Part II: Section 2.1 [*cross ref.*]) may be expected to be relatively insensitive to the degree of bulking, but that the efficiency of the tests for uniformity (using COYU, see Part II: Section 2.2 [*cross ref.*]) must be expected to decrease when the data are bulked. The COYU test for uniformity cannot be carried out if all plants within a plot are bulked.

### 6.2 Distinctness

6.2.1 In the COYD method for examining distinctness the basic values to be used in the analyses are the annual variety means. As bulk sampling also gives at least one value for each variety per year, it will usually still be possible to use the COYD method for distinctness purposes for any degree of bulking, as long as at least one value is recorded for each variety in each year and that the bulk samples are representative for the variety. However, some problems may be foreseen: the assumption of data being normal distributed may be better fulfilled when the mean of many individual measurements are analyzed instead of the mean of fewer measurements or, in the extreme, just a single measurement.

6.2.2 The efficiency of the test of distinctness may be expected to be lower when based on bulked samples than when it is based on the mean of all individual plants in a year. The loss will be from almost zero upwards, depending on the importance of the different sources of variations. The variation which is relevant for the efficiency of variety comparisons is formulated in the following model.

$$\sigma_{total}^2 = \sigma_{vy}^2 + \sigma_p^2 + \sigma_i^2 + \sigma_m^2$$

where

$\sigma_{total}^2$  is the total variance of a characteristic used for comparing varieties

The total variance is regarded as being composed of four sources of variations:

- 1:  $\sigma_{vy}^2$  the year in which the variety is measured
- 2:  $\sigma_p^2$  the plot in which the measurement was taken
- 3:  $\sigma_i^2$  the plant on which the measurement was taken
- 4:  $\sigma_m^2$  the inaccuracy in the measurement process

6.2.3 In cases where the data are not bulked the variance of the difference between two variety means,  $\sigma_{diff}^2$ , becomes:

$$\sigma_{dif}^2 = 2 \left\{ \frac{\sigma_{vy}^2}{a} + \frac{\sigma_p^2}{ab} + \frac{\sigma_i^2}{abc} + \frac{\sigma_m^2}{abc} \right\}$$

where

$a$  is the number of years used in the COYD method

$b$  is the number of replicates in each trial

$c$  is the number of plants in each plot

6.2.4 Assuming that each bulk sample has been composed in such a way that it represents an equal amount of material from all the individual plants which have been bulked into that sample, the variance between two varieties based on  $k$  bulked samples (each of  $l$  plants) becomes:

$$\sigma_{dif}^2 = 2 \left\{ \frac{\sigma_{vy}^2}{a} + \frac{\sigma_p^2}{ab} + \frac{\sigma_i^2}{abkl} + \frac{\sigma_m^2}{abk} \right\}$$

where

$k$  is the number of bulk samples

$l$  is the number of plants in each bulk sample

6.2.5 Thus if all plants in each plot are divided in  $k$  groups of  $l$  plants each and an average measurement is taken for each of the  $k$  groups, then only the last term in the expression for  $\sigma_{dif}^2$  has increased (as  $kl$  is equal to  $c$ ). For many characteristics it is found that the variance caused by the measurements process is small and hence the bulking of samples will only have a minor effect on the conclusions reached by the COYD method. Only if the variance caused by the measurement process is relatively large can bulking have a substantial effect on the distinctness tests using COYD.

#### *Example 1*

*Variances for comparing varieties were estimated (by the use of estimated variance components) for different degrees of bulking. The calculations were based on the weight of 100 seeds of 145 pea varieties grown in Denmark during 1999 and 2000. In this example, the contribution to the variance caused by the measurement process was relatively very small, which means that bulking will have a low influence on the test for distinctness. In a 3 year test with 30 plants in each of 2 blocks, the variance of a difference between two varieties was estimated to be 2.133 and 2.135, for no bulking and a single bulk sample per plot, respectively.*

*For other variables the variance component due to the measurement process may be relatively more important. However, it is likely that in most practical cases this variance component will be relatively small.*

6.2.6 In some cases each bulk sample is not drawn from a specific set of plants (say, plant 1 to 5 in bulk sample 1, plant 6 to 10 in bulk sample 2 etc.), but bulk samples are formed from mixed samples of all plants in a plot. This means that different bulk samples may contain material from the same plants. It must be expected that similar results apply here, although, in this situation, the effect of bulking may have an increased effect because there is no guarantee that all plants will be equally represented in the bulk samples.

## 6.3 Uniformity

### 6.3.1 Bulking within plot

6.3.1.1 In COYU the test is based on the standard deviation of the individual plant observations (within plots) as a measurement of uniformity. The log of the standard deviations plus one are analyzed in an over-years analysis; i.e. the values  $Z_{vy} = \log(s_{vy} + 1)$  are used in the analyses. The variance on these  $Z_{vy}$  values can be regarded as arising from two sources, a component that depends on the variety-by-year interaction and a component that depends on the number of degree of freedom used for estimating the standard deviation,  $s_{vy}$  (the fewer degrees of freedom the more variable the standard deviation will be). This can be written (note that the same symbols as used in the distinctness section will be used here with different meaning):

$$Var(Z_{vy}) = \sigma_{vy}^2 + \sigma_f^2$$

where this variance can be regarded as being composed of two sources of variations:

- 1:  $\sigma_{vy}^2$  the year in which the variety is measured
- 2:  $\sigma_f^2$  the number of degrees of freedom used in estimating  $s_{vy}$

$\sigma_f^2$  is approximately  $\frac{1}{2\nu} \left( \frac{\sigma}{\sigma + 1} \right)^2$  when the recorded variable is normally distributed and the

standard deviations do not vary too much. This last expression reduces to  $0.5/\nu$  when  $\sigma \gg 1$ .

Here  $\sigma$  is the mean value of the  $s_{vy}$  values and  $\nu$  is the number of degrees of freedom used in the estimation of  $s_{vy}$ .

6.3.1.2 The variance caused by the year in which the variety is measured may be assumed to be independent of whether the samples are bulked or not, whereas the variance caused by the number of degrees of freedom will increase when bulked samples are used because a lower number of degrees of freedom is available.

6.3.1.3 The variance of a difference between a  $Z_{vy}$  for a candidate variety and the mean of the reference varieties'  $Z_{vy}$  values may be written:

$$\sigma_{dif}^2 = (\sigma_{vy}^2 + \sigma_f^2) \left( \frac{1}{a} + \frac{1}{ar} \right)$$

where

$a$  is the number of year used in the test

$r$  is the number of reference varieties

#### *Example 2*

*The effect of bulking in the test for uniformity, an estimate was made using the same data as for Example 1 I Part II, Section 6.2.5 [cross ref.]. For a test using 50 reference varieties in 3 years with 30 plants per variety in each of 2 plots per trial the variance for comparing the  $Z_{vy}$  value for a candidate variety and the mean of the reference varieties'  $Z_{vy}$  will be 0.0004 if no bulking is done. This can be compared to 0.0041, 0.0016 and 0.0007 when 2, 4 and 10 bulk samples per plot were used. Thus, in this example, the effect of bulking has a great influence on the test for uniformity. The variance increased,*

*approximately by a factor of 10 when changing from individual plant records to just 2 bulk samples per plot. This means that the degree of non-uniformity must be much higher for it to be detected when 2 bulk samples are used instead of individual plant records.*

### 6.3.2 Bulking across plots

Bulking across plots means that part of the between plot (and block) variation will be included in the estimated standard deviation between bulked samples. If this variation is relatively large it will tend to mask any differences in uniformity between varieties. In addition some noise may also be added because the ratio of material from the different plots may vary from bulk to bulk. Finally the assumptions for the present recommended method, COYU, may not be fulfilled in such cases. Therefore it is recommended to bulk only within plots.

### 6.3.3 Taking just one bulk sample per plot

In general, if all plants in a plot are bulked such that only a single sample is available for each plot, it becomes impossible to calculate the within plot variability and in such cases no tests for uniformity can be performed. In rare cases, where non-uniformity may be judged from values that can only be found in mixtures, non-uniformity may be detected even where a single bulk sample for each plot is used. For example, in the characteristic “erucic acid” in oil seed rape, values between 2% and 45% can only arise because of a lack of uniformity. However this only applies in certain special cases and even here the non-uniformity may only show up under certain circumstances.

[End of document]