



TWA/41/19

ORIGINAL: English

DATE: April 20, 2012

INTERNATIONAL UNION FOR THE PROTECTION OF NEW VARIETIES OF PLANTS

Geneva

TECHNICAL WORKING PARTY FOR AGRICULTURAL CROPS

Forty-First Session
Angers, France, May 21 to 25, 2012

COMMENTS CONCERNING THE DRAFT TEST GUIDELINES FOR WHEAT
(DOCUMENT TG/3/12(PROJ.1))

Document prepared by experts from France

The structure of the document is as follows:

- I DRAFT WITH COMMENTS RECEIVED BY THE SUBGROUP
- II GENERAL COMMENTS RECEIVED BY THE SUBGROUP



TG/3/12(proj.1)
ORIGINAL: English
DATE: 03.04.12

INTERNATIONAL UNION FOR THE PROTECTION OF NEW VARIETIES OF PLANTS
GENEVA

DRAFT

WHEAT
UPOV Code: TRITI_AES
Triticum aestivum L. emend. Fiori et Paol.

GUIDELINES
FOR THE CONDUCT OF TESTS
FOR DISTINCTNESS, UNIFORMITY AND STABILITY

prepared by experts from France

to be considered by the

*Technical Working Party for Agricultural Crops
at its 41st session, to be held in France from May 21 to 25, 2012*

Alternative Names:*

<i>Botanical name</i>	<i>English</i>	<i>French</i>	<i>German</i>	<i>Spanish</i>
Triticum aestivum L. emend. Fiori et Paol.	Wheat			

The purpose of these guidelines ("Test Guidelines") is to elaborate the principles contained in the General Introduction (document TG/1/3), and its associated TGP documents, into detailed practical guidance for the harmonized examination of distinctness, uniformity and stability (DUS) and, in particular, to identify appropriate characteristics for the examination of DUS and production of harmonized variety descriptions.

ASSOCIATED DOCUMENTS

These Test Guidelines should be read in conjunction with the General Introduction and its associated TGP documents.

* These names were correct at the time of the introduction of these Test Guidelines but may be revised or updated. [Readers are advised to consult the UPOV Code, which can be found on the UPOV Website (www.upov.int), for the latest information.]

<u>TABLE OF CONTENTS</u>	<u>PAGE</u>
1. SUBJECT OF THESE TEST GUIDELINES	4
2. MATERIAL REQUIRED	4
3. METHOD OF EXAMINATION.....	4
3.1 Number of Growing Cycles.....	4
3.2 Testing Place	4
3.3 Conditions for Conducting the Examination	4
3.4 Test Design.....	4
3.5 Additional Tests	5
4. ASSESSMENT OF DISTINCTNESS, UNIFORMITY AND STABILITY.....	5
4.1 Distinctness	5
4.2 Uniformity.....	6
4.3 Stability	7
5. GROUPING OF VARIETIES AND ORGANIZATION OF THE GROWING TRIAL.....	7
6. INTRODUCTION TO THE TABLE OF CHARACTERISTICS	8
6.1 Categories of Characteristics	8
6.2 States of Expression and Corresponding Notes	8
6.3 Types of Expression	8
6.4 Example Varieties.....	8
6.5 Legend.....	9
7. TABLE OF CHARACTERISTICS/TABLEAU DES CARACTÈRES/MERKMALSTABELLE/TABLA DE CARACTERES.....	10
8. EXPLANATIONS ON THE TABLE OF CHARACTERISTICS	16
8.1 Explanations covering several characteristics.....	16
8.2 Explanations for individual characteristics.....	16
The descriptions of the growth stages of the Zadoks decimal code for cereals	25
9. LITERATURE.....	26
10. TECHNICAL QUESTIONNAIRE.....	27

1. Subject of these Test Guidelines

These Test Guidelines apply to all varieties of *Triticum aestivum* L. emend. Fiori et Paol.

2. Material Required

2.1 The competent authorities decide on the quantity and quality of the plant material required for testing the variety and when and where it is to be delivered. Applicants submitting material from a State other than that in which the testing takes place must ensure that all customs formalities and phytosanitary requirements are complied with.

2.2 The material is to be supplied in the form of seed.

2.3 The minimum quantity of plant material, to be supplied by the applicant, should be:

Seeds: 5 kg
Ears (if requested): 100

Comment [G1]:
DK suggest 3 kg

Comment [G2]:
UK proposes 3kg

Comment [G3]:
RO: The minim quantity of plant material, to be supplied by the applicant.
Ear (for the winter varieties) 150

The seed should meet the minimum requirements for germination, species and analytical purity, health and moisture content, specified by the competent authority. In cases where the seed is to be stored, the germination capacity should be as high as possible and should, be stated by the applicant.

If ears are requested, they should contain a sufficient number of viable seeds to establish a satisfactory row of plants for observation.

Comment [G4]:
UK: If ear row samples are requested, they should

2.4 The plant material supplied should be visibly healthy, not lacking in vigor, nor affected by any important pest or disease.

2.5 The plant material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If it has been treated, full details of the treatment must be given.

3. Method of Examination

3.1 *Number of Growing Cycles*

The minimum duration of tests should normally be two independent growing cycles.

3.2 *Testing Place*

Tests are normally conducted at one place. In the case of tests conducted at more than one place, guidance is provided in TGP/9 "Examining Distinctness".

3.3 *Conditions for Conducting the Examination*

3.3.1 The tests should be carried out under conditions ensuring satisfactory growth for the expression of the relevant characteristics of the variety and for the conduct of the examination.

3.3.2 Stage of development for the assessment

The optimum stage of development for the assessment of each characteristic is indicated by a number in the second column of the Table of Characteristics. The stages of development denoted by each number are described in the descriptions of the growth stages of the Zadoks decimal code for cereals at the end of the Chapter 8.

3.4 *Test Design*

3.4.1 Each test should be designed to result in a total of at least 2000 plants. The assessment for the characteristics "Seasonal type" should be carried out on at least 300 plants.

Comment [G5]:
DE: According to the CPVO-TP 500 plants. Was it reduced by intention?

3.4.2 The design of the tests should be such that plants or parts of plants may be removed for measurement or counting without prejudice to the observations which must be made up to the end of the growing cycle.

3.4.3 Single ear rows: if tests on ear rows are conducted, at least 100 ear rows should be observed.

3.4.4 In case of hybrids, the parent lines have to be included in the test and should be tested and assessed as any other self-pollinating variety. The observations on the hybrid variety itself should be made on at least 200 plants.

3.5 *Additional Tests*

Additional tests, for examining relevant characteristics, may be established.

4. Assessment of Distinctness, Uniformity and Stability

4.1 *Distinctness*

4.1.1 General Recommendations

It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding distinctness. However, the following points are provided for elaboration or emphasis in these Test Guidelines.

To assess distinctness of hybrids, the parent lines and the formula may be used according to the following recommendations:

- (i) description of parent lines according to the Test Guidelines;
- (ii) check of the originality of the parent lines in comparison with the variety collection, based on the characteristics in Chapter 7, in order to identify similar parent lines;
- (iii) check of the originality of the hybrid formula in relation to the hybrids in the variety collection, taking into account the most similar lines; and
- (iv) assessment of the distinctness at the hybrid level for varieties with a similar formula.

Further guidance is provided in documents TGP/9 "Examining Distinctness" and TGP/8 "Trial Design and Techniques Used in the Examination of Distinctness, Uniformity and Stability".

4.1.2 Consistent Differences

The differences observed between varieties may be so clear that more than one growing cycle is not necessary. In addition, in some circumstances, the influence of the environment is not such that more than a single growing cycle is required to provide assurance that the differences observed between varieties are sufficiently consistent. One means of ensuring that a difference in a characteristic, observed in a growing trial, is sufficiently consistent is to examine the characteristic in at least two independent growing cycles.

4.1.3 Clear Differences

Determining whether a difference between two varieties is clear depends on many factors, and should consider, in particular, the type of expression of the characteristic being examined, i.e. whether it is expressed in a qualitative, quantitative, or pseudo-qualitative manner. Therefore, it is important that users of these Test Guidelines are familiar with the recommendations contained in the General Introduction prior to making decisions regarding distinctness.

4.1.4 Number of Plants / Parts of Plants to be Examined

Unless otherwise indicated, for the purposes of distinctness, all observations on single plants should be made on 20 plants or parts taken from each of 20 plants and any other observations made on all plants in the test, disregarding any off-type plants.

In the case of observations of parts taken from single plants, the number of parts to be taken from each of the plants should be 1.

4.1.5 Method of Observation

The recommended method of observing the characteristic for the purposes of distinctness is indicated by the following key in the second column of the Table of Characteristics (see document TGP/9 "Examining Distinctness", Section 4 "Observation of characteristics"):

- MG: single measurement of a group of plants or parts of plants
- MS: measurement of a number of individual plants or parts of plants
- VG: visual assessment by a single observation of a group of plants or parts of plants
- VS: visual assessment by observation of individual plants or parts of plants

Type of observation: visual (V) or measurement (M)

"Visual" observation (V) is an observation made on the basis of the expert's judgment. For the purposes of this document, "visual" observation refers to the sensory observations of the experts and, therefore, also includes smell, taste and touch. Visual observation includes observations where the expert uses reference points (e.g. diagrams, example varieties, side-by-side comparison) or non-linear charts (e.g. color charts). Measurement (M) is an objective observation against a calibrated, linear scale e.g. using a ruler, weighing scales, colorimeter, dates, counts, etc.

Type of record: for a group of plants (G) or for single, individual plants (S)

For the purposes of distinctness, observations may be recorded as a single record for a group of plants or parts of plants (G), or may be recorded as records for a number of single, individual plants or parts of plants (S). In most cases, "G" provides a single record per variety and it is not possible or necessary to apply statistical methods in a plant-by-plant analysis for the assessment of distinctness."

In cases where more than one method of observing the characteristic is indicated in the Table of Characteristics (e.g. VG/MG), guidance on selecting an appropriate method is provided in document TGP/9, Section 4.2.

4.2 Uniformity

4.2.1 It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding uniformity. However, the following points are provided for elaboration or emphasis in these Test Guidelines:

4.2.2 Indication of sample size in the Table of Characteristics

The recommended sample size for the assessment of uniformity is indicated by the following key in the table of characteristics:

- {A} sample size of 100 plants/parts of plants/ear-rows
- {B} sample size of 2000 plants or parts of plants

4.2.3 Uniformity assessment on all plants in the test

For the assessment of uniformity in a sample of 2000 plants, a population standard of 0.3% and an acceptance probability of at least 95% should be applied. In the case of a sample size of 2000 plants, 10 off-types are allowed.

4.2.4 Uniformity assessment on a sub-sample

For the assessment of uniformity in a sample of 100 ear-rows, plants or parts of plants, a population standard of 1% and an acceptance probability of at least 95% should be applied. In the case of a sample size of 100 ear-rows, plants or parts of plants, 3 off-types are allowed.

An ear-row is considered to be an off-type ear-row if there is more than one off-type plant within that ear-row.

Comment [G6]:
DE Headings are for explanation in the template. Not necessary for the TG.

Comment [G7]:
DE proposes to delete this subtitle

Comment [G8]:
DE proposes to delete this subtitle

For “A” characteristics, with the exception of characteristic 1 and 2, the assessment of uniformity can be done in 2 steps. In a first step, 20 plants are observed. If no off-types are observed, the variety is declared to be uniform. If more than 3 off-types are observed, the variety is declared not to be uniform. If 1 to 3 off-types are observed, an additional sample of 80 plants or parts of plants must be observed.

4.2.6 Uniformity assessment where the parent formula is used

Where the assessment of a hybrid variety involves the parent lines, the uniformity of the hybrid variety should, in addition to an examination of the hybrid variety itself, also be assessed by examination of the uniformity of its parent lines.

For the assessment of uniformity of hybrids, a population standard of 10% and an acceptance probability of at least 95% should be applied. In the case of a sample size of 200 plants, 27 off-types are allowed.

Comment [G9]:
DE proposes to delete this subtitle

Comment [G10]:
UK: What are the standards for 3 way hybrids?

4.3 *Stability*

4.3.1 In practice, it is not usual to perform tests of stability that produce results as certain as those of the testing of distinctness and uniformity. However, experience has demonstrated that, for many types of variety, when a variety has been shown to be uniform, it can also be considered to be stable.

4.3.2 Where appropriate, or in cases of doubt, stability may be further examined by testing a new seed stock to ensure that it exhibits the same characteristics as those shown by the initial material supplied.

4.3.3 Stability assessment: hybrid varieties

Where appropriate, or in cases of doubt, the stability of a hybrid variety may, in addition to an examination of the hybrid variety itself, also be assessed by examination of the uniformity and stability of its parent lines.

5. Grouping of Varieties and Organization of the Growing Trial

5.1 The selection of varieties of common knowledge to be grown in the trial with the candidate varieties and the way in which these varieties are divided into groups to facilitate the assessment of distinctness are aided by the use of grouping characteristics.

5.2 Grouping characteristics are those in which the documented states of expression, even where produced at different locations, can be used, either individually or in combination with other such characteristics: (a) to select varieties of common knowledge that can be excluded from the growing trial used for examination of distinctness; and (b) to organize the growing trial so that similar varieties are grouped together.

5.3 The following have been agreed as useful grouping characteristics:

- (a) Straw: pith in cross section (half way between base of ear and stem node below) (characteristic 11)
- (b) Awns or scurs : presence (characteristic 15)
- (c) Ear : color (characteristic 17)
- (d) Seasonal type (characteristic 25)

Comment [G11]:
UK proposes to delete Straw: pith in cross section as a grouping characteristic
FR agrees

5.4 An alternative method to grouping characteristics is GAIA method. It has been developed to optimize trials, by avoiding the growing of some of the varieties in the variety collection. The principle is to compute a phenotypic distance between each pair of varieties, this distance being a sum of distances on each individual observed characteristic. The background of the method relies on the possibility given to the crop expert to express his confidence on the differences observed, by giving weights to the difference for each observed characteristic.

GAIA method is mainly used after a first growing cycle to identify those varieties in the variety collection which can be excluded from the subsequent growing cycle(s) because they are “Distinct Plus” (see TGP/8/1 Part II section 1.3.2.1) from all the candidate varieties. GAIA can also identify similar varieties, on which the DUS examiner will need to focus attention in the subsequent growing cycle.

Comment [G12]:
UK proposes that GAIA is not referred to specifically here, but that standard terms for databases that compare varieties is used.
DE: The method and the use of GAIA is explained in TGP/8 and TGP/9. I am not in favor to repeat such specific methods in individual TGs (we would also not do this for COY etc.).
I understand that GAIA is not an alternative to grouping but complements the process for D.

5.5 Guidance for the use of grouping characteristics, in the process of examining distinctness, is provided through the General Introduction and document TGP/9 “Examining Distinctness”.

6. Introduction to the Table of Characteristics

6.1 *Categories of Characteristics*

6.1.1 Standard Test Guidelines Characteristics

Standard Test Guidelines characteristics are those which are approved by UPOV for examination of DUS and from which members of the Union can select those suitable for their particular circumstances.

6.1.2 Asterisked Characteristics

Asterisked characteristics (denoted by *) are those included in the Test Guidelines which are important for the international harmonization of variety descriptions and should always be examined for DUS and included in the variety description by all members of the Union, except when the state of expression of a preceding characteristic or regional environmental conditions render this inappropriate.

6.2 *States of Expression and Corresponding Notes*

6.2.1 States of expression are given for each characteristic to define the characteristic and to harmonize descriptions. Each state of expression is allocated a corresponding numerical note for ease of recording of data and for the production and exchange of the description.

6.2.2 In the case of qualitative and pseudo-qualitative characteristics (see Chapter 6.3), all relevant states of expression are presented in the characteristic. However, in the case of quantitative characteristics with 5 or more states, an abbreviated scale may be used to minimize the size of the Table of Characteristics. For example, in the case of a quantitative characteristic with 9 states, the presentation of states of expression in the Test Guidelines may be abbreviated as follows:

State	Note
small	3
medium	5
large	7

However, it should be noted that all of the following 9 states of expression exist to describe varieties and should be used as appropriate:

State	Note
very small	1
very small to small	2
small	3
small to medium	4
medium	5
medium to large	6
large	7
large to very large	8
very large	9

6.2.3 Further explanation of the presentation of states of expression and notes is provided in document TGP/7 "Development of Test Guidelines".

6.3 *Types of Expression*

An explanation of the types of expression of characteristics (qualitative, quantitative and pseudo-qualitative) is provided in the General Introduction.

6.4 Example Varieties

Where appropriate, example varieties are provided to clarify the states of expression of each characteristic.

Comment [G13]:
UK: Example varieties should be more geographically diverse and relevant to all countries. Some varieties listed here (Cadenza) are very old. UK suggests using only photographs instead of example varieties as in revision of TGP/7 Example varieties.

6.5 *Legend*

(*) Asterisk characteristic – see Chapter 6.1.2

QL Qualitative characteristic – see Chapter 6.3

QN Quantitative characteristic – see Chapter 6.3

PQ Pseudo-qualitative characteristic – see Chapter 6.3

MG, MS, VG, VS – see Chapter 4.1.5

A, B _ see Chapter 4.2

(a) See Explanations on the Table of Characteristics in Chapter 8.1

(+) See Explanations on the Table of Characteristics in Chapter 8.2.

7. Table of Characteristics/Tableau des caractères/Merkmalstabelle/Tabla de caracteres

	English	français	deutsch	español	Example Varieties Exemples Beispielsorten Variedades ejemplo	Not No
1. (+)	00 VG	Seed: coloration with phenol	Semence : coloration au phenol	Samen: Phenolfärbung	Semilla: color con phenol	
QN	A	None or very light	Nulle ou très faible	Fehlend oder sehr hell		1
		light	Faible	Hell	claro	Soissons 3
		medium	moyenne	mittel	medio	Aerobic 5
		dark	Forte	Dunkel	Oscuro	Excelcior 7
		very dark	Très foncée	Sehr dunkel	Muy oscuro	Sideral 9
2. (+)	09-11 VG	Coleoptile: anthocyanin coloration	Coléoptile: pigmentation anthocyanique	Keimscheide: Anthocyanfärbung	Coleóptilo: pigmentación antocianica	
QN	A	absent or very weak	absente ou très faible	fehlend oder sehr gering	ausente o muy débil	Altigo 1
		weak	Faible	Gering	débil	Accor 3
		medium	Moyenne	Mittel	media	Premio 5
		strong	Forte	Stark	Fuerte	Arezzo 7
		very strong	Très forte	Sehr stark	Muy fuerte	Caphorn 9
3. (*)	25-29 VG	Plant: growth habit	Plante: port au tallage	Pflanze: Wuchsform	Planta: porte	
(+)	B	erect	dressé	aufrecht	erecto	Bandera 1
QN		semi erect	demi-dressé	halbaufrecht	semierecto	Esperia 3
		intermediate	demi-dressé à demi-étalé	mittel	medio	Crousty
		semi prostrate	demi-étalé	halbliiegend	semiprostrado	Euclide
		prostrate	étalé	liegend	postrado	Instinct
4. (+)	47-51 VG	Plant: frequency of plants with recurved flag leaves	Plante : fréquence de plantes avec la dernière feuille retombante	Pflanze: Häufigkeit von Pflanzen mit gebogenen obersten Blättern	Planta: frecuencia de plantas con ¿	
QN	B	absent or very low	nulle ou très faible	fehlend oder sehr gering	ausente o muy débil	Sorrial
		low	faible	gering	débil	Arezzo 3
		medium	moyenne	mittel	media	Courtot 5
		high	forte	stark	fuerte	Saturnus 7
		very high	très forte	sehr stark	muy fuerte	MV kolo 9

Comment [G14]:
CZ – to check example, in CPVO file is 3, in CZ is 4

Comment [G15]:
PL: proposes intermediate instead of medium
CA would like to propose the state 'intermediate' (as opposed to medium) for the third state of expression in characteristic 3: plant: growth habit CL:Intermediate

FR:(this proposal has already been introduced in the draft 1

Comment [G16]:
UK : What are the experiences of other countries? Are all these states necessary?

	English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Not No
5. (*)	50-52 MG	Time of ear emergence	Époque d'épiaison	Zeitpunkt des Ährenschiebens	Época de ¿	
(+)						
QN	B	very early	très précoce	sehr früh	muy temprana	Accor 1
		early	précoce	früh	temprana	Caphorn 3
		medium	moyenne	mittel	media	Richepain 5
		late	tardive	spät	tardía	Boncap 7
		very late	très tardive	sehr spät	muy tardía	9
6. (*)	60-65 VG	Flag leaf: glaucosity of sheath	Dernière feuille : glaucescence de la gaine	Oberstes Blatt: Bereifung der Blattscheide	Última hoja: ¿ de la vaina	
QN	B	absent or very weak	absente ou très faible	fehlend oder sehr gering	ausente o muy débil	Benedict 1
		weak	faible	gering	débil	Aerobic 3
		medium	moyenne	mittel	medio	Pakito 5
		strong	forte	stark	fuerte	Solehio 7
		very strong	très forte	sehr stark	muy fuerte	Illico 9
7.	60-65 VG	Flag leaf: glaucosity of blade (lower side)	Dernière feuille : glaucescence du limbe (face inférieure)	Oberste Blatt: Bereifung der Blattspreite	Última hoja: ¿ del limbo	
QN	B	absent or very weak	absente ou très faible	fehlend oder sehr gering	ausente o muy débil	Courtot 1
		weak	faible	gering	débil	Bologna 3
		medium	moyenne	mittel	medio	Amador 5
		strong	forte	stark	fuerte	Cezanne 7
		very strong	très forte	sehr stark	muy fuerte	Goncourt 9
8. (*)	60-69 VG	Ear: glaucosity	Epi : glaucescence	Aehre: Bereifung	¿: ¿	
QN	B	absent or very weak	absente ou très faible	fehlend oder sehr gering	ausente o muy débil	Soissons 1
		weak	faible	gering	débil	Bologna 3
		medium	moyenne	mittel	medio	Solehio 5
		strong	forte	stark	fuerte	Premio 7
		very strong	très forte	sehr stark	muy fuerte	Exelcior 9

	English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Not No
9.	60-69 VG	Culm: glaucosity of neck	Tige : glaucescence du col de l'épi	Halm: Bereifung des obersten Internodiums	ζ: ζ	
QN	B	absent or very weak	absente ou très faible	fehlend oder sehr gering	ausente o muy débil	Benedict 1
		weak	faible	gering	débil	Saturnus 3
		medium	moyenne	mittel	medio	Aubusson 5
		strong	forte	stark	fuerte	Arezzo 7
		very strong	très forte	sehr stark	muy fuerte	Exelcior 9
10.	75-92	Plant: length	Plante: longueur	Pflanze: Länge	Planta: altura (?)	
(*)	MG	very short	très courte	sehr niedrig	muy corta	Cordiale
						Comment [G17]: DK – change to VG/MG
(+)	B	short	courte	niedrig	corta	Renan 3
QN		medium	moyenne	mittel	media	Intérêt 5
		long / tall	longue / haute	lang / hoch	larga	Bagatelle 007 7
		very long / very tall	très longue / très haute	sehr lang / sehr hoch	muy larga	9
11.	80-92 (+) VG	Straw: pith in cross section	Paille : moëlle en section transversale	Halm: Füllung im Querschnitt	Tallo: grosor	
PQ	A	absent or very thin	Absente ou très mince	fehlend oder sehr dünn	Ausente o muy delgado	Pakito 1
		medium	moyenne	mittel	medio	Saturnus 2
		very thick or filled	Très épaisse ou pleine	Sehr dick oder ?	Muy grueso o ζ	Aerobic 3
12.	92	Ear: shape in profile	Epi: forme en vue de profil	Aehre : form in Seiten-ansicht		
(+)	VG					Comment [G20]: DE: To keep old order!
PQ	B	tapering	Pyramidal	Pyramiden-förmig		Sankara
		fusiform	Fusiforme	Spindle-förmig		Arezzo
		parallel sided	À bords parallèles	Parallel		Viscount
		semi-clavate	En demi-massue	Halb keulenförmig		Aura
		clavate	En massue	keulenförmig		Apache
						Comment [G21]: UK : What are the experiences of other countries? Are all these states necessary?
						Comment [G18]: CZ to avoid changes in dtb we propose to keep an original numbering, PQ char. allows it, to check AREZO in CPVO file is parallel
						Comment [G19]: JP: I think that judgment of difference between Note 4 and Note 5 is difficult. To be delete Note 4.

	English	français	deutsch	español	Example Varieties Exemples Beispielsorten Variedades ejemplo	Not No	
13 (*) (+)	80-92 Ear: density	Epi: compacité	Aehr: Dichte				
QN	VG,B very lax or MS, A lax	Très lâche	Sehr locker		Magno	1	
	medium	Demi-lâche à demi-compact	Mittel		Sponsor	Comment [G22]: CZ - MG should be deleted. If we want to follow an explanation, MS and A sample should be added as the second option FR proposes to put VG, B and MS, A DE: B is correct. We do not agree with CZ to add A.	
	dense	Compact	Dicht		Aubusson		
	very dense	Très compact	Sehr dicht		Premio		
					Rehti		9
14 (+)	80-92 Ear: length	Epi: longueur	Aehre : länge				
QN	VG very short	Très court	Sehr kurz			Comment [G23]: JP: Char. 14 have example varieties. So Ad.14 to be deleted. Char. 14 is QN characteristic. These pictures are not necessary.	
	B short	Court	Kurz		Ambello		Comment [G24]: DE proposes to add MS.
	medium	Moyen	Mittel		Soissons		Comment [G25]: PL: proposes VG/MG ... [1]
	long	Long	Lang		Aubusson		Comment [G26]: DK – change to VG/MG
	very long	Très long	Sehr lang		Folklor		
15. (*) (+)	80-92 Awns or scurs: presence	Barbes ou arêtes: présence	Grannen oder Spelzen-spitzen: Vorhandensein				
QL	VG both absent	Toutes les deux absentes	Beide fehlend		Genoveva	1	
	B scurs present	Arêtes présentes	Spelzenspitzen vorhanden		Aubusson	2	
	awns present	Barbes présentes	Grannen vorhanden		Arezzo	3	
16. (*) (+)	80-92 Awns or scurs at tip of ear: length	Barbes ou arêtes à l'extrémité de l'épi: longueur	Grannen oder Spelzen-spitzen an der Aehrenspitze : länge				
QN	VG very short	Très courtes	Sehr kurz		Ephoros	Comment [G27]: JP: Char. 16 have example varieties.	
	B short	Courtes	Kurz		Graindor		Comment [G28]: DE proposes to add MS
	medium	Moyennes	Mittel		Pakito		3
	long	Longues	Lang		As de cœur		5
	very long	Très longues	Sehr lang		Arezzo		7
						9	

	English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Not No
17.	80-92	Ear: color	Epi: couleur	Aehre: Farbe		
(*) (+)						
QL	VG	white	Blanc	Weiss	Arezzo	1
	B	colored	Coloré	gefärbt	Segor	2
18.	80-92	Apical rachis segment: extend of hairiness of convex surface	Article terminal du rachis: étendue de la pilosité de la face externe	Oberstes Spindelglied : äussere Behaarung		
(+)						
QN	VG	absent or very weak	Nulle ou très faible	Fehlend oder sehr gering	Graindor	
	A	weak	Faible	Gering	Crousty	
		medium	Moyenne	Mittel	Sirtaki	
		strong	Forte	Stark	Cadenza	7
		very strong	Très forte	Sehr stark	KWS Scirocco	9
19.	80-92	Lower glume: shoulder width	Glume inférieure: largeur de la troncature	Hüllspelze : Schulterbreite		
(+)						
QN	VG	absent or very narrow	Nulle ou très étroite	Fehlend oder sehr schmal	Courtot	1
	A	narrow	Étroite	Schmal	Altigo	3
	(a)	medium	Moyenne	Mittel	Apache	5
		broad	Large	Breit	Orvantis	7
		very broad	Très large	Sehr breit	Aglika	9
20.	80-92	Lower glume: shoulder shape	Glume inférieure: forme de la troncature	Hüllspelze : chulterform		
(+)						
QN	VG	sloping	Inclinée	Abfallend	Ambello	1
	A	slightly sloping	Légèrement inclinée	Leicht abfallend	Soissons	3
	(a)	straight	Droite	Gerade	Apache	5
		elevated	Echancrée	Gehoben	Aubusson	7
		strongly elevated with 2 nd point present	Fortement échancrée avec présence d'un 2ème bec	Stark gehoben mit vorhandener zweiter spitze	Fiorenzo	9

Comment [G29]:
CA proposes to amend the wording for characteristics 18 and 23 to make them consistent with one another. In both characteristics, it appears that it is the extent of the hairiness on the surfaces that is being observed
FR has modified the draft according to this proposal

	English	français	deutsch	español	Example Varieties Exemples Beispielsorten Variedades ejemplo	Not No
21.	80-92	Lower glume: beak length	Glume inférieure: longueur du bec	Hüllspelze : Zahnlänge		
(+)						
QN	VG	very short	Très court	Sehr kurz	Graindor	1
	A	short	Court	Kurz	Sollario	3
	(a)	medium	Moyen	Mittel	Quality	5
		long	Long	Lang	Soissons	7
		very long	Très long	Sehr lang	Bandera	
22.	80-92	Lower glume: beak shape	Glume inférieure: forme du bec	Hüllspelze : zahnform		
(*) (+)						
QN	VG	straight	Droit	Gerade	Premio	1
	A	slightly curved	Légèrement coudé	Leicht gebogen	Altigo	3
	(a)	moderately curved	Demi-coudé	Mittel gebogen	Sponsor	5
		strongly curved	Fortement coudé	Stark gebogen	Quebon	7
		geniculate	Genouillé	geknickt	Velocity	9
23.	80-92	Lower glume: extent of hairiness of internal surface	Glume inférieure: étendue de la pilosité de la face interne	Hüllspelze : verbreitung der inneren behaarung		
(*) (+)						
QN	VG	weak	Faible	Gering	Altigo	
	A	medium	Moyenne	Mittel	Alixan	
	(a)	strong	Forte	stark	Quality	
24.	92	Grain: color	Graine: couleur	Korn: farbe		
(+)						
QL	VG	white	Blanc	Weiss	Recital	
	A	red	Roux	rot	Apache	
25.	-	Seasonal type	Type de développement	Wechselverhalten		
(+) (*)						
PQ	VG	winter type	Type hiver	Winterform	Aubusson	
		alternative type	Type alternatif	Wechselform	Cezanne	
		spring type	Type printemps	sommerform	Josselin	

Comment [G30]:
CZ – to check example, in CPVO file is 5, in CZ is 6

Comment [G31]:
CA proposes to amend the wording for characteristics 18 and 23 to make them consistent with one another. In both characteristics, it appears that it is the extent of the hairiness on the surfaces that is being observed
FR has modified the draft according to this proposal

Comment [G32]:
JP: I would like to propose to delete (*).

Comment [G33]:
DE: To keep as it is. We do not agree with the comment from CZ.

Comment [G34]:
DK – assess the characteristic on the submitted seed? After 1 hour in a NaOH solution the expression White or Red on the grain are clear

Comment [G35]:
CZ – to change characteristic completely as follows:
Grain: coloration, light 1, medium 3, dark 5, QN instead of QL
There is no clear border between color

8. Explanations on the Table of Characteristics

8.1 Explanations covering several characteristics

Characteristics containing the following key in the second column of the Table of Characteristics should be examined as indicated below:

(a) Characteristics on lower glume must be observed at midthird of ear spikelet

8.2 Explanations for individual characteristics

Ad 1 : Grain : coloration with phenol

Method for Determination of Phenol Reaction

Number of grains per test	100 grains. The grains should not have been treated chemically
Concentration of solution	1 per cent Phenol-solution (freshly made up)
Preparation of grains	Soak grains in phenol solution during 4 hours with a constant move. Rinse out 4 or 5 times with clean water. Drain in half-light during at least 10 hours (reaction occurs in contact with air). Whole grain surface must be in contact with phenol solution.
Amount of solution	Laboratory
Place	Half-light, out of direct sunshine
Light	15 to 25°C
Temperature	At least 10 hours after rinsing
Time of recording	At least two of the example varieties should be included as a control
Note	

Comment [G36]:
DE: To keep as in CPVO-TP. We do not support any change of the method.

Comment [G37]:
CZ – The method works very well, there no reason for change, we are opinion that especially soaking before phenol test, which is missing here, is very important

Comment [G38]:
UK uses a different method for Grain: coloration of phenol.

Comment [G39]:
- BR agrees with CZ comments about the importance of soaking the seed for phenol reaction examination

Comment [G40]:
DK - 10 hours is not very practical if you have a 8 hours day of work - may add at least 10 hours after rinsing. The old less time consuming procedure for Grain phenol coloration works very well for us.

Comment [G41]:
UK: States 5,7 and 9 are difficult to see from this example photograph. UK proposes providing clearer photographs showing the states more clearly.


Comment [G42]:
FR: the pictures have been improved

Comment [G43]:
De: To keep as in CPVO-TP. We do not support any change of the method..

Comment [G44]:
CZ - The method works well, there is more flexible as regards light and temperature, there is no reason for change

Comment [G45]:
DK The old procedure for Coleoptile: anthocyanin coloration works very well for us and was more flexible.


Comment [G46]:
DK - 10 hours is not very practical if you have a 8 hours day of work - may add at least 10 hours after rinsing. The old less time consuming procedure for Grain phenol coloration works very well for us



Ad 2 : Coleoptile : anthocyanin coloration


Method for the Determination of Anthocyanin Coloration

Number of grains per test:	100 grains
Preparation of grains:	Set up non-dormant grains on moistened filter paper covered with a Petri dish lid during germination
Place:	Climate room with controlled temperature
Light:	Since the beginning of the test, 12h day / 12h night. Light intensity: 12000-15000 lux.
Temperature:	19°C
Time of recording:	Coleoptiles fully developed (about 10 days) at stage 09-11
Note:	At least two of the example varieties should be included as a control when testing for distinctness



Comment [G47]:
FR: these pictures have been improved





Ad 3 : Plant : growth habit



The growth habit should be assessed visually from the attitude of the leaves and tillers. The angle formed by the outer leaves and the tillers with an imaginary vertical axis should be used.






Le port doit être déterminé visuellement d'après le port des feuilles et des tiges. On utilisera l'angle formé par les feuilles externes et les tiges avec un axe vertical imaginaire.

Die Wuchsform sollte auf Grund der Haltung der Blätter und Triebe visuell erfasst werden. Der von den äusseren Blättern und Trieben mit einer vertikalen Achse gebildete Winkel sollte verwendet werden.

				
1	3	5	7	9
Erect	Semi-erect	Intermediate	Semi-prostrate	Prostrate

Ad 4 : Plant : frequency of plants with recurved flag leaves

- 1. all flag leaves are rectilinear
- 3 about 1/4 of the plants with recurved flag leaves
- 5 about 1/2 of the plants with recurved flag leaves
- 7 about 3/4 of the plants with recurved flag leaves
- 9 all flag leaves are recurved

				
1	3	5	7	9
Absent or very low	Low	Medium	High	Very high

Ad 5 : Time of ear emergence




Time of ear emergence should be scored when the first spikelet is visible on 50% of ears.

Ad 10 : Plant : length

The length of plant includes stem, ear, awns and scurs.











Ad 11 : Straw : pith in cross section

Pith in cross section should be observed halfway between base of ear and stem node below.

		
1	2	3
Absent or very thin	Medium	Very thick or filled

Ad 12 : Ear : shape in profile



Comment [G48]:
DE: To keep old order.

				
				
1	2	3	4	5
Tapering	Fusiform	Parallel sided	Semi-clavate	Clavate

Ad 13 : Ear : density

The density can be assessed either visually or as measurement of the ratio of the number of spikelets/ear length.






Comment [G49]:
CZ – measurement should be done from “A” sample, see our explanation above

				
1	3	5	7	9
Very lax	Lax	Medium	Dense	Very dense

Ad 14 : Ear : length




Length of ear should be **observed** excluding awns and scurs.

Comment [G50]:
DK - or measured

				
1	3	5	7	9
Very short	Short	Medium	Long	Very long

Ad 15 : Awns or scurs : presence






Comment [G51]:
UK proposes providing clearer photographs of ear characteristics.

		
1	2	3
Both absent	Scurs present	Awns present

Comment [G52]:
FR: pictures have been replaced by drawings

Ad 16 : Awns or scurs at tip of ear : length

Comment [G53]:
JP: Ad.16 should be indicated part for measurement or observation.






				
1	3	5	7	9
Very short	Short	Medium	Long	Very long

Comment [G54]:
FR: pictures have been replaced by drawings






Ad 17 : Ear : colour

	
1	2
White	Coloured

Ad 18 : Apical rachis segment : hairiness of convex surface

				
1	3	5	7	9
Absent or very weak	Weak	Medium	Strong	Very strong






Ad 19 : Lower glume : shoulder width

				
1	3	5	7	9
Absent or very narrow	Narrow	Medium	Broad	Very broad

Comment [G55]:
UK: The illustrations here exaggerate the expression of the lower glume: shoulder width compared to the diagram in the previous TG.






Comment [G56]:
FR: drawings have been improved

Ad 20 : Lower glume : shoulder shape






				
1	3	5	7	9
Sloping	Slightly sloping	Straight	Elevated	Strongly elevated with 2 nd point present

Comment [G57]:
FR: drawings have been improved

Ad 21 : Lower glume : beak length

				
1	3	5	7	9
Very short	Short	Medium	Long	Very long




Ad 22 : Lower glume : beak shape

				
1	3	5	7	9
Straight	Slightly curved	Moderately curved	Strongly curved	Geniculate



Comment [G58]:
CZ – new drawing is not correspond with old one – current 9 is 7 new one, current 7 is 5 new one.. Is the drawing really accordance with a real situation. If yes, current descriptions should be reclassified

Comment [G59]:
FR: drawings have been improved

Ad 23 : Lower glume : extent of internal hair

		
1	3	5
Weak	Medium	Strong

Ad 24 : Grain colour

	
1	2
White	Red

Ad 25 : Seasonal type

The seasonal type should be assessed on one or several plots sown in springtime. Example varieties should always be included in the plots. When the example varieties behave according to its description, the varieties under study can be described. At the time when the latest springtype variety is fully mature (stage 91/92 of the Zadoks decimal code), the growth stage reached by the respective variety should be assessed. The states of expression are defined as follows:

Winter type: the plants have reached stage 45 of the Zadoks decimal code (boots swollen) at maximum

Alternative type: the plants have exceeded stage 45 of the Zadoks decimal code---as a rule they have exceeded stage 75---and have reached stage 90 at maximum

Spring type: the plants have exceeded stage 90 of the Zadoks decimal code.

Comment [G60]:

DK:May be add comment:
Characteristic only meant for
applied varieties where
vernalisation is normally
necessary for the plants to exceed
stage 90

The descriptions of the growth stages of the Zadoks decimal code for cereals

Zadoks Decimal code	Description
00	Dry seed
01	Start of imbibition
03	Imbibition complete
05	Radicle emerged from seed
07	Coleoptile emerged from seed
09	Leaf just at coleoptile tip
10	First leaf through coleoptile
11	First leaf unfolded
12	2 leaves unfolded
13	3 leaves unfolded
14	4 leaves unfolded
15	5 leaves unfolded
16	6 leaves unfolded
17	7 leaves unfolded
18	8 leaves unfolded
19	9 or more leaves unfolded
20	Main shoot only
21	Main shoot and 1 tiller
22	Main shoot and 2 tillers
23	Main shoot and 3 tillers
24	Main shoot and 4 tillers
25	Main shoot and 5 tillers
26	Main shoot and 6 tillers
27	Main shoot and 7 tillers
28	Main shoot and 8 tillers
29	Main shoot and 9 or more tillers
30	Pseudo stem erection
31	1st node detectable
32	2nd node detectable
33	3rd node detectable
34	4th node detectable
35	5th node detectable
36	6th node detectable
37	Flag leaf just visible

39	Flag leaf ligule/collar just visible
40	-
41	Flag leaf sheath extending
45	Boots just swollen
47	Flag leaf sheath opening
49	First awns visible
50	First spikelet of inflorescence visible
53	1/4 of inflorescence emerged
55	1/2 of inflorescence emerged
57	3/4 of inflorescence emerged
59	Emergence of inflorescence completed
60	Beginning on anthesis
65	Anthesis half-way
69	Anthesis completed
70	-
71	Kernel watery ripe
73	Early milk
75	Medium milk
77	Late milk
80	-
83	Early dough
85	Soft dough
87	Hard dough
90	-
91	Kernel hard (difficult to divide with thumbnail)
92	Kernel hard (no longer dented with thumbnail)
93	Kernel loosening in daytime
94	Overripe, straw dead and collapsing
95	Seed dormant
96	Viable seed giving 50% germination
97	Seed not dormant
98	Secondary dormancy induced
99	Secondary dormancy lost

9. Literature

BEZAR, H.J., HADFIELD, P.D., 1982: "Identification of New Zealand Wheat Cultivars" Crop Research Division, D.S.I.R., Christchurch, NZ.

BRIGGLE, L.W., REITZ, L.P., 1963: "Classification of Triticum Species and of Wheat Varieties Grown in the United States" United States Department of Agriculture, Technical Bulletin No. 1278, US.

BUSTARRET, J., 1944: "Variétés et variations" Annales agronomiques, 14ème année, 336, 365, FR.

DE BACKER, A., 1983: "L'homogénéité des variétés de Blé" Mémoire de fin d'études, 122e promotion Beauvais, FR.

DHORNE, D., 1985: "Les cultivars de blé (Triticum spp) et leur identification" Mémoire pour l'obtention du titre d'Ingénieur D.P.E., Ecole Nationale Supérieure Agronomique de Toulouse, FR.

FEINS, G.K. et al, 1975: "Australian Wheat Varieties" CSIR Wheat Research Unit, North Rye, New South Wales, AU.

FRIEDBERG, L. 1958

HERVEY-MURRAY, C.G., 1980: "The Identification of Cereal Varieties", Cambridge University Press, GB.

JONARD, P., 1951: "Les blés tendres (Triticum vulgare vill) cultivés en France" Paris, Institut National de la Recherche Agronomique, FR.

MILATZ, R., 1970: "Kriterien der Getreidearten einschliesslich Mais und ihre Bewertung zur Sortenidentifizierung" Bonn, Verband Deutscher Pflanzenzüchter, DE.

PERCIVAL, J., 1921: "The Wheat Plant" monograph, London, Duckworth and Co., GB.

J.C. Zadoks, T.T. Chang and C.F. Konzak, EUCARPIA Bulletin No 7, 1974, pages 42 - 52.

Comment [G61]:
DE proposes to add this reference

10. Technical Questionnaire

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
		Application date: (not to be filled in by the applicant)
TECHNICAL QUESTIONNAIRE to be completed in connection with an application for plant breeders' rights		
In the case of hybrid varieties which are the subject of an application for plant breeders' rights, and where the parent lines are to be submitted as a part of the examination of the hybrid variety, this Technical Questionnaire should be completed for each of the parent lines, in addition to being completed for the hybrid variety.		
1. Subject of the Technical Questionnaire		
1.1 Botanical name	<input type="text" value="Triticum aestivum L. emend. Fiori et Paol."/>	
1.2 Common name	<input type="text" value="Wheat"/>	
2. Applicant		
Name	<input type="text"/>	
Address	<input type="text"/>	
Telephone No.	<input type="text"/>	
Fax No.	<input type="text"/>	
E-mail address	<input type="text"/>	
Breeder (if different from applicant)	<input type="text"/>	
3. Proposed denomination and breeder's reference		
Proposed denomination (if available)	<input type="text"/>	
Breeder's reference	<input type="text"/>	

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

#4. Information on the breeding scheme and propagation of the variety

4.1 Breeding scheme

Variety resulting from:

4.1.1 Crossing

(a) controlled cross []
(please state parent varieties)

(.....) x (.....)
female parent male parent

(b) partially known cross []
(please state known parent variety(ies))

(.....) x (.....)
female parent male parent

(c) unknown cross []

4.1.2 Mutation []
(please state parent variety)

.....

4.1.3 Discovery and development []
(please state where and when discovered and how developed)

.....

4.1.4 Other []
(please provide details)

.....

Authorities may allow certain of this information to be provided in a confidential section of the Technical Questionnaire.

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

4.2 Method of propagating the variety

4.2.1 Seed-propagated varieties

- (a) Self-pollination []
- (b) Hybrid []
- (c) Other []
(please provide details)

4.2.2 Other []
(please provide details)


In the case of hybrid varieties the production scheme for the hybrid should be provided on a separate sheet. This should provide details of all the parent lines required for propagating the hybrid e.g.

Single Hybrid

(.....) x (.....)
female parent male parent

Three-Way Hybrid

(.....) x (.....)
female line male line



(.....) x (.....)
single hybrid used as female parent male parent

and should identify in particular:

- (a) any male sterile lines
- (b) maintenance system of male sterile lines.

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

5. Characteristics of the variety to be indicated (the number in brackets refers to the corresponding characteristic in Test Guidelines; please mark the note which best corresponds).

Characteristics	Example Varieties	Note
5.1 Seasonal type : (25)		
Winter type	Aubusson	1 []
Alternative type	Cezanne	2 []
Spring type	Josselin	3 []
5.2 Time of emergence : (5)		
Very early	Accor	1 []
Early	Caphor	3 []
Medium	Richepain	5 []
Late	Boncap	7 []
Very late		9 []
5.3 Plant : length (10)		
Very short	Cordiale	1 []
Short	Renan	3 []
Medium	Intérêt	5 []
Long	Bagatelle 007	7 []
Very long	Vitus	9 []
5.4 Straw : pith in cross section (11)		
Absent or very thin	Pakito	1 []
Medium	Saturnus	2 []
Very thick	Aerobic	3 []

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:	
	Characteristics	Example Varieties	Note
5.5 (15)	Awns or scurs : presence		
	Both absent	Genoveva	1 []
	Scurs present	Aubusson	2 []
	Awns present	Arezzo	3 []
5.6 (17)	Ear : color		
	White	Arezzo	1 []
	Colored	Segor	2 []

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

6. Similar varieties and differences from these varieties

Please use the following table and box for comments to provide information on how your candidate variety differs from the variety (or varieties) which, to the best of your knowledge, is (or are) most similar. This information may help the examination authority to conduct its examination of distinctness in a more efficient way.

Denomination(s) of variety(ies) similar to your candidate variety	Characteristic(s) in which your candidate variety differs from the similar variety(ies)	Describe the expression of the characteristic(s) for the similar variety	Describe the expression of the characteristic(s) for your candidate variety
<i>Caphorn</i>	<i>Coleoptile:anthocyanin coloration</i>	9	6

Comments:

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

#7. Additional information which may help in the examination of the variety

7.1 In addition to the information provided in sections 5 and 6, are there any additional characteristics which may help to distinguish the variety?

Yes [] No []

(If yes, please provide details)

7.2 Are there any special conditions for growing the variety or conducting the examination?

Yes [] No []

(If yes, please provide details)

7.3 Other information

A representative color image of the variety should accompany the Technical Questionnaire.

8. Authorization for release

(a) Does the variety require prior authorization for release under legislation concerning the protection of the environment, human and animal health?

Yes [] No []

(b) Has such authorization been obtained?

Yes [] No []

If the answer to (b) is yes, please attach a copy of the authorization.

Authorities may allow certain of this information to be provided in a confidential section of the Technical Questionnaire.

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

9. Information on plant material to be examined or submitted for examination.

9.1 The expression of a characteristic or several characteristics of a variety may be affected by factors, such as pests and disease, chemical treatment (e.g. growth retardants or pesticides), effects of tissue culture, different rootstocks, scions taken from different growth phases of a tree, etc.

9.2 The plant material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If the plant material has undergone such treatment, full details of the treatment must be given. In this respect, please indicate below, to the best of your knowledge, if the plant material to be examined has been subjected to:

- | | | |
|---|---------|--------|
| (a) Microorganisms (e.g. virus, bacteria, phytoplasma) | Yes [] | No [] |
| (b) Chemical treatment (e.g. growth retardant, pesticide) | Yes [] | No [] |
| (c) Tissue culture | Yes [] | No [] |
| (d) Other factors | Yes [] | No [] |

Please provide details for where you have indicated "yes".

.....

II GENERAL COMMENTS RECEIVED BY THE SUBGROUP

(These comments don't include those from Austria, Italy and ESA which received the very first draft later because not listed among the interested bodies. If any, they will be considered during the subgroup meeting in May)

Proposals for deletion made by France

3. Flag leaf: anthocyanin coloration of auricles	49-51 VG	absent or very weak	nulle ou très faible	fehlend oder sehr gering	Soissons; Prinqual	1
Dernière feuille: pigmentation anthocyanique des oreillettes		weak	faible	gering	Niklas; Troll	3
		medium	moyenne	mittel	Cargidoc; -	5
Oberstes Blatt: Anthocyanfärbung der Auricula		strong	forte	stark	Cargo; Sunnan	7
		very strong	très forte	sehr stark	Recital; Dollar	9
23. Lowest lemma: beak (+) shape (as for 18)	80-92 VS	straight	droit	gerade	Soissons; Prinqual	1
Glumelle inférieure: forme du bec (comme pour 18)		slightly curved	légèrement coudé	leicht gebogen	Slejpner; Briscard	3
		moderately curved	demi-coudé	mittel gebogen	Sideral; Wim	5
Untere Deckspelze: Zahnform (wie unter 18)		strongly curved	fortement coudé	stark gebogen	Parade; Axona	7
		geniculate	genouillé	geknickt	Tara; -	9

Comment [G62]:
DK agrees

Comment [G63]:
RO : ;We would like to keep the characteristic ;flag leaf : anthocianin coloration of auricles

Comment [G64]:
CA: In the 'Proposals for deletion' section of the document: CA would like to keep the characteristic: flag leaf: anthocyanin colouration of auricles. We agree with your proposal to delete lowest lemma: beak shape.

Comment [G65]:
UK wishes to retain Lowest lemma: beak shape as it is a useful characteristic.

Comment [G66]: CZ agrees

AUS:

I would like to thank experts from France for the 1st draft of revision of wheat guideline. The addition of photographs in explaining the states is helpful and definitely is a major improvement over the existing one.

However, I have a question regarding the states of expression in Ear: shape in profile (Characteristic 12).

Please see the ear shape of a variety name 'Datatine' from the attached photo.



Can we call this Ear shape: Fusiform (state 2) or do we need a separate state to describe this shape?

FR thinks that the shape is fusiform but with a very dense and short ear

BR:

- BR agree with the deletion of the characteristics “flag leaf: anthocianin coloration of auricles” and “Lemma: beak shape”, because, according to Brazilian experts, these are problematic ones.

PL:

- to harmonize spelling for ‘color’ in the document – see table of characteristic and Ad. 17 and Ad. 24

- all state names should start with lowercase letter.

- example varieties for spring and winter type varieties should be provided.

UK

§ 3.4.5: UK uses one plot of 3 rows with 1000 plants and other countries may use different designs. UK suggests allowing individual testing authorities to decide on the layout they wish to use.

§ 4.2.3 UK has particular difficulties with aneuploids. What are the experiences of other countries, and should the population standard be raised to take this into consideration?

§ 4.2.5 UK proposes to delete this paragraph.

. UK proposes the following for inclusion: Lower Glume: surface roughness, Lower Glume: surface hairiness and electrophoresis.

CL:

Chili proposes this new characteristic

26.	92	Grain: ??	Graine: ??	Korn: ??	Grano: forma
(+)					
PQ	VG				Redondo
	A				Redondo eíptico
					Elíptico

Amendments in Spanish have not been introduced in the draft

Comment [G67]: These 2 points will be checked at UPOV Office and TC EDC levels

Comment [G68]:
CZ: We propose to indicate type (winter, spring) at example varieties

Comment [G69]: In the draft 1 all example varieties are winter type varieties except the varieties Scirocco (char 18 note 9) and Josselin (char 25 note 3)

Comment [G70]:
DE There are several steps and ways to test D according to TGP/9 and TGP/8. I think it is not appropriate to mention one very specific possible additional tool. FR This paragraph initially proposed by France has been deleted

Comment [G71]:
DE: The specific procedure is appropriate for the CPVO-TP but not for the UPOV-TG.
FR: This paragraph initially proposed by France has been deleted

DE

I understand that there are different views in relation to the illustrations in chapter 8. But we are of the opinion that many of the added images and some of the modified illustrations do not provide additional information and guidance for the experts in the field. In particular, the states in Ad 1, 2, 3, 4 12 and 13 are not all to the point. For Ad 14, 16 and 24 we wonder if images are necessary at all. The images in Ad 15 should be improved (better developed ear for 2, better image quality focus and color). We welcome the new illustrations for 21 and 23. For 18, 19, 20, 22 we prefer the previous version. In particular the beak in 19, 20, 22 is too blatant.

We would be happy for a review of chapter 8 for the next draft.

FR

This is a reference which explains that the soaking of seeds is not necessary for the observation of the characteristic coloration with phenol (char.1) :

Essai de classification des blés d'après leur réaction à l'acide phénique

L. Friedberg

impr. Marc Texier, 1933

40 pages

Document prepared by France , Beaucouzé April, 2nd, 2012

[End of Document]