



TG/COCOS(proj.2)

ORIGINAL: English

DATE: 2013-03-19

INTERNATIONAL UNION FOR THE PROTECTION OF NEW VARIETIES OF PLANTS

Geneva


COCONUT

UPOV Code: COCOS_NUC

Cocos nucifera L.**GUIDELINES****FOR THE CONDUCT OF TESTS****FOR DISTINCTNESS, UNIFORMITY AND STABILITY***prepared by experts from Brazil**to be considered by the**Technical Working Party for Fruit Crops**at its forty-fourth session, to be held in Napier, New Zealand, from April 29 to May 3, 2013*Alternative Names:^{*}

<i>Botanical name</i>	<i>English</i>	<i>French</i>	<i>German</i>	<i>Spanish</i>
<i>Cocos nucifera</i> L.	Coconut	Cocotier	Kokosnuß	Cocotero

The purpose of these guidelines ("Test Guidelines") is to elaborate the principles contained in the General Introduction (document TG/1/3), and its associated TGP documents, into detailed practical guidance for the harmonized examination of distinctness, uniformity and stability (DUS) and, in particular, to identify appropriate characteristics for the examination of DUS and production of harmonized variety descriptions.

ASSOCIATED DOCUMENTS

These Test Guidelines should be read in conjunction with the General Introduction and its associated TGP documents.

^{*} These names were correct at the time of the introduction of these Test Guidelines but may be revised or updated. [Readers are advised to consult the UPOV Code, which can be found on the UPOV Website (www.upov.int), for the latest information.]

<u>TABLE OF CONTENTS</u>	<u>PAGE</u>
1. SUBJECT OF THESE TEST GUIDELINES	3
2. MATERIAL REQUIRED	3
3. METHOD OF EXAMINATION	3
3.1 NUMBER OF GROWING CYCLES	3
3.2 TESTING PLACE	3
3.3 CONDITIONS FOR CONDUCTING THE EXAMINATION	3
3.4 TEST DESIGN	3
3.5 ADDITIONAL TESTS.....	3
4. ASSESSMENT OF DISTINCTNESS, UNIFORMITY AND STABILITY	4
4.1 DISTINCTNESS	4
4.2 UNIFORMITY	5
4.3 STABILITY.....	5
5. GROUPING OF VARIETIES AND ORGANIZATION OF THE GROWING TRIAL.....	5
6. INTRODUCTION TO THE TABLE OF CHARACTERISTICS.....	5
6.1 CATEGORIES OF CHARACTERISTICS	5
6.2 STATES OF EXPRESSION AND CORRESPONDING NOTES.....	6
6.3 TYPES OF EXPRESSION	6
6.4 EXAMPLE VARIETIES	6
6.5 LEGEND	7
7. TABLE OF CHARACTERISTICS/TABLEAU DES CARACTÈRES/MERKMALSTABELLE/TABLA DE CARACTERES	8
8. EXPLANATIONS ON THE TABLE OF CHARACTERISTICS.....	14
8.1 EXPLANATIONS COVERING SEVERAL CHARACTERISTICS.....	14
8.2 EXPLANATIONS FOR INDIVIDUAL CHARACTERISTICS	14
9. LITERATURE	24
10. TECHNICAL QUESTIONNAIRE.....	25

1. Subject of these Test Guidelines

These Test Guidelines apply to all varieties of *Cocos nucifera* L., including tall type, dwarf type and hybrid tall x dwarf, under controlled pollination.

2. Material Required

2.1 The competent authorities decide on the quantity and quality of the plant material required for testing the variety and when and where it is to be delivered. Applicants submitting material from a State other than that in which the testing takes place must ensure that all customs formalities and phytosanitary requirements are complied with.

2.2 The material is to be supplied in the form of seeds (fruits).

2.3 The minimum quantity of plant material, to be supplied by the applicant, should be:

20 seeds.

2.4 The plant material supplied should be visibly healthy, not lacking in vigor, nor affected by any important pest or disease.

2.5 The plant material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If it has been treated, full details of the treatment must be given.

3. Method of Examination

3.1 *Number of Growing Cycles*

3.1.1 The minimum duration of tests should normally be two independent growing cycles. In particular, it is essential that the palms produce a satisfactory crop of fruit in each of the two growing cycles.

3.1.2 The growing cycle is considered to be the period ranging from the beginning of active vegetative growth or flowering, continuing through active vegetative growth or flowering and fruit development and concluding with the harvesting of fruit.

3.2 *Testing Place*

Tests are normally conducted at one place. In the case of tests conducted at more than one place, guidance is provided in TGP/9 "Examining Distinctness".

3.3 *Conditions for Conducting the Examination*

The tests should be carried out under conditions ensuring satisfactory growth for the expression of the relevant characteristics of the variety and for the conduct of the examination. In particular, it is essential that the trees produce a satisfactory crop of fruit in each of the two growing cycles.

3.4 *Test Design*

3.4.1 Each test should be designed to result in a total of at least 12 palms.

3.4.2 The design of the tests should be such that plants or parts of plants may be removed for measurement or counting without prejudice to the observations which must be made up to the end of the growing cycle.

3.5 *Additional Tests*

Additional tests, for examining relevant characteristics, may be established.

4. Assessment of Distinctness, Uniformity and Stability

4.1 *Distinctness*

4.1.1 General Recommendations

It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding distinctness. However, the following points are provided for elaboration or emphasis in these Test Guidelines.

Further guidance is provided in documents TGP/9 “Examining Distinctness” and TGP/8 “Trial Design and Techniques Used in the Examination of Distinctness, Uniformity and Stability”.

4.1.2 Consistent Differences

The differences observed between varieties may be so clear that more than one growing cycle is not necessary. In addition, in some circumstances, the influence of the environment is not such that more than a single growing cycle is required to provide assurance that the differences observed between varieties are sufficiently consistent. One means of ensuring that a difference in a characteristic, observed in a growing trial, is sufficiently consistent is to examine the characteristic in at least two independent growing cycles.

4.1.3 Clear Differences

Determining whether a difference between two varieties is clear depends on many factors, and should consider, in particular, the type of expression of the characteristic being examined, i.e. whether it is expressed in a qualitative, quantitative, or pseudo-qualitative manner. Therefore, it is important that users of these Test Guidelines are familiar with the recommendations contained in the General Introduction prior to making decisions regarding distinctness.

4.1.4 Number of Plants / Parts of Plants to be Examined

Unless otherwise indicated, for the purposes of distinctness, all observations on single plants should be made on 12 plants or parts taken from each of 12 plants and any other observations made on all plants in the test, disregarding any off-type plants. In the case of observations of parts taken from single plants, the number of parts to be taken from each of the plants should be 2.

4.1.5 Method of Observation

The recommended method of observing the characteristic for the purposes of distinctness is indicated by the following key in the second column of the Table of Characteristics (see document TGP/9 “Examining Distinctness”, Section 4 “Observation of characteristics”):

MG: single measurement of a group of plants or parts of plants

MS: measurement of a number of individual plants or parts of plants

VG: visual assessment by a single observation of a group of plants or parts of plants

VS: visual assessment by observation of individual plants or parts of plants

Type of observation: visual (V) or measurement (M)

“Visual” observation (V) is an observation made on the basis of the expert’s judgment. For the purposes of this document, “visual” observation refers to the sensory observations of the experts and, therefore, also includes smell, taste and touch. Visual observation includes observations where the expert uses reference points (e.g. diagrams, example varieties, side-by-side comparison) or non-linear charts (e.g. color charts). Measurement (M) is an objective observation against a calibrated, linear scale e.g. using a ruler, weighing scales, colorimeter, dates, counts, etc.

Type of record: for a group of plants (G) or for single, individual plants (S)

For the purposes of distinctness, observations may be recorded as a single record for a group of plants or parts of plants (G), or may be recorded as records for a number of single, individual plants or parts of plants (S). In most cases, “G” provides a single record per variety and it is not

possible or necessary to apply statistical methods in a plant-by-plant analysis for the assessment of distinctness.

In cases where more than one method of observing the characteristic is indicated in the Table of Characteristics (e.g. VG/MG), guidance on selecting an appropriate method is provided in document TGP/9, Section 4.2.

4.2 *Uniformity*

4.2.1 It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding uniformity. However, the following points are provided for elaboration or emphasis in these Test Guidelines:

4.2.2. For the assessment of uniformity of controlled pollinated varieties, a population standard of 1% and an acceptance probability of at least 95% should be applied. In the case of a sample size of 12 plants, 1 off-type is allowed.

4.3 *Stability*

4.3.1 In practice, it is not usual to perform tests of stability that produce results as certain as those of the testing of distinctness and uniformity. However, experience has demonstrated that, for many types of variety, when a variety has been shown to be uniform, it can also be considered to be stable.

4.3.2 Where appropriate, or in cases of doubt, stability may be further examined by testing a new seed or plant stock to ensure that it exhibits the same characteristics as those shown by the initial material supplied.”

5. Grouping of Varieties and Organization of the Growing Trial

5.1 The selection of varieties of common knowledge to be grown in the trial with the candidate varieties and the way in which these varieties are divided into groups to facilitate the assessment of distinctness are aided by the use of grouping characteristics.

5.2 Grouping characteristics are those in which the documented states of expression, even where produced at different locations, can be used, either individually or in combination with other such characteristics: (a) to select varieties of common knowledge that can be excluded from the growing trial used for examination of distinctness; and (b) to organize the growing trial so that similar varieties are grouped together.

5.3 The following have been agreed as useful grouping characteristics:

- (a) Stem: diameter (characteristic 9)
- (b) Fruit: color (characteristic 27)
- (c) Fruit: weight (characteristic 28)
- (d) Fruit: shape (characteristic 29)
- (e) Nut: shape (characteristic 31)

5.4 Guidance for the use of grouping characteristics, in the process of examining distinctness, is provided through the General Introduction and document TGP/9 “Examining Distinctness”.

6. Introduction to the Table of Characteristics

6.1 *Categories of Characteristics*

6.1.1 Standard Test Guidelines Characteristics

Standard Test Guidelines characteristics are those which are approved by UPOV for examination of DUS and from which members of the Union can select those suitable for their particular circumstances.

6.1.2 Asterisked Characteristics

Asterisked characteristics (denoted by *) are those included in the Test Guidelines which are important for the international harmonization of variety descriptions and should always be examined for DUS and included in the variety description by all members of the Union, except when the state of expression of a preceding characteristic or regional environmental conditions render this inappropriate.

6.2 States of Expression and Corresponding Notes

6.2.1 States of expression are given for each characteristic to define the characteristic and to harmonize descriptions. Each state of expression is allocated a corresponding numerical note for ease of recording of data and for the production and exchange of the description.

6.2.2 In the case of qualitative and pseudo-qualitative characteristics (see Chapter 6.3), all relevant states of expression are presented in the characteristic. However, in the case of quantitative characteristics with 5 or more states, an abbreviated scale may be used to minimize the size of the Table of Characteristics. For example, in the case of a quantitative characteristic with 9 states, the presentation of states of expression in the Test Guidelines may be abbreviated as follows:

State	Note
small	3
medium	5
large	7

However, it should be noted that all of the following 9 states of expression exist to describe varieties and should be used as appropriate:

State	Note
very small	1
very small to small	2
small	3
small to medium	4
medium	5
medium to large	6
large	7
large to very large	8
very large	9

6.2.3 Further explanation of the presentation of states of expression and notes is provided in document TGP/7 "Development of Test Guidelines".

6.3 Types of Expression

An explanation of the types of expression of characteristics (qualitative, quantitative and pseudo-qualitative) is provided in the General Introduction.

6.4 Example Varieties

Where appropriate, example varieties are provided to clarify the states of expression of each characteristic.

6.5 *Legend*

- (*) Asterisk characteristic – see Chapter 6.1.2
- QL Qualitative characteristic – see Chapter 6.3
- QN Quantitative characteristic – see Chapter 6.3
- PQ Pseudo-qualitative characteristic – see Chapter 6.3
- MG, MS, VG, VS – see Chapter 4.1.5
- (a)-(d) See Explanations on the Table of Characteristics in Chapter 8.1
- (+) See Explanations on the Table of Characteristics in Chapter 8.2.

7. Table of Characteristics/Tableau des caractères/Merkmalstabelle/Tabla de caracteres

	English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
1. VG (*) (+)	Young plant: color of shoot:				Indonesia to check their examples	
PQ	yellow				Genjah Kuning Nias	1
	green				Genjah Salak	2
	red				Dalam Mapanget	3
	brown				Genjah Raja Brown	4
2. VG/MS (*) (+)	Young plant: number of leaves					
QN	few				Genjah Kuning Nias	3
	medium				Genjah Raja Brown	5
	many				Dalam Mapanget	7
3. VG (*) (+)	Young plant: time of leaflets splitting					
QN	early				GR ????	3
	medium				Genjah Kuning Bali	5
	late					7
4. MG (*) (+)	Time of appearance of first inflorescence					
QN	early				Genjah Tebing Tinggi	3
	medium				Dalam Mapanget	5
	late				DJP ????	7
5. VG (*) (+)	Leaves attitude of lower leaves					
QN (a)	upwards					1
	outwards					2
	downwards					3
6. VG (*) (+)	Stem: bole					
QL (a)	absent				Genjah Kuning Nias	1
	present					9
7. VG/MS (*) (+)	Stem: diameter of the bole					
QN (a)	small				Kelapa Hibrida Indonesia-1	1
	medium					3
	large				Dalam Mapanget	5

	English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
8. VG/ MS (*) (+)	Stem: height					
QN (a)	short					3
	medium					5
	long					7
9. VG/ MS (*) (+)	Stem: diameter					
QN (a)	small					3
	medium					5
	large					7
10. VG/ MS (+)	Petiole: length					
QN (a)	short				Genjah Kuning Nias	3
	medium				Kelapa Hibrida Indonesia-1	5
	long				Dalam Tenga	7
11. VG/ MS (+)	Petiole: thickness					
QN (a)	thin				Genjah Kuning Nias	3
	medium				Kelapa Hibrida Indonesia-2	5
	thick				Dalam Bali	7
12. VG/ MS (+)	Petiole: width					
QN (a)	narrow				Dalam Mapanget	3
	medium				Genjah Raja Brown	5
	broad				Dalam Kima Atas	7
13. VG	Petiole: color					
PQ (a)	yellow				Genjah Kuning Nias	1
	yellowish green					2
	yellowish red					3
	green				Genjah Salak	4
	greenish yellow					5
	greenish red					6
	red				Dalam Mapanget	7
	reddish yellow					8
	reddish green					9
	brown				Genjah Raja Brown	10

	English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
14. VG/ MS (+)	Leaf: length of rachis					
QN (a)	short				Genjah Kuning Nias	3
	medium				Kelapa Hibrida Indonesia-3	5
	long				Dalam Palu	7
15. VG/ MS (+)	Leaf: number of leaflets					
QN (a)	few				DTT ????	3
	medium				DJP ????	5
	many				Dalam Takome	7
16. VG/ MS (+)	Leaf: leaflet length					
QN (a)	short				Genjah Kuning Nias	3
	medium				DTT ????	5
	long				Dalam Kima Atas	7
17. VG/ MS (+)	Leaf: leaflet width					
QN (a)	narrow				Genjah Kuning Bali	3
	medium				DMA ????	5
	broad				DAG ????????	7
18. VG (+)	Leaflet: intensity of green color:					
QN (a)	light					1
	medium					2
	dark					3
19. VG/ MS (+)	Inflorescence: peduncle length					
QN (b)	short				Genjah Raja Brown	3
	medium				Kelapa Hibrida Indonesia-4	5
	long				Dalam Mapanget	7
20. VG/ MS (+)	Inflorescence: peduncle diameter					
QN (b)	small				Genjah Raja Brown	3
	medium				Kelapa Hibrida Indonesia-4	5
	large				Dalam Mapanget	7

	English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
21.	VG/ MS	Inflorescence: central axis length				
(+)						
QN	(b)	short			Genjah Raja Brown	3
		medium			Kelapa Hibrida Indonesia- 4	5
		long			Dalam Mapanget	7
22.	VG/ MS	Inflorescence: number of spikelets				
(+)						
QN	(b)	few			Dalam Tenga	3
		medium			Dalam Banyuwangi	5
		many			Dalam Kima Atas	7
23.	VG/ MS	Inflorescence: number of spikelets with female flowers				
(*) (+)						
QN	(b)	few				3
		medium				5
		many				7
24.	VG/ MS	Inflorescence: length of first spikelet bearing female flower				
(+)						
QN	(b)	short			Genjah Raja Brown	3
		medium			Kelapa Hibrida Indonesia- 4	5
		long			Dalam Mapanget	7
25.	VG/ MS	Bunch: number of fruits				
QN	(c)	few			Dalam Sawarna	3
		medium			Dalam Mapanget	5
		many			Dalam Takome	7
26.	VG/ MS	Bunch: peduncle length	BR: to delete, related char. 19			
(+)						
QN	(c)	short				3
		medium				5
		long				7

	English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
27.	VG					
(+)						
(*)						
PQ	(c)	yellow			Genjah Kuning Bali	1
		yellowish green				2
		yellowish red				3
		green			Genjah Hijau Nias	4
		greenish yellow				5
		greenish red				6
		red			GMW????	7
		reddish yellow				8
		reddish green				9
		brown			Genjah Raja Brown	10
28.	MS	Fruit: weight				
(*)						
(+)						
QN	(d)	low			GOS ?????	3
		medium			Genjah Tebing Tinggi	5
		high			Dalam Bali	7
29.	VG/ MS	Fruit: shape				
(*)						
(+)						
PQ	(d)	circular				1
		ovate				2
		elliptic				3
		obovate				4
		pyriform				5
30.	VG/ MS	Fruit:ratio weight fruit/husk				
(+)						
QN	(d)	low				1
		medium				3
		high				5
31.	VG/ MS	Nut: shape				
(*)						
(+)						
PQ	(d)	ovate				1
		elliptic				2
		circular				3
		oblate				4

	English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
32. VG/ MS	Nut: weight					
QN (d)	low				GOS ?????	3
	medium				Genjah Raja Brown	5
	high				Dalam Palu	7
33. VG/ MS (+)	Shell: thickness					
QN (d)	thin				Genjah Kuning Nias	3
	medium				Kelapa Hibrida Indonesia-1	5
	thick				Dalam Tenga	7
34. VG/ MS (*) (+)	Meat: weight					
QN (d)	low				GOS ?????	3
	medium				Dalam Tenga	5
	high				Dalam Bali	7
35. VG/ MS (*)	Meat: thickness					
QN (d)	thin				GKJ????	3
	medium				Dalam Sawarna	5
	thick				Dalam Mapanget	7
36.	Consistency of meat					
QL (d)	solid	compact???				1
	lose	jelly????				2

8. Explanations on the Table of Characteristics

8.1 *Explanations covering several characteristics*

Characteristics containing the following key in the second column of the Table of Characteristics should be examined as indicated below:

- (a) Palm, stem, petiole and petiole and leaf: Observations should be made when the eleventh leaf scars appears (see photo Ad. 5 to 9: leaf scars). Observations on leaf and petiole should be made at 14^a leaf on maturity leaf.
- (b) Inflorescence: Observations on inflorescence should be taken after the appearance of the fifth inflorescence, when female flowers are receptive.
- (c) Bunch, peduncle and fruit color: Observations on the bunch, peduncle and fruit color should be made at maturity for consumption as water (at 6-7 months age fruit), after the appearance of the sixth bunch (**we need to take off the fifth inflorescence to evaluate**).
- (d) Fruit, nut, shell and meat. Observations on the fruit, nut, shell and meat should be made at maturity for consumption as copra (at 11-12 months age fruit), after the appearance of the sixth bunch.

(a) Leaf scars (Ad. 5 to 18)



8.2 *Explanations for individual characteristics*

Ad. 1: Young plant: color of shoot

Should be observed as soon as shoot emergence.

Ad. 2: Young plant: number of leaves

Should be observed at age 6 months after the germination of the seed.

Ad. 3: Young plant: time of leaflets splitting


The leaflets splitting begins around age 6 months of the young plant.




Ad. 4: Time of appearance of first inflorescence

The time of appearance of first inflorescence should be observed when 50% of the plants have emitted the first inflorescence.

Ad. 5: Leaves attitude of lower leaves



Ad. 6: Stem: bole



Ad. 7 Stem: diameter of the bole

The diameter of the bole should be measured at 0.2 m at the soil.

Ad. 8: Stem: height


Ad. 9: Stem: diameter

The stem diameter should be measured from ground to 1.5 meters.



Ad. 10: Petiole: length


The petiole length should be measured from base to the most proximal leaflet of the rachis.



Ad. 11: Petiole: thickness

Ad. 12: Petiole: width

The petiole thickness and the petiole width should be measured at insertion of the first leaflet.



Ad. 14: Leaf: length of rachis


The length of rachis should be measured from to the most proximal leaflet to the tip of the rachis.

Ad. 16: Leaf: leaflet length

Ad. 17: Leaf: leaflet width

The leaflet length and the leaflet width should be measured at halfway of the rachis


- Ad. 19: Inflorescence: peduncle length
- Ad. 20: Inflorescence: peduncle diameter
- Ad. 21: Inflorescence: central axis length



- Ad. 22: Inflorescence: number of spikelets




Ad. 23: Inflorescence: number of spikelets with female flowers








Ad. 24: Inflorescence: length of first spikelet bearing female flower

Should be evaluated after the appearance of the fifth inflorescence, when the first female flower appears.

Ad. 27: Fruit: color



Ad. 29: Fruit: shape

		< broadest part >		
		(below middle)	at middle	(above middle)
< lateral outline >	rounded	 <p>2 ovate</p>	 <p>3 elliptic</p>	 <p>4 obovate</p>
	rounded with neck		 <p>1 circular</p>	 <p>5 pyriform</p>

Ad. 30: Fruit: the ratio weight fruit/husk

Ad. 31: Nut: shape





Ad. 33: Shell: thickness

Ad. 34: Meat: weight



1.	exocarp	husk = exocarp + mesocarp
2.	mesocarp	
3.	endocarp (shell)	nut = endocarp + endosperm
4.	endosperm (meat)	
5.	embryo	

Ad. 31: Nut: shape

		← Broadest part →		
		below middle	middle	above middle
ratio width/length	↑ elongated	 1 ovate	 2 elliptic	
	medium		 3 circular	
	↓ compressed		 4 oblate	

Ad. 36: Consistency of meat



A

B

1

solid ???
compact???

2

lose ?????
jelly????

9. Literature

IPGRI, 1995: Descriptors for Coconut (*Cocos nucifera* L.). International Plant Genetic Resources Institute. Rome, IT, 61 pp.

The Minimum List of Descriptors for coconut – from CIRAD with same other countries contributions (China, Brasil, France, Mexico, Indonesia, Malaysia, Philippines, Viet Nam), ano....

Ribeiro, F.E., de Siqueira, E.R., Aragão, W.M., Tupinambá, E.A., 1999: O coqueiro anão no Brasil. Aracaju: Embrapa Tabuleiros Costeiros, 23p. (Embrapa-Tabuleiros Costeiros. Documentos, 8).

Ribeiro, F.E., de Siqueira, E.R., Aragão, W.M., Tupinambá, E.A., 2000: Ecótipos de coqueiro gigante no Brasil. Aracaju: Embrapa Tabuleiros Costeiros, 25p. (Embrapa-Tabuleiros Costeiros. Documentos, 17).

Ribeiro, F.E., de J. Ribeiro, M.M., 2011: Caracterização de populações de coqueiro gigante no Nordeste do Brasil. Aracaju: Embrapa Tabuleiros Costeiros, 16p. (Embrapa-Tabuleiros Costeiros. Boletim de Pesquisa e Desenvolvimento, 59).

Aragão, W.M., Ribeiro, F.E., Tupinambá, E.A., de Siqueira, E.R., 2003: Variedades e híbridos. In: Fontes, H.R., Ribeiro, F.E., Fernandes, M.F., (Ed.). Coco produção: Aspectos técnicos. Aracaju: Embrapa Tabuleiros Costeiros, p. 21.

de Siqueira, E.R., Ribeiro, F.E., Aragão, W.M., Tupinambá, E.A., 1998: Melhoramento genético do coqueiro. In: Ferreira, J.M.S.; Warwick, D.R.N.; Siqueira, L.A. (Ed.). A cultura do coqueiro no Brasil. 2. Ed. rev. Amp. Brasília: Embrapa-SPI, 292p.

Aragão, W.M., Ribeiro, F.E., de V. Melo, M.F. 2009: Cultivares de coqueiro para produção de coco seco: coqueiro gigante vs híbridos. In: Cintra, F.L.D., Fontes, H.R., Passos, E.E.M., Ferreira, J.M.S., (Ed.). Fundamentos tecnológicos para a revitalização das áreas cultivadas com coqueiro gigante do Brasil. Aracaju: Embrapa Tabuleiros Costeiros, 232p.

Wuidart, W., Rognon, F., 1978: L'analyse de composant de la noix de cocotier: Méthode de détermination du coprah. *Oléagineux*, **33**(5):225-33.

10. Technical Questionnaire

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
		Application date: (not to be filled in by the applicant)
TECHNICAL QUESTIONNAIRE to be completed in connection with an application for plant breeders' rights		
1. Subject of the Technical Questionnaire		
1.1 Botanical name	<input type="text" value="Cocos nucifera L."/>	
1.2 Common name	<input type="text" value="Coconut"/>	
2. Applicant		
Name	<input type="text"/>	
Address	<input type="text"/>	
Telephone No.	<input type="text"/>	
Fax No.	<input type="text"/>	
E-mail address	<input type="text"/>	
Breeder (if different from applicant)	<input type="text"/>	
3. Proposed denomination and breeder's reference		
Proposed denomination (if available)	<input type="text"/>	
Breeder's reference	<input type="text"/>	

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

#4. Information on the breeding scheme and propagation of the variety

4.1 Breeding scheme

Variety resulting from:

4.1.1 Crossing

(a) controlled cross
(please state parent varieties)

(.....) x (.....)
female parent male parent

(b) partially known cross
(please state known parent variety(ies))

(.....) x (.....)
female parent male parent

(c) unknown cross

4.1.2 Mutation
(please state parent variety)

4.1.3 Discovery and development
(please state where and when discovered and how developed)

4.1.4 Other

(please provide details)

Authorities may allow certain of this information to be provided in a confidential section of the Technical Questionnaire.

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

4.2 Method of propagating the variety

4.2.1 Seed-propagated varieties

- (a) Self-pollination []
- (b) Cross-pollination []
 - (i) population []
 - (ii) synthetic variety []
- (c) Hybrid []
- (d) Other []
(please provide details)

[]

4.2.2 Vegetative propagation

- (a) cuttings []
- (b) *in vitro* propagation []
- (c) other (state method) []

[]

4.2.3 Other []
(please provide details)

[]

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

5. Characteristics of the variety to be indicated (the number in brackets refers to the corresponding characteristic in Test Guidelines; please mark the note which best corresponds).

Characteristics	Example Varieties	Note
5.1 Stem: diameter (9)		
very small		1[]
very small to small		2[]
small		3[]
small to medium		4[]
medium		5[]
medium to large		6[]
large		7[]
large to very large		8[]
very large		9[]
5.2 Fruit: color (27)		
yellow	Genjah Kuning Bali	1[]
yellowish green		2[]
yellowish red		3[]
green	Genjah Hijau Nias	4[]
greenish yellow		5[]
greenish red		6[]
red	GMW???	7[]
reddish yellow		8[]
reddish green		9[]
brown	Genjah Raja Brown	10[]

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

Characteristics	Example Varieties	Note
5.3 Fruit: weight (28)		
very low		1[]
very low to low		2[]
low	GOS???	3[]
low to medium		4[]
medium	Genjah Tebing Tinggi	5[]
medium to high		6[]
high	Dalam Bali	7[]
high to very high		8[]
very high		9[]
5.4 Fruit: shape (29)		
circular		1[]
ovate		2[]
elliptic		3[]
obovate		4[]
pyriform		5[]
5.5 Nut: shape (31)		
ovate		1[]
elliptic		2[]
circular		3[]
oblate		4[]

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

6. Similar varieties and differences from these varieties

Please use the following table and box for comments to provide information on how your candidate variety differs from the variety (or varieties) which, to the best of your knowledge, is (or are) most similar. This information may help the examination authority to conduct its examination of distinctness in a more efficient way.

Denomination(s) of variety(ies) similar to your candidate variety	Characteristic(s) in which your candidate variety differs from the similar variety(ies)	Describe the expression of the characteristic(s) for the similar variety(ies)	Describe the expression of the characteristic(s) for your candidate variety
---	---	--	--

Genjah Hijau Nias	<i>Fruit: color</i>	<i>green</i>	<i>yellow</i>
-------------------	---------------------	--------------	---------------

--	--	--	--

--	--	--	--

--	--	--	--

Comments:

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

#7. Additional information which may help in the examination of the variety

7.1 In addition to the information provided in sections 5 and 6, are there any additional characteristics which may help to distinguish the variety?

Yes [] No []

(If yes, please provide details)

7.2 Are there any special conditions for growing the variety or conducting the examination?

Yes [] No []

(If yes, please provide details)

7.3 Other information

Main use of the variety:

(a) pot plant []

(b) garden plant []

(c) other []

A representative color image of the variety should accompany the Technical Questionnaire.

8. Authorization for release

(a) Does the variety require prior authorization for release under legislation concerning the protection of the environment, human and animal health?

Yes [] No []

(b) Has such authorization been obtained?

Yes [] No []

If the answer to (b) is yes, please attach a copy of the authorization.

Authorities may allow certain of this information to be provided in a confidential section of the Technical Questionnaire.

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

9. Information on plant material to be examined or submitted for examination.

9.1 The expression of a characteristic or several characteristics of a variety may be affected by factors, such as pests and disease, chemical treatment (e.g. growth retardants or pesticides), effects of tissue culture, different rootstocks, scions taken from different growth phases of a tree, etc.

9.2 The plant material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If the plant material has undergone such treatment, full details of the treatment must be given. In this respect, please indicate below, to the best of your knowledge, if the plant material to be examined has been subjected to:

- | | | |
|---|---------|--------|
| (a) Microorganisms (e.g. virus, bacteria, phytoplasma) | Yes [] | No [] |
| (b) Chemical treatment (e.g. growth retardant, pesticide) | Yes [] | No [] |
| (c) Tissue culture | Yes [] | No [] |
| (d) Other factors | Yes [] | No [] |

Please provide details for where you have indicated "yes".

.....

10. I hereby declare that, to the best of my knowledge, the information provided in this form is correct:

Applicant's name

Signature

Date

[End of document]