

**Test Guidelines for Oilseed Rape: TG/36/7**  
**Additional Characteristic(s)**

Submitting Authority:

Contact Expert: Name:

Date:

Organization:

Tel.:

E-mail:

		English	français	deutsch	español	Example Varieties/ Exemples/ Beispielssorten/ Variedades ejemplo	[iii] Note/ Nota
<b>New 1.</b>	<b>MS</b>	<b>Cotyledon: lamina base to wide point (lbtwp)</b>					
	<b>QN</b>	short	court	kurz	corto	Hattrick	3
		medium	moyen	mittel	medio	DK Exmoji	5
		long	long	lang	largo	DK Excited	7
<b>New 2.</b>	<b>MS</b>	<b>Cotyledon: lbtwp/width ratio</b>					
	<b>QN</b>	low	bas	klein	baja	ES Mambo	3
		medium	moyen	mittel	media	DK Exmoji	5
		high	élevé	groß	alta	Simona	7

- [i] indicate type of expression (QL, PQ, QN)
- [ii] indicate method of observation (VG, VS, MG, MS)
- [iii] example varieties to be provided for at least 2 states.

**Explanation / Illustration (including extent of the use of the characteristic(s)):**

These characteristics have been included in the United Kingdom Test Protocol for many years. In the United Kingdom these are measured using image analysis, so the data is collected routinely, however the information is only used in the granting of plant breeder's rights in a small number of cases.

The characteristics are observed, as per the illustration, on cotyledons from 40 seedlings over two replicates:

