



BMT/13/25

ORIGINAL: English

DATE: November 10, 2011

**INTERNATIONAL UNION FOR THE PROTECTION OF NEW VARIETIES OF PLANTS**  
GENEVA

**WORKING GROUP ON BIOCHEMICAL AND MOLECULAR  
TECHNIQUES, AND DNA-PROFILING IN PARTICULAR**

**Thirteenth Session**  
**Brasilia, November 22 to 24, 2011**

**USE OF MOLECULAR MARKERS TO IDENTIFY SOYBEAN VARIETIES: THE  
EXPERIENCE OF A PUBLIC SOYBEAN BREEDING PROGRAM**

*Document prepared by experts from Brazil*

The aim of this work is to demonstrate some cases where molecular markers were useful to identify soybean varieties in order to help breeders and the official service of protection of varieties in Brazil. The partnership between Embrapa/Epamig/Fundação Triângulo, located at Uberaba, Minas Gerais State, released some important soybean varieties adapted for the Brazilian Savannah region. One of the most important varieties developed by these partners, which was a very important variety in the recent soybean history in Brazil and also in the world, was MG/BR 46 (Conquista). 'Conquista' was released in 1995 and seemed to be just more one soybean variety for the use of soybean farmers in the Minas Gerais State, but the history proved that it was very different of the other varieties. With high yield stability, adapted to a broad range of environmental conditions (especially unfavourable conditions) and with a very rustic root system (including resistance to both root rot nematode species more common in Brazil, *Meloidogyne javanica* and *M. incognita*), 'Conquista' was sown in more than three million hectares in Brazil in a growing season. It is still sown widely (about 100.00 hectares in the growing season of 2010/2011) and was also largely used in crosses by Brazilian soybean breeders. A famous chapter of 'Conquista' history was the dispute between Fundação Triângulo (Minas Gerais State) and Fundação MT (Mato Grosso State), both of them Embrapa partners. Fundação MT released a new variety named MT/BR 49 (Pioneira), a few months after 'Conquista' was released, and a breeder suspected that 'Pioneira' was very similar to 'Conquista'. The question was clarified only with the use of

molecular markers and the conclusion was that they presented the same genotype. Another history was that 'Conquista' was used to develop some RR (Roundup Ready) transgenic derived varieties (for example BRS Valiosa RR and BRS Favorita RR), and a problem of identification arose during the process of genetic seed production of BRS Valiosa RR and BRS Favorita RR, but in this case the SSR molecular markers used did not solve the problem. Another case of the use of molecular markers in Embrapa's soybean breeding program developed by the partners of the Minas Gerais State is to clear a doubt about the origin of the variety BRSMG 800A, the first soybean Brazilian with brown tegument, and that was suspected to be a mutation for tegument color of the variety BRSMG 790A.

FRONZA, V.<sup>1</sup>; ABDELNOOR, R.V.<sup>2</sup>; ARIAS, C.A.A.<sup>2</sup>; PACHECO, L.G.A.<sup>3</sup>; ARANTES, N.E.<sup>4</sup>; ZITO, R.K.<sup>5</sup>; SÁ, M.E.L.<sup>6</sup>

<sup>1</sup> Pesquisador Embrapa Soja, C.P. 311, CEP 38060-040, Uberaba-MG. E-mail: [vanoli@cnpsa.embrapa.br](mailto:vanoli@cnpsa.embrapa.br).

<sup>2</sup> Pesquisador Embrapa Soja, C.P. 231, CEP 86001-970, Londrina-PR.

<sup>3</sup> Engº Agrº Serviço Nacional de Proteção de Cultivares/MAPA, Esplanada dos Ministérios, Bloco D, Anexo A, Sala 251, CEP 70043-900, Brasília-DF.

<sup>4</sup> Pesquisador bolsista Fapemig/Epamig, C.P. 311, CEP 38060-040, Uberaba-MG.

<sup>5</sup> Pesquisador Embrapa Soja, C.P. 179, CEP 75375-000, Santo Antônio de Goiás-GO.

<sup>6</sup> Pesquisadora Epamig/Embrapa Recursos Genéticos e Biotecnologia, C.P. 02372, CEP 70770-917, Brasília-DF.

[End of document]