



TC/49/40

ORIGINAL: English

DATE: February 7, 2013

**INTERNATIONAL UNION FOR THE PROTECTION OF NEW VARIETIES OF PLANTS**

Geneva

**TECHNICAL COMMITTEE****Forty-Ninth Session  
Geneva, March 18 to 20, 2013**

## PARTIAL REVISION OF THE TEST GUIDELINES FOR TOMATO (DOCUMENT TG/44/11)

*Document prepared by the Office of the Union*

1. The Technical Working Party for Vegetables (TWV), at its forty-fifth session, held in Monterey, United States of America, from July 25 to 29, 2011, agreed to propose to the Technical Committee (TC) to adopt a partial revision of the Test Guidelines for Tomato (document TG/44/11) in order to include:

- (a) a revised format for disease resistance characteristics according to the explanations for disease resistance characteristics in Test Guidelines; as set out in document TGP/12/2 Draft 2 "Guidance on Certain Physiological Characteristics", Section 2.4; and
- (b) a gene-specific marker method for examination of resistance to Tomato Spotted Wilt topovirus (TSWV) - Race 0.

2. The TC, at its forty-eighth session held in Geneva from March 26 to 28, 2012, noted that, in response to a number of technical questions concerning disease resistance, raised by interested experts after the TWV session, it had been agreed by the TWV Chairperson, former TWV Chairperson, and the Leading Expert to consider a new draft of the partial revision of the Test Guidelines for Tomato at the forty-sixth session of the TWV (see document TC/48/22 "Report on Conclusions", paragraph 147).

3. The TWV at its forty-sixth session, held near the city of Venlo, Netherlands, June 11 to 15, 2012, considered document TWV/46/19 and agreed to propose the partial revision of the Test Guidelines for Tomato (document TG/44/11) as set out in the Annexes to this document:

ANNEX I Proposal to correct diseases names in Chapters: 5.3, 7, 8 and 10

ANNEX II Inclusion of a revised format for disease resistance characteristics according to the explanations for disease resistance characteristics in Test Guidelines; as set out in document TGP/12/2 Draft 2 "Guidance on Certain Physiological Characteristics", Section 2.4 (current and proposed new wording are presented on opposite pages)

ANNEX III Addition of Literature References to Chapter 9: Literature;

4. The partial revision to document TG/44/11 would be adopted as document TG/44/11 Rev..

[Annex follows]

Proposal to Correct Disease Names in Chapters: 5.3, 7 ,8 and 10. TQ*Current wording:*

|            | English  | français  | deutsch  | español  | Example Varieties<br>Exemples<br>Beispielssorten<br>Variedades ejemplo | Note/<br>Nota |
|------------|--|---|--|--|--|---------------|
| 49.<br>(+) | VG<br>Resistance to <i>Fusarium oxysporum</i> f. sp. <i>radicis lycopersici</i> (Forl) | Résistance à <i>Fusarium oxysporum</i> f. sp. <i>radicis lycopersici</i> (Forl) | Resistenz gegen <i>Fusarium oxysporum</i> f. sp. <i>radicis lycopersici</i> (Forl) | Resistencia a <i>Fusarium oxysporum</i> f. sp. <i>radicis lycopersici</i> (Forl) |  |               |

*Proposed new wording:*

|            |  |   |  |  |  |  |
|------------|--|---|--|--|--|--|
| 49.<br>(+) | VG<br>Resistance to <i>Fusarium oxysporum</i> f. sp. <i>radicis-lycopersici</i> (Forl) | Résistance à <i>Fusarium oxysporum</i> f. sp. <i>radicis-lycopersici</i> (Forl) | Resistenz gegen <i>Fusarium oxysporum</i> f. sp. <i>radicis-lycopersici</i> (Forl) | Resistencia a <i>Fusarium oxysporum</i> f. sp. <i>radicis-lycopersici</i> (Forl) |  |  |
|------------|--|---|--|--|--|--|

*Current wording:*

|            | English  | français   | deutsch  | español  | Example Varieties<br>Exemples<br>Beispielssorten<br>Variedades ejemplo | Note/<br>Nota |
|------------|--|--|--|--|--|---------------|
| 51.<br>(+) | VG<br>Resistance to Tomato Mosaic Tobamovirus (ToMV) | Résistance au virus de la mosaïque de la tomate (ToMV) | Resistenz gegen das Tomatenmosaikvirus, Tobamovirus (ToMV) | Resistencia al virus del mosaico del tomate (ToMV) |  |               |

*Proposed new wording:*

|            |  |  |   |  |  |  |
|------------|--|--|---|--|--|--|
| 51.<br>(+) | VG<br>Resistance to Tomato mosaic virus (ToMV) | Résistance au virus de la mosaïque de la tomate (ToMV) | Resistenz gegen das Tomatenmosaikvirus (ToMV) | Resistencia al virus del mosaico del tomate (ToMV) |  |  |
|------------|--|--|---|--|--|--|

*Current wording:*

|            | English                                | français                        | deutsch                            | español                          | Example Varieties<br>Exemples<br>Beispielssorten<br>Variedades ejemplo | Note/<br>Nota |
|------------|--|---------------------------------|------------------------------------|----------------------------------|--|---------------|
| 54.<br>(+) | VG<br>Resistance to <i>Stemphylium</i> | Résistance à <i>Stemphylium</i> | Resistenz gegen <i>Stemphylium</i> | Resistencia a <i>Stemphylium</i> |  |               |

*Proposed new wording:*

|            |  |   |  |  |  |  |
|------------|--|---|--|--|--|--|
| 54.<br>(+) | VG<br>Resistance to <i>Stemphylium</i> spp. (Ss) | Résistance à <i>Stemphylium</i> spp. (Ss) | Resistenz gegen <i>Stemphylium</i> spp. (Ss) | Resistencia a <i>Stemphylium</i> spp. (Ss) |  |  |
|------------|--|---|--|--|--|--|

*Current wording:*

|            | English   | français  | deutsch  | español  | Example Varieties<br>Exemples<br>Beispielssorten<br>Variedades ejemplo | Note/<br>Nota |
|------------|---|---|--|--|--|---------------|
| 57.<br>(+) | <b>VG</b><br><b>Resistance to Tomato Yellow Leaf Curl Begomovirus (TYLCV)</b> | <b>Résistance au bégomovirus des feuilles jaunes en cuillère de la tomate (TYLCV)</b> | <b>Resistenz gegen gelbes Tomatenblattrollvirus, Begomovirus (TYLCV)</b> | <b>Resistencia a Begomovirus del rizado amarillo de la hoja del tomate (TYLCV)</b> |  |               |

*Proposed new wording:*

|            |   |   |   |  |  |  |
|------------|---|---|---|--|--|--|
| 57.<br>(+) | <b>VG</b><br><b>Resistance to Tomato yellow leaf curl virus (TYLCV)</b> | <b>Résistance au virus des feuilles jaunes en cuillère de la tomate (TYLCV)</b> | <b>Resistenz gegen gelbes Tomatenblattrollvirus (TYLCV)</b> | <b>Resistencia a virus del rizado amarillo de la hoja del tomate (TYLCV)</b> |  |  |
|------------|---|---|---|--|--|--|

*Current wording:*

|            | English   | français   | deutsch  | español  | Example Varieties<br>Exemples<br>Beispielssorten<br>Variedades ejemplo | Note/<br>Nota |
|------------|---|--|--|--|--|---------------|
| 58.<br>(+) | <b>VG</b><br><b>Resistance to Tomato Spotted Wilt Tospovirus (TSWV)</b> | <b>Résistance au virus de la tache bronzée de la tomate (TSWV)</b> | <b>Resistenz gegen das Tomatenbronzenfleckenvirus, Tospovirus (TSWV)</b> | <b>Resistencia a Tospovirus del bronceado de tomate (TSWV)</b> |  |               |
|            | <b>- Race 0</b>   | <b>- Pathotype 0</b>   | <b>- Pathotyp 0</b>  | <b>- Raza 1</b>  |  |               |

*Proposed new wording:*

|            |  |  |  |   |  |  |
|------------|--|--|--|---|--|--|
| 58.<br>(+) | <b>VG</b><br><b>Resistance to Tomato spotted wilt virus (TSWV)</b> | <b>Résistance au virus de la tache bronzée de la tomate (TSWV)</b> | <b>Resistenz gegen das Tomatenbronzenfleckenvirus (TSWV)</b> | <b>Resistencia a virus del bronceado de tomate (TSWV)</b> |  |  |
|            | <b>- Race 0</b>  | <b>- Pathotype 0</b>   | <b>- Pathotyp 0</b>  | <b>- Raza 1</b>   |  |  |

*Current wording:*

|            | English   | français   | deutsch  | español   | Example Varieties<br>Exemples<br>Beispielssorten<br>Variedades ejemplo | Note/<br>Nota |
|------------|---|--|--|---|--|---------------|
| 61.<br>(+) | <b>VG</b><br><b>Resistance to Tomato Torrado Virus (ToTV)</b> | <b>Résistance au virus Tomato Torrado (ToTV)</b> | <b>Resistenz gegen Tomato Torrado Virus (ToTV)</b> | <b>Resistencia al virus del torrado del tomate (ToTV)</b> |  |               |

*Proposed new wording:*

|            |   |  |  |   |  |  |
|------------|---|--|--|---|--|--|
| 61.<br>(+) | <b>VG</b><br><b>Resistance to Tomato torrado virus (ToTV)</b> | <b>Résistance au virus tomato torrado (ToTV)</b> | <b>Resistenz gegen Tomato Torrado Virus (ToTV)</b> | <b>Resistencia al virus del torrado del tomate (ToTV)</b> |  |  |
|------------|---|--|--|---|--|--|

Proposal to Include a Revised Format for Disease Resistance Characteristics  
(Current and Proposed New Wording are presented on opposite pages)

*Current wording:*

Ad. 46: Resistance to *Meloidogyne incognita* (Mi)

Method

Maintenance of strain

Type of medium: on roots of susceptible varieties

Special conditions avoid rotting of roots

Execution of test

Temperature: not over 28° C

Growing method: preferably in the greenhouse

Method of inoculation: plants are sown in infested soil

Duration of test

- from sowing to inoculation: inoculation before sowing,
- from inoculation to reading: 30 to 45 days

Number of plants tested: 10 to 20

Remarks: avoid rotting of roots avoid high temperature

Notation: number of root knots contaminated with eggs and root deformation

Standard varieties: susceptible: Clairvil, Casaque Rouge  
moderately resistant: Madyta, Vinchy  
highly resistant: Anabel, Anahu, F1 Anahu x Monalbo

*Proposed new wording:*

Ad 46: Resistance to *Meloidogyne incognita* (Mi)

|   |   |
|---|---|
| 1. Pathogen .....   | <i>Meloidogyne incognita</i>  |
| 3. Host species .....   | <i>Solanum lycopersicum</i>   |
| 4. Source of inoculum .....   | Naktuinbouw (NL <sup>1</sup> ) or GEVES <sup>2</sup> (FR)   |
| 5. Isolate .....  | non-resistance breaking   |
| 6. Establishment isolate identity .....   | use rootstock or tomato standards   |
| 7. Establishment pathogenicity .....  | use susceptible rootstock or tomato standard  |
| 8. Multiplication inoculum  |   |
| 8.1 Multiplication medium .....   | living plant  |
| 8.2 Multiplication variety .....  | preferably resistant to powdery mildew  |
| 8.3 Plant stage at inoculation .....  | see 10.3  |
| 8.5 Inoculation method .....  | see 10.4  |
| 8.6 Harvest of inoculum .....   | root systems are cut with scissors into pieces of about 1 cm length   |
| 8.7 Check of harvested inoculum .....   | visual check for presence of root knots   |
| 8.8 Shelf life/viability inoculum .....   | 1 day   |
| 9. Format of the test .....   |   |
| 9.1 Number of plants per genotype ....  | 20 plants   |
| 9.2 Number of replicates .....  | 1 replicate   |
| 9.3 Control varieties .....   |   |
| Susceptible: .....  | Clairvil, Casaque Rouge   |
| Moderately resistant : .....  | Campeon, Madyta, Vinchy   |
| Highly resistant: .....   | Anabel, Anahu, Anahu x Casaque Rouge  |
| 9.4 Test design .....   | include standard varieties  |
| 9.5 Test facility .....   | greenhouse or climate room  |
| 9.6 Temperature .....   | not over 28° C  |
| 9.7 Light .....   | at least 12 h per day   |
| 10. Inoculation .....   |   |
| 10.1 Preparation inoculum .....   | small pieces of diseased root mixed with soil<br>mix soil and infested root pieces                                    |
| 10.2 Quantification inoculum .....  | soil: root ratio = 8:1, or depending on experience  |
| 10.3 Plant stage at inoculation .....   | seed, or cotyledons   |
| 10.4 Inoculation method .....   | plants are sown in infested soil or contamination of soil after sowing when plantlets are at cotyledon stage          |
| 10.7 Final observations .....   | 28 to 45 days after inoculation   |
| 11. Observations .....  |   |
| 11.1 Method .....   | root inspection   |
| 11.2 Observation scale .....  | Symptoms:<br>galling, root malformation,<br>growth reduction, plant death   |
| 11.3 Validation of test .....   | evaluation of variety resistance should be calibrated with results of resistant and susceptible controls on standards |
| 12. Interpretation of test results in comparison with control varieties:  |   |
| To consider that resistant varieties may have a few plants with a few galls. These are not considered as off-types. |   |
| absent (susceptible) .....  | [1] growth strongly reduced, high gall count  |
| intermediate (moderately resistant) .....   | [2] medium growth reduction, medium gall count  |
| present (highly resistant) .....  | [3] present; no growth reduction, no galls  |
| 13. Critical control points:  |   |
| Avoid rotting of roots; high temperature causes breakdown of resistance.  |   |

<sup>1</sup> Naktuinbou; [resistentie@naktuinbouw.nl](mailto:resistentie@naktuinbouw.nl)

<sup>2</sup> GEVES; [Valerie.GRIMAULT@geves.fr](mailto:Valerie.GRIMAULT@geves.fr)

*Current wording:*

Ad. 47: Resistance to *Verticillium* sp.(Va and Vd)

#### Method

##### Maintenance of strains

Race 0 represented by strain Toreilles 4-1-4-1 is used. Race 0 is the common race defined by its ability to infect plants with the Ve gene.

Long term storage of strains: conidia suspended in glycerol solution at -80°C.  
Strain can be subcultured on Potato Dextrose Agar (PDA) medium.

##### Execution of test

##### Growth stage of plants

Plants are grown in greenhouse or growth chamber. Inoculation can be done from the cotyledon stage (first leaves emerging) to 2 expanded leaves stage.

The following varieties can be used as controls. As a minimum, there should be one resistant and one susceptible control in the test. The heterozygous variety will help interpretation of results in case of aggressive test. Clarion could be interesting to add to susceptible controls as it is less susceptible and could also help to check the inoculation pressure of the test. These 2 varieties are optional.

| Standard variety         | Vd:0 |
|--------------------------|------|
| Marmande verte, Flix     | S    |
| Clarion                  | s    |
| Monalbo x Marmande verte | RH   |
| Monalbo, Elias           | R    |

R resistance present; no symptoms  
RH resistance present; sometimes very weak symptoms  
s resistance absent; weak symptoms  
S resistance absent; clear symptoms

##### Temperature:

Test performed under controlled conditions at 20 to 22°C.

##### Inoculum:

*Verticillium* sp. is grown on liquid Czapek Dox Broth or S of Messiaen media for 3 to 7 days in the dark, at 20 to 25°C with aeration. Spores are harvested and adjusted to 10<sup>6</sup>sp/ml.

##### Method of inoculation

Plantlets are harvested, roots are cut and soaked for 5 to 15 min in the inoculum suspension. Plantlets are then transplanted in soil.

##### Duration of test

At least 33 days from sowing to notation.

##### Number of plants tested:

At least 20 plants.

##### Notation:

25-30 days after inoculation.

##### Notation scale and interpretation of results:

R: no symptoms

S: chlorosis in the lower leaves, growth reduced and brown vessels or growth not reduced and brown vessels.

Analysis of results should be calibrated with results on R and S controls.

*Proposed new wording:*

Ad 47: Resistance to *Verticillium* sp. (Va and Vd)

|   |  |
|---|--|
| 1. Pathogen .....   | <i>Verticillium dahliae</i> or <i>Verticillium albo-atrum</i>  |
| 3. Host species .....   | <i>Solanum lycopersicum</i>  |
| 4. Source of inoculum .....   | Naktuinbouw <sup>3</sup> (NL) and GEVES <sup>4</sup> (FR)  |
| 5. Isolate .....  | Race 0 (e.g. strain Toreilles 4-1-4-1)   |
| 8. Multiplication inoculum .....  |  |
| 8.1 Multiplication medium .....   | Potato Dextrose Agar, Agar Medium "S" of Messiaen  |
| 8.4 Inoculation medium.....   | water (for scraping agar plates) or Czapek Dox broth (3-7 d-old  |
| .....   | aerated culture at 20-25°C, in darkness)   |
| 8.6 Harvest of inoculum .....   | filter through double muslin cloth   |
| 8.7 Check of harvested inoculums.....                                   | spore count; adjust to 10 <sup>6</sup> per ml  |
| 8.8 Shelf life/viability inoculums.....                                 | 1 d at 4°C   |
| 9. Format of the test.....  |  |
| 9.1 Number of plants per genotype.....                                  | 35 seeds for 24 plants   |
| 9.2 Number of replicates.....   | 1 replicate  |
| 9.3 Control varieties.....  |  |
| Susceptible .....   | Flix, Marmande verte, Clarion, Santonio, Anabel  |
| Resistant .....   | Monalbo, Elias, Monalbo x Marmande verte, Daniela,<br>Marmande VR  |
| 9.4 Test design.....  | 20 plants inoculated at least, 2 blanks at least   |
| 9.5 Test facility.....  | greenhouse or climate room   |
| 9.6 Temperature.....  | optimal 20-25°C, 20-22°C after inoculation   |
| 9.7 Light.....  | 12 h or longer   |
| 10. Inoculation .....   |  |
| 10.1 Preparation inoculums.....   | aerated, liquid culture (8.4)  |
| 10.2 Quantification inoculums.....                                      | count spores, adjust to 10 <sup>6</sup> per ml   |
| 10.3 Plant stage at inoculation.....                                    | cotyledon to 3rd leaf  |
| 10.4 Inoculation method.....  | roots are immersed for 4 to 15 min in spore suspension.  |
| 10.7 Final observations.....  | 14-33 d after inoculation  |
| 11. Observations.....   |  |
| 11.1 Method.....  | visual   |
| 11.2 Observation scale.....   | growth retardation, wilting, chlorosis, and vessel browning  |
| 11.3 Validation of test.....  | evaluation of variety resistance should be calibrated with<br>results of resistant and susceptible controls. |
| 12. Interpretation of test results in comparison with control varieties |  |
| absent .....  | [1] severe symptoms  |
| present.....  | [9] no or mild symptoms  |

13. Critical control points:

All symptoms may be present in resistant varieties, but the severity will be distinctly less than in susceptible varieties. Usually resistant varieties will show significantly less growth retardation than susceptible varieties. Observation of vessel browning is important for diagnosis. Usually, vessel browning will not extend to the 1<sup>st</sup> leaf in resistant varieties. Many hybrid varieties are heterozygous and appear to have mild symptoms in the biotest.

<sup>3</sup> Naktuinbouw: [resistentie@naktuinbouw.nl](mailto:resistentie@naktuinbouw.nl)

<sup>4</sup> GEVES; [Valerie.GRIMAULT@geves.fr](mailto:Valerie.GRIMAULT@geves.fr)

*Current wording:*

Ad. 48.1 + 48.2 + 48.3: Resistance to *Fusarium oxysporum* f. sp. *lycopersici* (Fol) -Race 0 (ex 1), Race 1 (ex 2) and Race 2 (ex 3)

Method

Maintenance of strains

Long term storage of strains: at -80°C in 20% glycerol.  
Race 0 (ex 1) represented by strains Orange 71 or PRI 20698 or Fol 071 and race 1 represented by strains 4152 (more aggressive) or PRI40698 or RAF 70 (less aggressive) are used.  
Strains can be multiplied on PDA or S of Messiaen media.

Execution of test

Growth stage of plants

Plants are grown in greenhouse or growth chamber for 10 to 18 days (cotyledons to first leaf stages).

The following varieties are used as controls. Each line will be represented by at least one variety which can be chosen in the varieties indicated; the resistance phenotype to the two pathotypes of Fol is indicated. The heterozygous variety has a resistance phenotype usually weaker than in homozygous lines. This weak resistance can be used to calibrate the borderline between resistance and susceptibility. The heterozygous control for Fol:1 is optional.

| <u>Controls for Fol:0 resistance test</u> | Fol:0 | Fol:1* |
|---|-------|--------|
| Marmande, Marmande verte, Resal           | S     | S      |
| Marporum x Marmande verte (heterozygous)  | R     | S      |
| Marporum, Larissa                         | R     | S      |
| Motelle, Gourmet, Mohawk                  | R     | R      |

\* For information

| <u>Controls for Fol:1 resistance test</u> | Fol:0* | Fol:1 |
|---|--------|-------|
| Cherry Belle, Roma, Marmande verte        | S      | S     |
| Ranco**, Marporum                         | R      | S     |
| Motelle x Marmande verte                  | R      | R     |
| Tradiro, Odisea                           | R      | R     |

\* For information

\*\* For Ranco: weak resistance to Fol:0 with many escapes

R = resistance present  
S = resistance absent

Temperature:

Test performed in climatic chambers or greenhouse at 24-28°C. In case of aggressive test, temperature can be decreased to 20-24°C.

Inoculum:

*Fusarium oxysporum* f. sp. *lycopersici* is grown on PDA or S of Messiaen media or in aerated Czapek-Dox liquid cultures for 7 to 10 days. Spores are harvested and adjusted to 10<sup>6</sup>sp/ml for strains grown on media. In case of very aggressive isolate, inoculum concentration can be decreased.

Method of inoculation

Soaking of roots (cutting of roots optional) and of hypocotyls axis for 5 to 15 min in the inoculum suspension and transplantation of inoculated plantlets in soil.

Duration of test

At least 28 days from sowing to notation.

Number of plants tested:

At least 20 plants.

Notation:

At least 21 days after inoculation.

Notation scale:

4 classes:

- 0: no symptoms,
- 1: external healthy aspect of plant (without growth reduction) with brown vessels (sometimes extending above cotyledons, generally remaining below cotyledons),
- 2: growth reduction and brown vessels above cotyledons,
- 3: dead plant.

Interpretation of scale:

Generally, 0 and 1 are equivalent to resistant, 2 and 3 are susceptible but analysis of results should be calibrated with results of R and S controls.

Proposed new wording:

Ad 48: Resistance to *Fusarium oxysporum* f. sp. *lycopersici* (Fol)

|  |   |
|--|---|
| 1. Pathogen .....  | <i>Fusarium oxysporum</i> f. sp. <i>lycopersici</i>   |
| 3. Host species .....  | <i>Solanum lycopersicum</i>   |
| 4. Source of inoculum .....  | Naktuinbouw <sup>5</sup> (NL) and GEVES <sup>6</sup> (FR)   |
| 5. Isolate .....   | Race 0 (ex 1) (e.g. strains Orange 71 or PRI 20698 or Fol 071 1 (ex 2) (e.g. strains 4152 or PRI40698 or RAF 70 and 2 (ex 3)<br>Individual strains may vary in pathogenicity<br>use differential varieties (see 9.3)<br>on susceptible tomato varieties |
| 6. Establishment isolate identity .....  |   |
| 7. Establishment pathogenicity .....   |   |
| 8. Multiplication inoculum.....  |   |
| 8.1 Multiplication medium .....  | Potato Dextrose Agar, Medium "S" of Messiaen  |
| 8.4 Inoculation medium .....   | water for scraping agar plates or Czapek-Dox culture medium<br>(7 d-old aerated culture)  |
| 8.6 Harvest of inoculum .....  | filter through double muslin cloth  |
| 8.7 Check of harvested inoculum.....   | spore count; adjust to 106 per ml   |
| 8.8 Shelf-life/viability inoculum .....  | 4-8 h, keep cool to prevent spore germination   |
| 9. Format of the test.....   |   |
| 9.1 Number of plants per genotype .....  | at least 20 plants  |
| 9.2 Number of replicates.....  | 1 replicate   |
| 9.3 Control varieties for the test with<br>race 0 (ex 1)   |   |
| Susceptible .....  | Marmande, Marmande verte, Resal   |
| Resistant for race 0 only .....  | Marporum, Larissa, "Marporum x Marmande verte", Marsol, Anabel  |
| Resistant for race 0 and 1 .....   | Motelle, Gourmet, Mohawk  |
| Control varieties for the test with<br>race 1 (ex 2)   |   |
| Susceptible .....  | Marmande verte, Cherry Belle, Roma  |
| Resistant for race 0 only .....  | Marporum, Ranco   |
| Resistant for race 0 and 1 .....   | Tradiro, Odisea   |
| Remark: .....  | Ranco is slightly less resistant than Tradiro   |
| Control varieties for the test with<br>race 2 (ex 3)   |   |
| Susceptible for race 0, 1 and 2 .....  | Marmande verte, Motelle, Marporum   |
| Resistant for race 0, 1 and 2.....   | Tributes, Murdoch, Marmande verte x Florida   |
| 9.4 Test design .....  | >20 plants; e.g. 35 seeds for 24 plants, including 2 blanks   |
| 9.5 Test facility.....   | glasshouse or climate room  |
| 9.6 Temperature .....  | 24-28°C (severe test, with mild isolate)<br>20-24°C (mild test, with severe isolate)  |
| 9.7 Light .....  | 12 hours per day or longer  |
| 9.8 Season.....  | all seasons   |
| 9.9 Special measures .....   | slightly acidic peat soil is optimal;<br>keep soil humid but avoid water stress   |
| 10. Inoculation.....   |   |
| 10.1 Preparation inoculums .....   | aerated Messiaen or PDA or Agar Medium S of Messiaen or<br>Czapek Dox culture or scraping of plates   |
| 10.2 Quantification inoculums.....   | spore count, adjust to 106 spores per ml,<br>lower concentration for a very aggressive isolate  |
| 10.3 Plant stage at inoculation.....   | 10-18 d, cotyledon to first leaf  |
| 10.4 Inoculation method.....   | roots and hypocotyls are immersed in spore suspension<br>for 5-15 min; trimming of roots is an option   |
| 10.7 Final observations.....   | 14-21 days after inoculation  |
| 11. Observations.....  |   |
| 11.1 Method .....  | visual  |
| 11.2 Observation scale .....   | Symptoms:<br>growth retardation, wilting, yellowing,<br>vessel browning extending above cotyledon   |
| 11.3 Validation of test.....   | evaluation of variety resistance should be calibrated with results of resistant and<br>susceptible controls. Standards near borderline R/S will help to compare<br>between labs.  |
| 12. Interpretation of test results in comparison with control varieties  |   |
| absent.....  | [1] severe symptoms   |
| present .....  | [9] mild or no symptoms   |
| 13. Critical control points  |   |
| Test results may vary slightly in inoculum pressure due to differences in isolate, spore concentration, soil humidity and temperature. |   |

<sup>5</sup> Naktuinbouw: resistantie@naktuinbouw.nl

<sup>6</sup> GEVES; Valerie.GRIMAULT@geves.fr

*Current wording:*

Ad. 49: Resistance to *Fusarium oxysporum* f. sp. *radicis lycopersici* (Forl)

Method

Maintenance of race

Type of medium: on PDA or synthetic medium (according to Messiaen)

Special conditions: fridge 4° C

Execution of test

Growth stage of plants: appearance of third leaf

Temperature: day: 22° C, night: 16° C

Light: 14 hours

Growing method: climate room or glasshouse

Method of inoculation: soaking of roots and of hypocotyl axis for five minutes in the inoculum.

Duration of test

- from sowing to inoculation: 18 to 20 days
- from inoculation to reading: 10 days

Number of plants tested: 10 to 20 plants

Remarks: need for frequent renewal of races because of loss of pathogenicity

Standard varieties:

- susceptible: Motelle
- resistant:
  - Momor (homozygote)
  - F1 Momor x Motelle (heterozygote)
  - the Frl gene does not completely control the disease in the heterozygous stage.

*Proposed new wording:*

Ad 49: Resistance to *Fusarium oxysporum* f. sp. *radicis-lycopersici* (Forl)

|   |   |
|---|---|
| 1. Pathogen .....   | <i>Fusarium oxysporum</i> f. sp. <i>radicis-lycopersici</i>   |
| 3. Host species .....   | <i>Solanum lycopersicum</i>   |
| 4. Source of inoculum .....   | Naktuinbouw <sup>7</sup> (NL) and GEVES <sup>8</sup> (FR)   |
| 5. Isolate .....  |   |
| 7. Establishment pathogenicity .....  | symptoms on susceptible tomato  |
| 8.1 Multiplication medium .....   | Multiplication inoculum<br>Potato Dextrose Agar or Medium agar "S" of Messiaen  |
| 8.4 Inoculation medium .....  | water for scraping agar plates or<br>Czapek-Dox (7 d-old aerated culture)   |
| 8.6 Harvest of inoculum .....   | filter through double muslin cloth  |
| 8.7 Check of harvested inoculum .....   | spore count; adjust to 10 <sup>6</sup> per ml   |
| 8.8 Shelf life/viability inoculum .....   | 4-8 h, keep cool to prevent spore germination   |
| 9. Format of the test .....   |   |
| 9.1 Number of plants per genotype .....   | at least 20 plants  |
| 9.2 Number of replicates .....  | 1 replicate   |
| 9.3 Control varieties .....   |   |
| Susceptible: .....  | Motelle, Moneymaker   |
| Resistant: .....  | Momor, "Momor x Motelle"  |
| Remark: .....   | "Momor x Motelle" has slightly weaker resistance than Momor   |
| 9.4 Test design .....   | >20 plants; e.g. 35 seeds for 24 plants, including 2 blanks   |
| 9.5 Test facility .....   | glasshouse or climate room  |
| 9.6 Temperature .....   | 24-28°C (severe test, with mild isolate)<br>17-24°C (mild test, with severe isolate)  |
| 9.7 Light .....   | at least 12 hours per day   |
| 9.8 Season .....  | all seasons   |
| 9.9 Special measures .....  | slightly acidic peat soil is optimal;<br>keep soil humid but avoid water stress   |
| 10. Inoculation .....   |   |
| 10.1 Preparation inoculum .....   | aerated culture or scraping of plates   |
| 10.2 Quantification inoculum .....  | spore count, adjust to 10 <sup>6</sup> spores per ml  |
| 10.3 Plant stage at inoculation .....   | 12-18 d, cotyledon to third leaf  |
| 10.4 Inoculation method .....   | roots and hypocotyls are immersed in spore suspension<br>for 5-15 min   |
| 10.7 Final observations .....   | 10-21 days after inoculation  |
| 11. Observations .....  |   |
| 11.1 Method .....   | visual; a few plants are lifted at the end of the test  |
| 11.2 Observation scale .....  | Symptoms:<br>plant death, growth retardation caused by root degradation<br>root degradation, necrotic pinpoint and necrotic lesions on<br>stems |
| 11.3 Validation of test .....   | evaluation of variety resistance should be calibrated with results<br>of resistant and susceptible controls                                     |
| 12. Interpretation of test results in comparison with control varieties   |   |
| absent .....  | [1] symptoms  |
| present .....   | [9] no symptoms   |
| 13. Critical control points:  |   |
| Temperature should never exceed 27°C during the test period; frequent renewal of races may be needed because of loss of pathogenicity |   |

<sup>7</sup> Naktuinbouw: [resistentie@naktuinbouw.nl](mailto:resistentie@naktuinbouw.nl)

<sup>8</sup> GEVES; [Valerie.GRIMAULT@geves.fr](mailto:Valerie.GRIMAULT@geves.fr)

*Current wording:*

Ad. 50.1 – 50.6 Resistance to *Fulvia fulva* (Ff) (ex *Cladosporium fulvum*)

Method

Maintenance of races

Type of medium: PDA or synthetic medium

Special conditions: subculturing of isolates

Execution of test

Growth stage of plants: 3 leaves expanded

Temperature: day: 24° C, night: 16° C

Light: 12 hours

Growing method: in climate room, highest possible humidity, with reduced growth a few days before inoculation by irrigation of roots with ALAR 85 (daminazoide), or in glasshouse with high humidity, for example under a polyethylene cover.

Method of inoculation: spraying of a solution with the fungus on leaves.

Duration of test

- from sowing to inoculation: 22 to 25 days  
- from inoculation to reading: 20 to 25 days

Number of plants tested: 30 plants

Remarks: the level of expression of symptoms may vary between plants due to complex resistance genetics

Standard varieties:

- susceptible: Monalbo
- resistant: has to be chosen with the concerned alleles
  - cf1: Stirling Castle
  - cf2: Vetomold
  - cf3: V 121
  - cf4: Purdue 135
  - cf5: IVT 1149
  - cf2 cf4: Vagabond
  - cf2 cf5: F1 "Vetomold x IVT 1149"
  - cf2 cf4 cf5: F1 "Vagabond x IVT 1149"
  - cf6: F 77-38
  - cf9: IVT 1154

Race 0: Angela, Estrella, Sonatine, Sonato, Vemone  
Group A: Angela, Estrella, Sonatine, Sonato  
Group B: Angela, Estrella, Sonatine, Sonato, Vemone  
Group C: Angela, Estrella, Sonatine  
Group D: Estrella, Sonatine, Vemone  
Group E: Sonatine

*Proposed new wording:*

Ad 50: Resistance to *Fulvia fulva* (Ff) (ex *Cladosporium fulvum*)

|  |   |
|--|---|
| 1. Pathogen .....  | <i>Fulvia fulva</i> (ex <i>Cladosporium fulvum</i> )  |
| 3. Host species .....  | <i>Solanum lycopersicum</i>   |
| 4. Source of inoculum .....  | Naktuinbouw <sup>9</sup> (NL) or GEVES <sup>10</sup> (FR)   |
| 5. Isolate .....   | Race group 0, A, B, C, D, and E   |
| 6. Establishment isolate identity .....  | with genetically defined differentials from GEVES (FR)<br>A breaks Cf-2, B Cf-4, C Cf-2&4, D Cf-5, E Cf-2&4&5<br>symptoms on susceptible tomato   |
| 7. Establishment pathogenicity .....   |   |
| 8. Multiplication inoculum .....   | Potato Dextrose Agar or Malt Agar or a synthetic medium   |
| 8.1 Multiplication medium .....  | 4 hours, keep cool  |
| 8.8 Shelf life/viability inoculum.....   |   |
| 9. Format of the test.....   |   |
| 9.1 Number of plants per genotype .....  | more than 20 plants   |
| 9.2 Number of replicates.....  | 1 replicate   |
| 9.3 Control varieties.....   |   |
| Susceptible: .....   | Monalbo, Money maker  |
| Resistant for race 0:.....   | Angela, Estrella, Sonatine, Sonato, Vemone, Vagabond, IVT<br>1149, Vagabond x IVT 1149, IVT 1154  |
| Resistant for race group A: .....  | Angela, Estrella, Sonatine, Sonato  |
| Resistant for race group B: .....  | Angela, Estrella, Sonatine, Sonato, Vemone  |
| Resistant for race group C: .....  | Angela, Estrella, Sonatine  |
| Resistant for race group D: .....  | Estrella, Sonatine, Vemone  |
| Resistant for race group E: .....  | Sonatine, Jadviga, Rhianna, IVT 1154  |
| 9.5 Test facility .....  | glasshouse or climate room  |
| 9.6 Temperature .....  | day: 22° C, night: 20° or day: 25° C, night 20° C   |
| 9.7 Light .....  | 12 hours or longer  |
| 9.9 Special measures .....   | depending on facility and weather, there may be a need to<br>raise the humidity<br>e.g. humidity tent closed 3-4 days after inoculation<br>and after this, 66% until 80% closed during day, until end |
| 10. Inoculation .....  |   |
| 10.1 Preparation inoculum.....   | prepare evenly colonized plates, e.g. 1 for 36 plants;<br>remove spores from plate by scraping with water with Tween20;<br>filter through double muslin cloth   |
| 10.2 Quantification inoculum .....   | count spores; adjust to 10 <sup>5</sup> spores per ml or more   |
| 10.3 Plant stage at inoculation .....  | 19-20 d (incl. 12 d at 24°), 2-3 leaves   |
| 10.4 Inoculation method .....  | spray on dry leaves   |
| 10.7 Final observations .....  | 14 days after inoculation   |
| 11. Observations.....  |   |
| 11.1 Method.....   | visual inspection of abaxial side of inoculated leaves  |
| 11.2 Observation scale .....   | Symptom: velvety, white spots   |
| 11.3 Validation of test .....  | evaluation of variety resistance should be calibrated with results<br>of resistant and susceptible controls   |
| 12. Interpretation of test results in comparison with control varieties                      |   |
| absent .....   | [1] symptoms  |
| present.....   | [9] no symptoms   |
|  | Excessively high humidity may cause rugged brown spot on all<br>leaves. These are not to be considered as off-types.  |
| 13. Critical control points:   |   |
| Ff spores have a variable size and morphology. Small spores are also viable.                 |   |
| Fungal plates will gradually become sterile after 6-10 weeks. Store good culture at -80°C.   |   |
| For practical purposes, it is not possible to keep plants longer than 14 days inside a tent. |   |

<sup>9</sup> Naktuinbouw: resistentie@naktuinbouw.nl

<sup>10</sup> GEVES; Valerie.GRIMAULT@geves.fr

*Current wording:*

Ad. 51.1 – 51.3: Resistance to Tomato Mosaic Tobamovirus (ToMV)- Strains 0, 1 and 2

Method

Maintenance of strains

Strains are long term stored as desiccated leaves below 10°C.  
Race 0 represented by isolate INRA Avignon 6-5-1-1 (aucuba mosaic strain) is used.  
Virus should be multiplied on the susceptible control before being used for inoculation of the test.

Execution of test

Growth stage of plants

Plants are grown in greenhouse or growth chamber until cotyledons (first leaves emerging) to two expanded leaves have appeared.

Within each test at least one resistant and one susceptible standard variety is included.

The following varieties are used as controls. Each line will be represented by at least one resistance phenotype which can be chosen from the varieties indicated; the resistance phenotype to the 3 pathotypes of ToMV is indicated. Mobaci and Moperou will allow checking the pathotype identity of the virus. Monalbo x Momor will help the interpretation of the distinct resistance phenotype with necrosis.

| Variety           | Resistance phenotype |        |        |
|-------------------|----------------------|--------|--------|
|                   | ToMV:0               | ToMV:1 | ToMV:2 |
| Marmande, Monalbo | S                    | S      | S      |
| Mobaci            | R                    | S      | R      |
| Moperou           | R                    | R      | S      |
| Monalbo x Momor   | RN                   | RN     | RN     |
| Momor, Gourmet    | R                    | R      | R      |

R = resistance present; no symptoms

RN = resistance present; a variable proportion of plants showing some or extensive necrosis; all other plants have no symptoms.

S = resistance absent; mosaic symptoms

Temperature:

Test performed in climatic chambers or greenhouse at 24 to 26°C. At higher temperatures, resistance can break down.

Inoculum and method of inoculation

Mechanical inoculation by rubbing cotyledons (first leaves emerging) or two expanded leaves with an inoculum solution consisting of symptomatic leaves grinded in a buffer with carborundum added. Leaves can be rinsed after inoculation. Light is important for symptom expression.

Duration of test

24 to 42 days from sowing to notation.

Number of plants tested:

At least 20 plants.

Notation:

12-21 days after inoculation when symptoms are well developed on susceptible control.

Notation scale and interpretation results:

R: without symptoms or with necrosis (necrosis can be observed on plants heterozygous for resistance gene, these plants are noted resistant)

S: mosaic symptoms.

*Proposed new wording:*

Ad 51: Resistance to Tomato mosaic virus (ToMV)

|   |  |
|---|--|
| 1. Pathogen.....                        | Tomato mosaic virus  |
| 3. Host species.....                    | <i>Solanum lycopersicum</i>  |
| 4. Source of inoculum .....             | Naktuinbouw <sup>11</sup> (NL) or GEVES <sup>12</sup> (FR)   |
| 5. Isolate.....                         | Strain 0 (e.g. isolate INRA Avignon 6-5-1-1) 1 and 2   |
| 6. Establishment isolate identity ..... | genetically defined tomato standards<br>Mobaci (Tm1), Moperou (Tm2), Momor (Tm2 <sup>2</sup> )<br>on susceptible plant   |
| 7. Establishment pathogenicity.....     |  |
| 8. Multiplication inoculum              |  |
| 8.1 Multiplication medium.....          | living plant   |
| 8.2 Multiplication variety .....        | e.g. Moneymaker, Marmande  |
| 8.7 Check of harvested inoculum .....   | option: on <i>Nicotiana tabacum</i> "Xanthi",<br>check lesions after 2 days<br>fresh>1 day, desiccated>1year   |
| 8.8 Shelf life/viability inoculum ..... |  |
| 9. Format of the test .....             |  |
| 9.1 Number of plants per genotype ..... | at least 20 plants   |
| 9.2 Number of replicates.....           | 1 replicate  |
| 9.3 Control varieties .....             |  |
| Susceptible.....                        | Marmande, Monalbo  |
| Resistant for ToMV: 0 and 2.....        | Mobaci   |
| Resistant for ToMV: 0 and 1 .....       | Moperou  |
| Resistant with necrosis .....           | "Monalbo x Momor"  |
| Resistant .....                         | Gourmet  |
| 9.4 Test design.....                    | blank treatment with PBS and carborundum or similar buffer   |
| 9.5 Test facility .....                 | Glasshouse or climate room   |
| 9.6 Temperature.....                    | 24 to 26°C   |
| 9.7 Light .....                         | 12 hours or longer   |
| 9.8 Season .....                        | symptoms are more pronounced in summer   |
| 10. Inoculation.....                    |  |
| 10.1 Preparation inoculum .....         | 1 g leaf with symptoms with 10 ml PBS or similar buffer<br>homogenize, add carborundum to buffer (1 g/30ml)  |
| 10.3 Plant stage at inoculation .....   | cotyledons or 2 leaves   |
| 10.4 Inoculation method .....           | gentle rubbing   |
| 10.7 Final observations .....           | 11-21 days after inoculation   |
| 11. Observations .....                  |  |
| 11.1 Method .....                       | visual   |
| 11.2 Observation scale .....            | Symptoms of susceptibility:<br>mosaic in top, leaf malformation<br>Symptoms of resistance (based on hypersensitivity):<br>local necrosis, top necrosis, systemic necrosis<br>evaluation of variety resistance should be calibrated with results of<br>resistant and susceptible controls |
| 11.3 Validation of test.....            |  |

Remark: in some heterozygous varieties a variable proportion of plants may have severe systemic necrosis or some necrotic spots while the other plants have no symptoms. This proportion may vary between experiments

12. Interpretation of test results in comparison with control varieties

|              |     |   |
|--------------|-----|---|
| absent.....  | [1] | symptoms of susceptibility                            |
| present..... | [9] | no symptoms, or symptoms of hypersensitive resistance |

13. Critical control points:

Temperature and light may influence the development of necrosis. More light means more necrosis. At temperatures above 26°C the resistance may break down.

Resistant heterozygous varieties may have symptomless plants and plants with severe necrosis; in spite of apparent segregation the sample may be evaluated as uniform for resistance

Note: Strain INRA Avignon 6-5-1-1 is recommended for ToMV: 0. This strain causes a striking yellow Aucuba mosaic.

<sup>11</sup> Naktuinbouw: resistentie@naktuinbouw.nl

<sup>12</sup> GEVES: Valerie.GRIMAULT@geves.fr

*Current wording:*

Ad. 52: Resistance to *Phytophthora infestans* (Pi)

Method

Maintenance of race

Type of medium: on agar medium

Special conditions: 18° C

Execution of test

Growth stage of plants: 10 leaves developed

Temperature: 18° C

Light: after inoculation darkness during 24 hours, thereafter 10 hours darkness per day

Growing method: climatic room or glasshouse

Method of inoculation: spraying of spore suspension, isolate harvested freshly from leaves

Duration of test

- from sowing to inoculation: 6 to 7 weeks

- from inoculation to reading: 7 to 8 days

Hygrometry: very high during the first four days after inoculation (cover plants with polyethylene cover)

Remarks: heterozygotes may show a lower level of expression of resistance

Standard varieties:

- susceptible: Saint Pierre, Heinz 1706
- resistant: Pieraline, Heline, Pyros, F1 "Pieraline x Pieralbo"

*Proposed new wording:*

Ad 52: Resistance to *Phytophthora infestans* (Pi)

|   |   |
|---|---|
| 1. Pathogen .....   | <i>Phytophthora infestans</i>   |
| 3. Host species .....   | <i>Solanum lycopersicum</i>   |
| 4. Source of inoculum .....   |   |
| 5. Isolate .....  | highly pathogenic on tomato   |
| 6. Establishment isolate identity .....                                 | biotest   |
| 7. Establishment pathogenicity .....                                    | biotest   |
| 8. Multiplication inoculum .....  |   |
| 8.1 Multiplication medium .....   | V8 Agar or PDA or Malt Agar medium  |
| 8.2 Multiplication variety .....  | susceptible tomato variety  |
| 8.3 Plant stage at inoculation .....                                    | 4 weeks   |
| 8.4 Inoculation medium .....  | water   |
| 8.5 Inoculation method .....  | spraying  |
| 8.6 Harvest of inoculum .....   | wash spores from wetted plates  |
| 8.7 Check of harvested inoculum .....                                   | count sporangiospores   |
| 8.8 Shelf life/viability inoculum .....                                 | 4 h after chilling at 8-10°C  |
| 9. Format of the test .....   |   |
| 9.1 Number of plants per genotype .....                                 | 20 plants   |
| 9.2 Number of replicates .....  | 1 replicate   |
| 9.3 Control varieties .....   |   |
| Susceptible .....   | Saint Pierre, Heinz 1706  |
| Resistant .....   | Pieraline, Heline, Pyros, "Pieraline x Pieralbo", Fline   |
| Remark: heterozygote varieties may have a                               | slightly lower level of expression of resistance.   |
| 9.5 Test facility .....   | glasshouse  |
| 9.6 Temperature .....   | 18°C  |
| 9.7 Light .....   | after inoculation darkness during 24 hours, thereafter 10 hour<br>darkness per 24 hours   |
| 9.9 Special measures .....  | humidity tent during four days after inoculation  |
| 10. Inoculation .....   |   |
| 10.1 Preparation inoculum .....   | wash spores from sporulating leaves, chill at 8-10°C<br>Chilling will induce zoospore release   |
| Remark .....  | Use fresh spores from repeated infection cycles on tomato<br>plants during 3 weeks before inoculation<br>count sporangiospores; adjust to 104 spores per ml<br>10 leaves developed (6 to 7 weeks) |
| 10.2 Quantification inoculum .....                                      |   |
| 10.3 Plant stage at inoculation .....                                   | 10 leaves developed (6 to 7 weeks)  |
| 10.4 Inoculation method .....   | spraying  |
| 10.7 Final observations .....   | 5-7 days after inoculation  |
| 11. Observations .....  |   |
| 11.1 Method .....   | visual  |
| 11.2 Observation scale .....  | Symptoms: water-soaked lesions, yellowing, and death  |
| 11.3 Validation of test .....   | evaluation of variety resistance should be calibrated with results<br>of resistant and susceptible controls   |
| 12. Interpretation of test results in comparison with control varieties |   |
| absent .....  | [1] severe symptoms   |
| present .....   | [9] no or mild symptoms   |
| 13. Critical control points:  |   |
| Resistance is only well-expressed in the adult plant.                   |   |



*Proposed new wording:*

Ad 53: Resistance to *Pyrenochaeta lycopersici* (PI)

|   |  |
|---|--|
| 1. Pathogen .....   | <i>Pyrenochaeta lycopersici</i>  |
| 3. Host species .....   | <i>Solanum lycopersicum</i>  |
| 4. Source of inoculum .....   | -  |
| 5. Isolate .....  | -  |
| 7. Establishment pathogenicity .....  | biotest  |
| 8. Multiplication inoculum .....  |  |
| 8.1 Multiplication medium .....   | V8 Agar  |
| 8.2 Multiplication variety .....  | susceptible tomato variety   |
| 8.3 Plant stage at inoculation .....  | seed   |
| 8.4 Inoculation medium.....   | mixture of soil, e.g. (70%), sand (20%) and inoculum (10.1) (10%)<br>or soil mixed with diseased roots cut to small pieces                         |
| 8.5 Inoculation method .....  | sowing, or transplanting at fruit maturity   |
| 8.6 Harvest of inoculum .....   | diseased roots are harvested after 2-4 months  |
| 8.7 Check of harvested inoculum .....   | visual inspection of lesions on roots  |
| 8.8 Shelf-life/viability inoculum .....   | the fungus will not die quickly, but may lose its pathogenicity within a week after isolation on an agar medium                                    |
| 9. Format of the test.....  |  |
| 9.1 Number of plants per genotype .....   | 20 plants  |
| 9.2 Number of replicates .....  | 1 replicate  |
| 9.3 Control varieties.....  |  |
| susceptible:.....   | Montfavet H 63.5   |
| resistant: .....  | Kyndia, Moboglan, Pyrella  |
| 9.5 Test facility .....   | greenhouse or climate cell   |
| 9.6 Temperature .....   | day 24°C, night 14°C   |
| 9.7 Light .....   | 12 h minimum   |
| 10. Inoculation .....   |  |
| 10.1 Preparation inoculum.....  | e.g. double-autoclaved mixture of soil with 10% oatmeal added<br>e.g. Incubate for 10-14 d at 20°C with occasional, repeated turning               |
| 10.3 Plant stage at inoculation .....   | 6 weeks  |
| 10.4 Inoculation method .....   | transplanting into mixture of soil, sand and inoculum (8.4)<br>or soil mixed with diseased roots cut to small pieces<br>or naturally infected soil |
| 10.7 Final observations .....   | 6-8 weeks after transplanting (flowering plant)  |
| 11. Observations.....   |  |
| 11.1 Method.....  | visual   |
| 11.2 Observation scale.....   | Symptoms: brown lesions on roots   |
| 11.3 Validation of test .....   | evaluation of variety resistance should be calibrated with results of resistant and susceptible controls   |
| 12. Interpretation of test results in comparison with control varieties   |  |
| absent .....  | [1] symptoms   |
| present.....  | [9] no symptoms  |
| 13. Critical control points:  |  |
| The fungus loses its pathogenicity quickly after isolation on an agar medium. It is advisable to keep the isolate alive on living plants. |  |

*Current wording:*

Ad. 54: Resistance to *Stemphylium*

Method

Maintenance of isolate

Type of medium: on PDA or synthetic medium

Special conditions: fridge 4° C without light

Execution of test

Growth stage of plants: three leaves expanded

Temperature: constant, day: 24° C, night: 24° C

Light: 12 hours

Growing method: glasshouse or climate room

Method of inoculation: pulverisation on leaves

Duration of test

- from sowing to inoculation: 20 to 22 days
- from inoculation to reading: 10 days

Number of plants tested: 30 plants

Remarks: production of inoculum on medium V8 under light

Standard varieties: susceptible: Monalbo  
resistant: Motelle, F1 Motelle x Monalbo

*Proposed new wording:*

Ad 54: Resistance to *Stemphylium* spp. (Ss)

|   |   |
|---|---|
| 1. Pathogen .....   | <i>Stemphylium</i> spp. e.g. <i>Stemphylium solani</i>  |
| 3. Host species .....   | <i>Solanum lycopersicum</i>   |
| 4. Source of inoculum .....   | GEVES <sup>13</sup> (FR)  |
| 5. Isolate .....  |   |
| 7. Establishment pathogenicity .....                                    | biotest   |
| 8. Multiplication inoculum .....  |   |
| 8.1 Multiplication medium .....   | PDA (12 hours per day under near-ultraviolet light to induce sporulation) or V8   |
| 9. Format of the test.....  |   |
| 9.1 Number of plants per genotype .....                                 | at least 20 plants  |
| 9.2 Number of replicates.....   | 1 replicate   |
| 9.3 Control varieties.....  |   |
| Susceptible: .....  | Monalbo   |
| Resistant: .....  | Motelle, F1 Motelle x Monalbo   |
| 9.5 Test facility .....   | greenhouse or climate cell  |
| 9.6 Temperature .....   | 24°C  |
| 9.7 Light .....   | 12 hours minimum  |
| 9.9 Special measures .....  | incubation in tunnel with 100 % relative humidity or humidity tent closed 5 days after inoculation, after this, 80% until end   |
| 10. Inoculation .....   |   |
| 10.1 Preparation inoculum.....  | sporulating plates (8.1) are scraped and air-dried overnight<br>The next day plates are soaked and stirred for 30 min in a beaker with demineralized water, or sporulating plates are scraped with water with Tween<br>The spore suspension is sieved through a double layer of muslin. |
| 10.2 Quantification inoculum .....                                      | 5.10 <sup>3</sup> – 10 <sup>5</sup> spores per ml   |
| 10.3 Plant stage at inoculation .....                                   | 20-22 days (three expanded leaves)  |
| 10.4 Inoculation method .....   | praying   |
| 10.7 Final observations .....   | 4 -10 days after inoculation  |
| 11. Observations.....   |   |
| 11.1 Method.....  | visual  |
| 11.2 Observation scale .....  | Symptoms:<br>necrotic lesions on cotyledons and leaves;<br>yellowing of leaves  |
| 11.3 Validation of test .....   | evaluation of variety resistance should be calibrated with results of resistant and susceptible controls  |
| 12. Interpretation of test results in comparison with control varieties |   |
| absent  | [1] symptoms (11.2)   |
| present   | [9] no symptoms, or less than resistant standard  |
| 13. Critical control points: .....                                      | 8.1 and 10.1  |

Note: Some isolates of *Stemphylium* cannot be classified easily as either *Stemphylium solani* or a related species. These *Stemphylium* isolates may still be useful for identifying resistance to *Stemphylium solani*.

<sup>13</sup> GEVES : Valerie.GRIMAULT@geves.fr

*Current wording:*

Ad. 55: Resistance to *Pseudomonas syringae* pv. tomato (Pst)

Method

Maintenance of races

Type of medium: on King B medium  
Special conditions: 20 - 22° C in the dark, transplantation every 10 days

Execution of test

Growth stage of plants: three leaves expanded  
Temperature: day: 22° C, night: 16° C  
Light: 12 hours  
Growing method: climatic room in summer, glasshouse in winter  
Method of inoculation: pulverisation on leaves

Duration of test

- from sowing to inoculation: 20 to 22 days  
- from inoculation to reading: 8 days

Number of plants tested: 30 plants  
Remarks: races to be renewed each year

Standard varieties: susceptible: Monalbo  
resistant: Ontario 7710, F1 Monalbo x Ontario 7710

*Proposed new wording:*

Ad 55: Resistance to *Pseudomonas syringae* pv. *tomato* (Pst)

|   |  |
|---|--|
| 1. Pathogen .....   | <i>Pseudomonas syringae</i> pv. <i>tomato</i>  |
| 3. Host species .....   | <i>Solanum lycopersicum</i>  |
| 4. Source of inoculum .....   | GEVES <sup>14</sup> (FR) or Naktuinbouw <sup>15</sup> (NL)   |
| 5. Isolate .....  |  |
| 6. Establishment isolate identity .....                                 |  |
| 7. Establishment pathogenicity .....                                    | biotest  |
| 8. Multiplication inoculum .....  |  |
| 8.1 Multiplication medium .....   | King's B agar medium, darkness   |
| 8.2 Multiplication variety .....  | Susceptible variety  |
| 8.4 Inoculation medium .....  | water  |
| 8.8 Shelf life/viability inoculum .....                                 | plates become old after 10 days  |
| 9. Format of the test .....   |  |
| 9.1 Number of plants per genotype .....                                 | at least 20 plants   |
| 9.2 Number of replicates .....  | 1 replicate  |
| 9.3 Control varieties .....   |  |
| Susceptible: .....  | Monalbo  |
| Resistant: .....  | Ontario 7710, "Monalbo x Ontario 7710", Tradiro, Hypeel 45   |
| 9.5 Test facility .....   | greenhouse or growth chamber   |
| 9.6 Temperature .....   | day: 22° C, night: 16° C or 20° C  |
| 9.7 Light .....   | 12 hours   |
| 9.9 Special measures .....  | humidity tent needed for 3 days or longer  |
| 10. Inoculation .....   |  |
| 10.1 Preparation inoculum .....   | wash off spores from plate. Plate should be less than 2-4 days old.                                      |
| 10.2 Quantification inoculum .....                                      | dilution plating, density 10 <sup>6</sup> colony forming units per ml                                    |
| 10.3 Plant stage at inoculation .....                                   | three leaves expanded (20-22 days)   |
| 10.4 Inoculation method .....   | spraying a bacterial suspension on leaves  |
| 10.7 Final observations .....   | 8 days after inoculation or longer   |
| 11. Observations .....  |  |
| 11.1 Method .....   | visual   |
| 11.2 Observation scale .....  | bacterial speck, greasy in appearance with marginal chlorosis<br>pinpoint lesions < 1.0 mm               |
| 11.3 Validation of test .....   | evaluation of variety resistance should be calibrated with results of resistant and susceptible controls |
| 12. Interpretation of test results in comparison with control varieties |  |
| absent .....  | [1] bacterial speck  |
| present .....   | [9] no symptoms or pinpoint lesions  |
| 13. Critical control points: .....                                      | Strains may lose virulence in storage  |

<sup>14</sup> GEVES; Valerie.GRIMAULT@geves.fr

<sup>15</sup> Naktuinbouw; resistentie@naktuinbouw.nl

*Current wording:*

Ad. 56: Resistance to *Ralstonia solanacearum* (Rs) – Race 1

Method

Maintenance of race : Two races may affect Tomato: race 1 (active between 25-30° C) and race 3 (active between 20-23° C)

Type of medium: Freezing at -80° C; culture in PYDAC immersed in oil; suspension in sterile distilled water

Special conditions: conservation at 15° C in sterile distilled water

Execution of test

Growth stage of plants: three to four well-developed leaves

Temp. (in climatic chamber): day: 26-30° C; night: 25° C

Light: 10 - 12 hours

Growing method: 2 possibilities:  
- in climatic chamber: rapid test  
- in the field: long test (applicable in tropical climate only)

Method of inoculation: deposit of at least 2 ml of inoculum, adjusted to  $10^7$  colonies per ml, at the foot of each plantlet prior to planting

Duration of test

- from sowing to inoculation: 3 to 4 weeks  
- from inoculation to reading:  
- 3 weeks for the fast test  
- 2 months for the long test

Number of plants tested: minimum of 30

Remarks: maintain high humidity

Standard varieties:  
- susceptible: Floradel  
- resistant: Caraïbo

*Proposed new wording:*

Ad 56: Resistance to *Ralstonia solanacearum*, race 1 (Rs)

|  |   |
|--|---|
| 1. Pathogen .....  | <i>Ralstonia solanacearum</i> (ex <i>Pseudomonas solanacearum</i> )   |
| 2. Quarantine status .....   | yes   |
| 3. Host species .....  | <i>Solanum lycopersicum</i>   |
| 4. Source of inoculum .....  |   |
| 5. Isolate .....   | Race 1 has a wide host range, including tomato.<br>Race 3 has a narrow host range, also including tomato    |
| 8. Multiplication inoculum .....   |   |
| 8.1 Multiplication medium .....  | Yeast Peptone Glucose (YPG) Agar or PYDAC   |
| Special conditions:.....   | 25-30°C (Race 3 usually needs 20-23°C)  |
| 8.5 Inoculation method .....   | 2 ml of inoculum placed at the foot of each plantlet<br>prior to transplanting                              |
| 8.8 Shelf life/viability inoculum.....   | suspension in sterile distilled water at 15°C (<1 year)   |
| 9. Format of the test.....   |   |
| 9.1 Number of plants per genotype .....  | 20 plants   |
| 9.2 Number of replicates.....  | 1 replicate   |
| 9.3 Control varieties.....   |   |
| Susceptible: .....   | Floradel  |
| Resistant: .....   | Caraibo   |
| 9.5 Test facility .....  | climate room  |
| 9.6 Temperature .....  | day: 26-30° C; night: 25° C   |
| 9.7 Light .....  | 10 - 12 hours   |
| 9.9 Special measures .....   | high humidity   |
| 10. Inoculation .....  |   |
| 10.2 Quantification inoculum .....   | density 10 <sup>7</sup> colony forming units per ml   |
| 10.3 Plant stage at inoculation .....  | three to four well-developed leaves (3 weeks)   |
| 10.4 Inoculation method .....  |   |
| 10.7 Final observations .....  | 3 weeks after inoculation   |
| 11. Observations.....  | In intermediate resistance varieties, bacteria could be present in<br>the lower part of the plant           |
| 11.3 Validation of test .....  | evaluation of variety resistance should be calibrated with results<br>of resistant and susceptible controls |
| 12. Interpretation of test results in comparison with control varieties                                |   |
| absent .....   | [1] symptoms  |
| present.....   | [9] no symptoms, or less than resistant standard  |
| 13. Critical control points:   |   |
| <i>Ralstonia solanacearum</i> has a quarantine status in some countries and is on the EPPO alert list. |   |

*Current wording:*

Ad. 57: Resistance to Tomato Yellow Leaf Curl Begomovirus (TYLCV)

Method

Execution of test: Plants are tested under field crop conditions respecting a period of planting and a place where the disease has been proven to exist. 100% contaminated plants are grown of susceptible local varieties to ensure natural transmission by *Bemisia* insect and repeatability of the results

Growth stage of plants: on adult plants of field crop outside

Method of inoculation: natural inoculation by *Bemisia*

Duration of test

- from sowing to inoculation: 6 weeks minimum
- from inoculation to reading: 2.5 months maximum

Number of plants tested: 20 plants minimum

Remarks:

Standard varieties: - susceptible: local varieties  
- resistant: TY 20 or accessions from *L. pimpinellifolium* and from *L. peruvianum*

*Proposed new wording:*

Ad 57: Resistance to Tomato yellow leaf curl virus (TYLCV)

|  |  |
|--|--|
| 1. Pathogen .....  | Tomato yellow leaf curl virus  |
| 2. Quarantine status .....   | yes  |
| 3. Host species .....  | <i>Solanum lycopersicum</i>  |
| 4. Source of inoculum .....  | -  |
| 5. Isolate .....   | -  |
| 8. Multiplication inoculum .....   |  |
| 8.6 Harvest of inoculum .....  | symptomatic leaves may be stored at -70°C  |
| 9. Format of the test.....   |  |
| 9.1 Number of plants per genotype .....  | 20 plants  |
| 9.2 Number of replicates.....  | 1 replicate  |
| 9.3 Control varieties.....   |  |
| Susceptible: .....   | Montfavet H 63.5   |
| Resistant: .....   | TY 20, Anastasia, Mohawk   |
| 9.5 Test facility .....  | field with natural disease pressure  |
| 9.9 Special measures .....   | prevent spread of white-flies  |
| 10. Inoculation .....  |  |
| 10.3 Plant stage at inoculation .....  | 6-12 weeks (adult plants)  |
| 10.4 Inoculation method .....  | vector (Bemisia white-flies carrying TYLCV)  |
| 10.7 Final observations .....  | 1-2 months after inoculation   |
| 11. Observations.....  |  |
| 11.1 Method.....   | visual   |
| 11.2 Observation scale.....  | Symptoms: leaf yellowing and curling   |
| 11.3 Validation of test .....  | evaluation of variety resistance should be calibrated with results of resistant and susceptible controls |
| 12. Interpretation of test results in comparison with control varieties  |  |
| absent .....   | [1] severe symptoms  |
| present.....   | [9] no or mild symptoms  |
| 13. Critical control points:   |  |
| TYLCV is endemic in many tropical and subtropical areas and has a quarantine status in many countries with a temperate climate. TYLCV is on the EPPO alert list. Some TYLCV resistant varieties may be susceptible to the closely related virus Tomato yellow leaf curl Sardinia virus (TYLCSV). |  |

*Current wording:*

Ad. 58: Resistance to Tomato Spotted Wilt Tospovirus (TSWV) - Race 0

Method

Maintenance of races

Type of medium: on tomato plants or frozen at  $-70^{\circ}\text{C}$

Special conditions:

Execution of test

Growth stage of plants: one or two leaves expanded

Temperature: day:  $20^{\circ}\text{C}$ , night:  $20^{\circ}\text{C}$

Light: extra light in winter

Growing method: glasshouse

Method of inoculation: mechanical, rubbing with carborundum on cotyledons, inoculum suspension  $< 10^{\circ}\text{C}$

Duration of test

- from sowing to inoculation: 20 days
- from inoculation to reading: 14 to 20 days

Number of plants tested: 15 to 30 plants

Remarks: be aware of thrips

Standard varieties:  
- susceptible: Monalbo  
- resistant: Tsunami, Bodar, Lisboa

*Proposed new wording:*

Ad 58: Resistance to Tomato spotted wilt virus (TSWV)

|   |   |
|---|---|
| 1. Pathogen .....   | Tomato spotted wilt virus   |
| 2. Quarantine status .....  | yes (see note below)  |
| 3. Host species .....   | <i>Solanum lycopersicum</i>   |
| 4. Source of inoculum .....   | Naktuinbouw <sup>16</sup> (NL), GEVES (FR)  |
| 5. Isolate .....  | race 0, preferably a thrips-transmission deficient variant  |
| 7. Establishment pathogenicity .....  | biotest   |
| 8. Multiplication inoculum .....  |   |
| 6 Harvest of inoculum .....   | symptomatic leaves may be stored at -70°C   |
| 9. Format of the test.....  |   |
| 9.1 Number of plants per genotype .....   | 20 plants   |
| 9.2 Number of replicates.....   | 1 replicate   |
| 9.3 Control varieties .....   |   |
| Susceptible: .....  | Monalbo, Momor, Montfavet H 63.5  |
| Resistant: .....  | Tsunami, Bodar, Mospomor, Lisboa  |
| 9.5 Test facility .....   | glasshouse or climatic chamber  |
| 9.6 Temperature .....   | 20°C  |
| 9.7 Light .....   | 12 hours or longer  |
| 9.9 Special measures .....  | prevent or combat thrips  |
| 10. Inoculation .....   |   |
| 10.1 Preparation inoculum.....  | press symptomatic leaves in ice-cold buffer<br>0,01 M PBS, pH 7.4, with 0,01 M sodium sulfite or similar buffer<br>Option: sieve the leaf sap through double muslin |
| 10.3 Plant stage at inoculation .....   | one or two expanded leaves  |
| 10.4 Inoculation method .....   | mechanical, rubbing with carborundum on cotyledons, inoculum<br>suspension < 10° C  |
| 10.7 Final observations .....   | 7-21 days after inoculation   |
| 11. Observations.....   |   |
| 11.1 Method.....  | visual  |
| 11.2 Observation scale.....   | Symptoms: top mosaic, bronzing, various malformations,<br>necrosis  |
| 11.3 Validation of test.....  | evaluation of variety resistance should be calibrated with results<br>of resistant and susceptible controls   |
| 12. Interpretation of test results in comparison with control varieties   |   |
| absent.....[1]  | symptoms  |
| present.....[9]   | no symptoms   |
| 13. Critical control points:  |   |
| TSWV has a quarantine status in some countries. TSWV is transmitted by <i>Thrips tabaci</i> and Western flower thrips ( <i>Frankliniella occidentalis</i> ). Pathotype 0 is defined by its inability to break resistance in tomato varieties carrying the resistance gene Sw-5. |   |

<sup>16</sup> Naktuinbouw: [resistentie@naktuinbouw.nl](mailto:resistentie@naktuinbouw.nl)

GEVES; Valerie.GRIMAULT@geves.fr

*Current wording:*

Ad. 59: Resistance to *Leveillula taurica* (Lt)

Method

Maintenance of races

Type of medium: tomato plants  
Special conditions:

Execution of test

Growth stage of plants: on adult plants of field crop outside

Method of inoculation: natural infection

Duration of test

- from sowing to inoculation: infection possible from planting stage to full grown plants  
- from inoculation to reading: before harvest

Number of plants tested: 20 plants

Remarks: Yellow chlorotic spots on upper side of leaves, mycelium on lower side of leaves.  
Check cleistothecia under microscope if it really concerns *Leveillula* and not another powdery mildew.

Standard varieties: - susceptible: Monalbo  
- resistant: Atlanta

*Proposed new wording:*

Ad 59: Resistance to *Leveillula taurica* (Lt)

|   |   |
|---|---|
| 1. Pathogen .....   | <i>Leveillula taurica</i>   |
| 3. Host species .....   | <i>Solanum lycopersicum</i>   |
| 4. Source of inoculum .....   | no long term storage method is available  |
| 5. Isolate .....  |   |
| 8.1 Multiplication medium .....   | detached leaves of a susceptible host plant   |
| 9. Format of the test.....  |   |
| 9.1 Number of plants per genotype .....   | 20 plants   |
| 9.2 Number of replicates.....   | 1 replicate   |
| 9.3 Control varieties.....  |   |
| Susceptible: .....  | Monalbo , Montfavet H 63.5  |
| Resistant:.....   | Atlanta   |
| 10. Inoculation .....   |   |
| 10.3 Plant stage at inoculation .....   | adult plants  |
| 10.4 Inoculation method .....   | natural infection, mainly by wind dispersal of spores   |
| 10.7 Final observations .....   | before harvest  |
| 11. Observations.....   |   |
| 11.1 Method.....  | visual  |
| 11.2 Observation scale .....  | Symptoms: Yellow chlorotic spots on upper side of leaves,<br>mycelium on abaxial side of leaves             |
| 11.3 Validation of test .....   | evaluation of variety resistance should be calibrated with results<br>of resistant and susceptible controls |
| 12. Interpretation of test results in comparison with control varieties                                       |   |
| absent .....  | [1] symptoms  |
| present.....  | [9] no symptoms, or less than resistant standard  |
| 13. Critical control points:  |   |
| Check cleistothecia under microscope to confirm presence of <i>Leveillula</i> and not another powdery mildew. |   |

*Current wording:*

Ad. 60: Resistance to *Oidium neolycopersici* (On) (ex *Oidium lycopersicum* (Ol))

Method

Maintenance of races

Type of medium: on tomato plants

Special conditions: climatic room

Execution of test

Growth stage of plants: 3 weeks

Temperature: 24°C during the day; 18°C during the night

Light: 12 hours

Method of inoculation: - by spraying ( $10^4$  conidia/ml) on leaves  
- by dredging (uncontrolled inoculum) on leaves

Execution of test

Duration of test

- from sowing to inoculation: 18 - 20 days

- from inoculation to reading: 15 – 18 days

Number of plants tested: 30 plants/lot

Remarks:

Scale of notes: - no sporulation }  
- sporulation without extension }Resistant  
(necrotic points) }  
- moderate sporulation }  
- abundant sporulation }Susceptible

Standard varieties: - susceptible: Momor (*L. esculentum*)  
- resistant: *L. hirsutum* PI-247087 (accession), Romiror

*Proposed new wording:*

Ad 60: Resistance to *Oidium neolycopersici* (On)

|   |  |
|---|--|
| 1. Pathogen .....   | <i>Oidium neolycopersici</i> (Powdery mildew)  |
| 3. Host species .....   | <i>Solanum lycopersicum</i>  |
| 4. Source of inoculum .....   |  |
| 5. Isolate see remark under 13.....                                     |  |
| 7. Establishment pathogenicity .....                                    | biotest  |
| 8. Multiplication inoculum .....  |  |
| 8.1 Multiplication medium .....   | plant  |
| 8.3 Plant stage at inoculation .....                                    | 3 weeks  |
| 8.4 Inoculation medium.....   | water  |
| 8.5 Inoculation method .....  | see 10.4   |
| 8.6 Harvest of inoculum.....  | by washing off   |
| 8.7 Check of harvested inoculum .....                                   | check for contaminants under microscope  |
| 8.8 Shelf-life/viability inoculum .....                                 | 1-2 hours  |
| 9. Format of the test.....  |  |
| 9.1 Number of plants per genotype .....                                 | 20 plants  |
| 9.2 Number of replicates.....   | 1 replicate  |
| 9.3 Control varieties.....  |  |
| Susceptible: .....  | Momor, Montfavet H 63.5  |
| Resistant tomato: .....   | Atlanta, Romiro, PI-247087   |
| 9.5 Test facility .....   | glasshouse   |
| 9.6 Temperature .....   | 20°C or 18/24°C  |
| 9.7 Light .....   | 12 hours   |
| 10. Inoculation .....   |  |
| 10.1 Preparation inoculum.....  | collect spores in water  |
| 10.2 Quantification inoculum .....                                      | 10 <sup>4</sup> conidia/ml   |
| 10.3 Plant stage at inoculation .....                                   | 3 weeks  |
| 10.4 Inoculation method .....   | by spraying on leaves or dredging of leaves  |
| 10.7 Final observations .....   | 7-18 days after inoculation  |
| 11. Observations.....   |  |
| 11.1 Method.....  | visual   |
| 11.2 Observation scale .....  | 0. no sporulation<br>1. necrotic points and sometimes locally restricted sporulation<br>2. moderate sporulation<br>3. abundant sporulation |
| 11.3 Validation of test .....   | evaluation of variety resistance should be calibrated with results of resistant and susceptible controls                                   |
| 12. Interpretation of test results in comparison with control varieties |  |
| absent.....   | [1] moderate or abundant sporulation   |
| present.....  | [9] no or restricted sporulation   |

13. Critical control points:

Resistance-breaking isolates should be avoided. Resistance to *O. neolycopersici* is usually race-specific. However, as long as a differential series of tomato genotypes with well defined resistances is lacking, it will remain hard to conclude that different races of *O. neolycopersici* exist.

*Current wording:*

Ad. 61: Resistance to Tomato Torrado Virus (ToTV)

Method

Maintenance of races

Type of medium: plant material with symptoms, stored at –80° C

Multiplication: on *N. tabacum* 'Xanthi' 3 weeks before start of experiment

Special conditions: use Quarantine procedures

Remarks: white fly may be a vector of ToTV

Execution of test

Growth stage of plants: inoculate when cotyledons are fully grown, re-inoculate 7 days later on first true leaves one or two leaves

Temperature: day: 23° C, night: 21° C; avoid temperature above 25°C

Light: extra light in winter, 16 h day, 8 h night

Growing method: Quarantine facilities; glasshouse

Method of inoculation: with ice-cold 0,01 M PBS pH 7 and carborundum

Duration of test

- from sowing to inoculation: 14 days

- from inoculation to reading: 14-21 days

Number of plants tested: 20 to 30 plants

Remarks: Necrotic spots on the top leaves of susceptible plants

Standard varieties: Resistant standard variety: Matias

Note: Patents pending on part of the method: WO2006/085749 and WO2008/150158 and equivalents. Use solely for DUS purposes and for the development of variety descriptions by UPOV and authorities of UPOV members, courtesy of De Ruiters Seeds R&D B.V./Monsanto Invest N.V.

*Proposed new wording:*

Ad 61: Resistance to Tomato torrado virus (ToTV)

|   |  |
|---|--|
| 1. Pathogen .....   | Tomato Torrado Virus   |
| 2. Quarantine status .....  | in regions with temperate climate  |
| 3. Host species .....   | <i>Solanum lycopersicum</i>  |
| 4. Source of inoculum .....   | -  |
| 5. Isolate .....  | -  |
| 7. Establishment pathogenicity .....  | biotest  |
| 8. Multiplication inoculum  |  |
| 8.1 Multiplication medium .....   | Nicotiana tabacum 'Xanthi'   |
| 8.3 Plant stage at inoculation .....  | cotyledon to first leaf  |
| 8.5 Inoculation method .....  | see 10.4   |
| 8.6 Harvest of inoculum .....   | after 3 weeks  |
| 8.7 Check of harvested inoculum .....   | plants yellow, systemic infection  |
| 8.8 Shelf-life/viability inoculum .....   | instable at room temperature   |
| 9. Format of the test.....  |  |
| 9.1 Number of plants per genotype .....   | 20 plants  |
| 9.2 Number of replicates.....   | 1 replicate  |
| 9.3 Control varieties.....  |  |
| Susceptible: .....  | Daniela  |
| Resistant tomato: .....   | Matias   |
| 9.5 Test facility .....   | glasshouse   |
| 9.6 Temperature .....   | 23°C during the day; 21°C during the night   |
| 9.7 Light .....   | 16 hours   |
| 10. Inoculation .....   |  |
| 10.3 Plant stage at inoculation .....   | 14 days  |
| 10.4 Inoculation method .....   | with ice-cold 0,01 M PBS pH 7 and carborundum  |
| 10.5 First observation .....  | 7 days after inoculation   |
| 10.6 Second observation .....   | 14 days after inoculation  |
| 10.7 Final observations .....   | 18 days after inoculation  |
| 11. Observations.....   |  |
| 11.1 Method.....  | visual   |
| 11.2 Observation scale .....  | necrotic spots on the top leaves   |
| 11.3 Validation of test .....   | evaluation of variety resistance should be calibrated with results of resistant and susceptible controls |
| 12. Interpretation of test results in comparison with control varieties                                       |  |
| absent.....[1]  | necrotic spots present   |
| present.....[9]   | no symptoms  |
| 13. Critical control points:  |  |
| ToTV is transmitted by white fly ( <i>Bemisia tabaci</i> ). Produce inoculum with ice-cold mortar and pestle. |  |
| During inoculation the temperature should be below 25°C   |  |

Note: Patents pending on part of the method: WO2006/085749 and WO2008/150158 and equivalents. Use solely for DUS purposes and for the development of variety descriptions by UPOV and authorities of UPOV members, courtesy to De Ruiter Seeds R&D B.V./Monsanto Invest N.V.

[Annex III follows]

Proposal to Add the Following Literature References to Chapter 9: Literature

Arens P., Mansilla C., Deinum D., Cavellini L., Moretti A., Rolland S., van der Schoot H., Calvache D., Ponz F., Collonnier C., Mathis R., Smilde D., Caranta C., Vosman B., 2010. Development and evaluation of robust molecular markers linked to disease resistance in tomato for distinctness, uniformity and stability testing. *Theoretical and applied genetics*. 120(3): 655-64

Bai, Y. 2004. The genetics and mechanisms of resistance to tomato powdery mildew (*Oidium neolycopersici*) in *Lycopersicon* species. Thesis Wageningen University, The Netherlands.

Barbieri, M., et al., 2010. Introgressions of resistance to two Mediterranean virus species causing tomato yellow leaf curl into a valuable traditional tomato variety. *Journal of Plant Pathology* 92(2):485-493

Garcia, S., et al., 2009. Resistance driven selection of begomoviruses associated with the TYLCV. *Virus research* 146: 66-72

Garland, S., Sharman, M., Persley, D. and McGrath, D. (2005) The development of an improved PCR-based marker system for Sw-5, an important TSWV resistance gene of tomato. *Australian Journal of Agricultural Research*, 56 (3): 285-289.

Gordillo, L.F. and M. R. Stevens (2008) Screening two *Lycopersicon peruvianum* collections for resistance to Tomato spotted wilt virus. *Plant Disease* 92(5): 694-704

Hubbeling, N., 1978. Breakdown of resistance to the Cf-5 gene in tomato by another new race of *Fulvia fulva*. *Mededelingen van de Faculteit Landbouwwetenschappen Universiteit Gent* 42/2

Martin, G. B., A. Frary, T. Wu, S. Brommonschenkel, J. Chunwongse, E. D. Earle, S. D. Tanksley (1994) A member of the tomato Pto family confers sensitivity to fenthion resulting in rapid cell death. *The Plant Cell* 6: 1543-1552

[http://www.worldseed.org/isf/pathogen\\_coding\\_3.html](http://www.worldseed.org/isf/pathogen_coding_3.html) (International Seed Federation (ISF), Trade Issues, Phytosanitary Matters, Pathogen coding, Strain Denomination, Differential sets)

[End of Annex III and of document]