



TG/125/7(proj.5)  
ORIGINAL: English  
DATE: 2017-02-15

## INTERNATIONAL UNION FOR THE PROTECTION OF NEW VARIETIES OF PLANTS

Geneva

DRAFT

### WALNUT

UPOV Code(s):

JUGLA\_REG

*Juglans regia L.*

### GUIDELINES

#### FOR THE CONDUCT OF TESTS

#### FOR DISTINCTNESS, UNIFORMITY AND STABILITY

*prepared by experts from China  
to be considered by the  
Technical Committee  
at its fifty-third session, to be held in Geneva  
from 2017-04-03 to 2017-04-05*

*Disclaimer: this document does not represent UPOV policies or guidance*

Alternative names:<sup>\*</sup>

Botanical name	English	French	German	Spanish
<i>Juglans regia L.</i>	Walnut, English Walnut	Noyer	Walnuß	Nogal

The purpose of these guidelines (“Test Guidelines”) is to elaborate the principles contained in the General Introduction (document TG/1/3), and its associated TGP documents, into detailed practical guidance for the harmonized examination of distinctness, uniformity and stability (DUS) and, in particular, to identify appropriate characteristics for the examination of DUS and production of harmonized variety descriptions.

#### ASSOCIATED DOCUMENTS

These Test Guidelines should be read in conjunction with the General Introduction and its associated TGP documents.

\* These names were correct at the time of the introduction of these Test Guidelines but may be revised or updated. [Readers are advised to consult the UPOV Code, which can be found on the UPOV Website ([www.upov.int](http://www.upov.int)), for the latest information.]

<u>TABLE OF CONTENTS</u>	<u>PAGE</u>
1. SUBJECT OF THESE TEST GUIDELINES.....	<u>4</u>
2. MATERIAL REQUIRED.....	<u>4</u>
3. METHOD OF EXAMINATION.....	<u>5</u>
3.1    Number of Growing Cycles.....	<u>5</u>
3.2    Testing Place.....	<u>5</u>
3.3    Conditions for Conducting the Examination.....	<u>5</u>
3.4    Test Design.....	<u>5</u>
3.5    Additional Tests.....	<u>5</u>
4. ASSESSMENT OF DISTINCTNESS, UNIFORMITY AND STABILITY.....	<u>6</u>
4.1    Distinctness.....	<u>6</u>
4.2    Uniformity.....	<u>7</u>
4.3    Stability.....	<u>7</u>
5. GROUPING OF VARIETIES AND ORGANIZATION OF THE GROWING TRIAL.....	<u>8</u>
6. INTRODUCTION TO THE TABLE OF CHARACTERISTICS.....	<u>9</u>
6.1    Categories of Characteristics.....	<u>9</u>
6.2    States of Expression and Corresponding Notes.....	<u>9</u>
6.3    Types of Expression.....	<u>9</u>
6.4    Example Varieties.....	<u>10</u>
6.5    Legend.....	<u>11</u>
7. TABLE OF CHARACTERISTICS/TABLEAU DES CARACTÈRES/MERKMALSTABELLE/TABLA DE CARACTERES.....	<u>12</u>
8. EXPLANATIONS ON THE TABLE OF CHARACTERISTICS.....	<u>23</u>
8.1    Explanations covering several characteristics.....	<u>23</u>
8.2    Explanations for individual characteristics.....	<u>23</u>
9. LITERATURE.....	<u>23</u>
10. TECHNICAL QUESTIONNAIRE.....	<u>34</u>

1. Subject of these Test Guidelines

These Test Guidelines apply to all varieties of *Juglans regia* L. for fruit use.

2. Material Required

- 2.1 The competent authorities decide on the quantity and quality of the plant material required for testing the variety and when and where it is to be delivered. Applicants submitting material from a State other than that in which the testing takes place must ensure that all customs formalities and phytosanitary requirements are complied with.
- 2.2 The material is to be supplied in the form of dormant shoots sufficient for grafting 5 plants or in the form of grafted plants on a rootstock specified by the testing authority.
- 2.3 The minimum quantity of plant material, to be supplied by the applicant, should be:  

10 dormant shoots for grafting or 5 one-year-old grafted plants.
- 2.4 The plant material supplied should be visibly healthy, not lacking in vigor, nor affected by any important pest or disease.
- 2.5 The plant material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If it has been treated, full details of the treatment must be given.

3. Method of Examination

3.1 *Number of Growing Cycles*

- 3.1.1 The minimum duration of tests should normally be two independent growing cycles.
- 3.1.2 The growing cycle is considered to be the duration of a single growing season, beginning with bud burst (flowering and/or vegetative), flowering and fruit harvest and concluding when the following dormant period ends with the swelling of new season buds.

3.2 *Testing Place*

Tests are normally conducted at one place. In the case of tests conducted at more than one place, guidance is provided in TGP/9 "Examining Distinctness".

3.3 *Conditions for Conducting the Examination*

The tests should be carried out under conditions ensuring satisfactory growth for the expression of the relevant characteristics of the variety and for the conduct of the examination.

3.4 *Test Design*

- 3.4.1 Each test should be designed to result in a total of at least 5 plants.
- 3.4.2 The design of the tests should be such that plants or parts of plants may be removed for measurement or counting without prejudice to the observations which must be made up to the end of the growing cycle.

3.5 *Additional Tests*

Additional tests, for examining relevant characteristics, may be established.

#### 4. Assessment of Distinctness, Uniformity and Stability

##### 4.1 *Distinctness*

###### 4.1.1 General Recommendations

It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding distinctness. However, the following points are provided for elaboration or emphasis in these Test Guidelines.

###### 4.1.2 Consistent Differences

The differences observed between varieties may be so clear that more than one growing cycle is not necessary. In addition, in some circumstances, the influence of the environment is not such that more than a single growing cycle is required to provide assurance that the differences observed between varieties are sufficiently consistent. One means of ensuring that a difference in a characteristic, observed in a growing trial, is sufficiently consistent is to examine the characteristic in at least two independent growing cycles.

###### 4.1.3 Clear Differences

Determining whether a difference between two varieties is clear depends on many factors, and should consider, in particular, the type of expression of the characteristic being examined, i.e. whether it is expressed in a qualitative, quantitative, or pseudo-qualitative manner. Therefore, it is important that users of these Test Guidelines are familiar with the recommendations contained in the General Introduction prior to making decisions regarding distinctness.

###### 4.1.4 Number of plants or parts of plants to be Examined

Unless otherwise indicated, for the purposes of distinctness, all observations on single plants should be made on 5 plants or parts of plants taken from each of 5 plants and any other observations made on all plants in the test, disregarding any off-type plants.

In the case of observations of parts taken from single plants, the number of parts to be taken from each of the plants should be 2.

###### 4.1.5 Method of Observation

The recommended method of observing the characteristic for the purposes of distinctness is indicated by the following key in the second column of the Table of Characteristics (see document TGP/9 "Examining Distinctness", Section 4 "Observation of characteristics"):

MG: single measurement of a group of plants or parts of plants

MS: measurement of a number of individual plants or parts of plants

VG: visual assessment by a single observation of a group of plants or parts of plants

VS: visual assessment by observation of individual plants or parts of plants

Type of observation: visual (V) or measurement (M)

"Visual" observation (V) is an observation made on the basis of the expert's judgment. For the purposes of this document, "visual" observation refers to the sensory observations of the experts and, therefore, also includes smell, taste and touch. Visual observation includes observations where the expert uses reference points (e.g. diagrams, example varieties, side-by-side comparison) or non-linear charts (e.g. color charts). Measurement (M) is an objective observation against a calibrated, linear scale e.g. using a ruler, weighing scales, colorimeter, dates, counts, etc.

Type of record: for a group of plants (G) or for single, individual plants (S)

For the purposes of distinctness, observations may be recorded as a single record for a group of plants or parts of plants (G), or may be recorded as records for a number of single, individual plants or parts of plants (S). In most cases, "G" provides a single record per variety and it is not possible or necessary to apply statistical methods in a plant-by-plant analysis for the assessment of distinctness.

In cases where more than one method of observing the characteristic is indicated in the Table of Characteristics (e.g. VG/MG), guidance on selecting an appropriate method is provided in document TGP/9, Section 4.2.

#### 4.2 *Uniformity*

- 4.2.1 It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding uniformity. However, the following points are provided for elaboration or emphasis in these Test Guidelines:
- 4.2.2 The assessment of uniformity should be according to the recommendations for cross-pollinated varieties in the General Introduction.
- 4.2.3 For the assessment of uniformity of vegetatively propagated varieties, a population standard of 1% and an acceptance probability of at least 95% should be applied. In the case of a sample size of 5 plants, no off-types are allowed.

#### 4.3 *Stability*

- 4.3.1 In practice, it is not usual to perform tests of stability that produce results as certain as those of the testing of distinctness and uniformity. However, experience has demonstrated that, for many types of variety, when a variety has been shown to be uniform, it can also be considered to be stable.
- 4.3.2 Where appropriate, or in cases of doubt, stability may be further examined by testing a new plant stock to ensure that it exhibits the same characteristics as those shown by the initial material supplied.

5. Grouping of Varieties and Organization of the Growing Trial

- 5.1 The selection of varieties of common knowledge to be grown in the trial with the candidate varieties and the way in which these varieties are divided into groups to facilitate the assessment of distinctness are aided by the use of grouping characteristics.
- 5.2 Grouping characteristics are those in which the documented states of expression, even where produced at different locations, can be used, either individually or in combination with other such characteristics: (a) to select varieties of common knowledge that can be excluded from the growing trial used for examination of distinctness; and (b) to organize the growing trial so that similar varieties are grouped together.
- 5.3 The following have been agreed as useful grouping characteristics:
- (a) Female flower: number per cluster (characteristic 6)
  - (b) Female flower: intensity of yellow color of stigma (characteristic 7)
  - (c) Fruit: setting type (characteristic 8)
  - (d) Nut: shape in lateral view (characteristic 10)
  - (e) Nut: thickness of shell (characteristic 24)
  - (f) Kernel: color of endopileura (characteristic 25)
  - (g) Time of male flowering compared to female flowering (characteristic 30)
- 5.4 Guidance for the use of grouping characteristics, in the process of examining distinctness, is provided through the General Introduction and document TGP/9 "Examining Distinctness".

6. Introduction to the Table of Characteristics

6.1 *Categories of Characteristics*

6.1.1 Standard Test Guidelines Characteristics

Standard Test Guidelines characteristics are those which are approved by UPOV for examination of DUS and from which members of the Union can select those suitable for their particular circumstances.

6.1.2 Asterisked Characteristics

Asterisked characteristics (denoted by \*) are those included in the Test Guidelines which are important for the international harmonization of variety descriptions and should always be examined for DUS and included in the variety description by all members of the Union, except when the state of expression of a preceding characteristic or regional environmental conditions render this inappropriate.

6.2 *States of Expression and Corresponding Notes*

6.2.1 States of expression are given for each characteristic to define the characteristic and to harmonize descriptions. Each state of expression is allocated a corresponding numerical note for ease of recording of data and for the production and exchange of the description.

6.2.2 In the case of qualitative and pseudo-qualitative characteristics (see Chapter 6.3), all relevant states of expression are presented in the characteristic. However, in the case of quantitative characteristics with 5 or more states, an abbreviated scale may be used to minimize the size of the Table of Characteristics. For example, in the case of a quantitative characteristic with 9 states, the presentation of states of expression in the Test Guidelines may be abbreviated as follows:

State	Note
small	3
medium	5
large	7

However, it should be noted that all of the following 9 states of expression exist to describe varieties and should be used as appropriate:

State	Note
very small	1
very small to small	2
small	3
small to medium	4
medium	5
medium to large	6
large	7
large to very large	8
very large	9

6.2.3 Further explanation of the presentation of states of expression and notes is provided in document TGP/7 “Development of Test Guidelines”.

6.3 *Types of Expression*

An explanation of the types of expression of characteristics (qualitative, quantitative and pseudo-qualitative) is provided in the General Introduction.

6.4 *Example Varieties*

Where appropriate, example varieties are provided to clarify the states of expression of each characteristic.

## 6.5 Legend

	English		français		deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
1	2	3	4	5	6	7		
	Name of characteristics in English	Nom du caractère en français		Name des Merkmals auf Deutsch	Nombre del carácter en español			
	states of expression	types d'expression		Ausprägungsstufen	tipos de expresión			

- 1 Characteristic number
- 2 (\*) Asterisked characteristic – see Chapter 6.1.2
- 3 Type of expression
  - QL Qualitative characteristic – see Chapter 6.3
  - QN Quantitative characteristic – see Chapter 6.3
  - PQ Pseudo-qualitative characteristic – see Chapter 6.3
- 4 Method of observation (and type of plot, if applicable)
  - MG, MS, VG, VS – see Chapter 4.1.5
- 5 (+) See Explanations on the Table of Characteristics in Chapter 8.2
- 6 (a)-(e) See Explanations on the Table of Characteristics in Chapter 8.1
- 7 Not applicable

7. Table of Characteristics/Tableau des caractères/Merkmalstabelle/Tabla de caracteres

	English		français		deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplos	Note/ Nota
1. (*)	QN	VG	(+)	(a)				
	Tree: growth habit		Arbre : port		Baum: Wuchsform	Árbol: hábito de crecimiento		
	upright		dressé		aufrecht	erecto	Corne, Sorrento, Daifeng, Zhonglin 1, Fenghui, Xinzaofeng, Daixiang	1
	semi-upright		demi-dressé		halbaufrecht	semierecto	Franquette, Hartley, Marbot, Liaoning 1, Liaoning 4, Shaanhe 1, Chuanhe 2, Alsószentiváni 117	2
	spreading		étalé		breitwüchsig	extendido	Gustine, Payne, Shangsong 6, Vina, Jinfeng, Jinlong 1, Jinlong 2, Xilin 2, Zhonglin 5, Luguang, Milotai 10	3
2.	QN	VG	(+)	(a), (b)				
	Tree: branching		Arbre : ramification		Baum: Verzweigung	Árbol: ramificación		
	weak		faible		gering	escasa	Vina, Chico, Huashan 5, Shangluo 3, Xisiyu 1	3
	medium		moyenne		mittel	media	Franquette, Hartley, Marbot, Shangsong 6, Lubo, Xinzaofeng, Qinyou 1, Chuanhe 2	5
	strong		forte		stark	abundante	Corne, Jinfeng, Liaoning 1, Liaoning 4, Parisienne, Shaanhe 1, Xiangling, Xilin 2, Zhonglin 1, Zhonglin 5, Luguang, Daixiang, Luguo 2, Xifu 2	7

		English		français		deutsch		español		Example Varieties Exemples Beispielssorten Variedades ejemplo		Note/ Nota		
3. (*)		PQ	VG	(+)	(b)									
	Bud: shape		Bourgeon : forme		Knospe: Form		Yema: forma							
	circular		circulaire		kreisförmig		circular		Jinlong 1, Xiangling, Xilin 2, Zhonglin 1, Luguang, Xinzaofeng, Daixiang, Luguo 2, Milotai 10			1		
	semi-circular		semi-circulaire		halb kreisförmig		semicircular		Fenghui			2		
	triangular		triangulaire		dreieckig		triangular		Liaoning 4, Zhenzhuhetao, Chuanhe 2			3		
4. (*)	QN	VG	(+)											
	Lateral leaflet: shape		Foliole latérale : forme		Seitenfiederblatt: Form		Folíolo lateral: forma							
	narrow elliptic		elliptique étroite		schmal elliptisch		elíptico estrecho		Hartley, Payne, Shangsong 6, Vina, Daifeng, Liaoning 1, Daixiang			3		
	medium elliptic		elliptique moyenne		mittel elliptisch		elíptico medio		Corne, Franquette, Marbot			5		
	broad elliptic		elliptique large		breit elliptisch		elíptico ancho		Chase D 9, Adam 10			7		
5.	QL	VG	(+)											
	Plant: second flowering		Plante : deuxième floraison		Pflanze: Zweite Blüte		Planta: segunda floración							
	absent		absente		fehlend		ausente		Jinlong 1, Milotai 10			1		
	present		présente		vorhanden		presente		Liaoning 4			9		
6. (*)	QN	MG	(+)	(c)										
	Female flower: number per cluster		Fleur femelle : nombre par bouquet		Weibliche Blüte: Anzahl pro Gruppe		Flor femenina: número por grupo							
	1-2		1-2		1-2		1-2		Jinlong 1, Xiangling, Xilin 2, Luguang			1		
	3-4		3-4		3-4		3-4		Shaanhe 1			2		
	5-10		5-10		5-10		5-10					3		
	11-20		11-20		11-20		11-20		Qinyou 1, Tisa			4		
	more than 20		plus de 20		mehr als 20		más de 20		Chuanhetao			5		

		English		français		deutsch		español		Example Varieties Exemples Beispielssorten Variedades ejemplo		Note/ Nota
7.	(*)	QN	VG		(c)							
		Female flower: intensity of yellow color of stigma		Fleur femelle : intensité de la couleur jaune du stigmate		Weibliche Blüte: Intensität der gelben Farbe der Narbe		Flor femenina: intensidad del color amarillo del estigma				
		light		claire		hell		claro		Daifeng, Daixiang, Milotai 10		1
		medium		moyenne		mittel		medio		Jinlong 1, Jinlong 2, Xiangling, Zhonglin 1, Zhonglin 5, Xinzaofeng		2
		dark		foncée		dunkel		oscuro		Xifu 2		3
8.	(*)	PQ	VG	(+)								
		Fruit: setting type		Fruit : type de position		Frucht: Ansatztyp		Fruto: tipo de fructificación				
		solitary		solitaire		einzeln		en solitario		Jinlong 1, Milotai 10		1
		binate		par deux		doppelt		en pares		Daifeng, Jinlong 1, Liaoning 1, Liaoning 4, Xiangling, Xilin 2, Zhonglin 5, Luguang, Fenghui, Daixiang, Luguo 2		2
		fascicled		en paquet		in Büscheln		en fascículos		Shaanhe 1		3
		bunchy		en grappe		in Trauben		en racimos		Chuanzihetao		4
9.		QN	MG/VG		(d)							
		Nut: size		Noix : grosseur		Nuß: Größe		Nuez: tamaño				
		small		petite		klein		pequeña		Chico, Grandjean, Zhenzhuhetao		3
		medium		moyenne		mittel		mediana		Franquette, Liaoning 4, Shaanhe 1, Honghetao		5
		large		grosse		groß		grande		Hartley, Daifeng, Jinlong 1, Jinlong 2, Lübo, Xiangling, Xilin 2, Zhonglin 1, Zhonglin 5, Luguang, Fenghui, Xinzaofeng, Daixiang, Luguo 2, Milotai 10, Sunland		7

		English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
10.	(*)	PQ	VG	(+)	(d)		
	Nut: shape in lateral view	Noix : forme en vue latérale		Nuß: Form in Seitenansicht	Nuez: forma en vista lateral		
	broad ovate	ovale large		breit eiförmig	oval ancha	Marbot, Payne, Serr	1
	ovate	ovale		eiförmig	oval	Gustine, Jinfeng	2
	triangular	triangulaire		dreieckig	triangular	Hartley	3
	elliptic	elliptique		elliptisch	elíptica	Corne, Sorrento, Franquette, Daifeng, Xilin 2	4
	circular	circulaire		kreisförmig	circular	Jinlong 1, Jinlong 2, Liaoning 4, Meylannaise, Xiangling, Zhonglin 1, Zhonglin 5, Milotai 10	5
	broad elliptic	elliptique large		breit elliptisch	elíptica ancha	Parisienne, Luguang	6
	oblong	oblongue		breitrund	oblonga	Mumahetao, Milotai bötermő, Sunland	7
11.	(*)	PQ	VG	(+)	(d)		
	Nut: shape in ventral view	Noix : forme en vue ventrale		Nuß: Form in Bauchansicht	Nuez: forma en vista ventral		
	broad ovate	ovale large		breit eiförmig	oval ancha	Payne, Serr, Xiangling	1
	triangular	triangulaire		dreieckig	triangular	Hartley	2
	ovate	ovale		eiförmig	oval	Gustine, Jinfeng	3
	circular	circulaire		kreisförmig	circular	Meylannaise, Milotai 10	4
	broad elliptic	elliptique large		breit elliptisch	elíptica ancha	Franquette	5
	oblanceolate	arrondie aplatie		abgeplattet	achatada	Yuanbao	6

		English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
12.	(*)	PQ	VG	(+)	(d)		
	Nut: shape in cross section	Noix : forme en section transversale		Nuß: Form im Querschnitt	Nuez: forma en sección transversal		
	reniform	réniforme		nierenförmig	reniforme		1
	oblite	arrondie aplatie		abgeplattet	achatada	Franquette, Chico, Liaoning 1, Jupiter	2
	elliptic	elliptique		elliptisch	elíptica	Corne, Hartley, Serr	3
	circular	circulaire		kreisförmig	circular	Marbot, Payne, Xiangling, Milotai 10, Victoria	4
13.	(*)	PQ	VG	(+)	(d)		
	Nut: shape of base in lateral view (facing the suture)	Noix : forme de la base en vue latérale (en face de la suture)		Nuß: Form der Basis in Seitenansicht (der Naht zugewandt)	Nuez: forma de la base en vista lateral (con la sutura de frente)		
	cuneate	cunée		keilförmig	cuneada	Corne, Milotai bötermő	1
	rounded	arrondie		abgerundet	redondeada	Franquette, Payne, Chico, Serr, Xiangling	2
	truncate	tronquée		stumpf	truncada	Parisienne	3
	emarginate	récurrente		eingesenkt	emarginada	Hartley	4
14.	(*)	PQ	VG	(+)	(d)		
	Nut: shape of apex in lateral view (facing the suture, excluding tip)	Noix : forme du sommet en vue latérale (en face de la suture, à l'exclusion de la pointe)		Nuß: Form der Spitze in Seitenansicht (der Naht zugewandt, ohne Spitze)	Nuez: forma del ápice en vista lateral (con la sutura de frente, excluida la punta)		
	obtuse	obtus		abgestumpft	obtuso	Vina	1
	rounded	arrondi		abgerundet	redondeado	Zhonglin 1	2
	truncate	tronqué		stumpf	truncado	Zhonglin 5, Milotai bötermő	3
	emarginate	récurrent		eingesenkt	emarginado	Xiangling	4

		English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
15.	(*)	QN	VG	(+)	(d)		
		Nut: length of tip	Noix : longueur de la pointe	Nuß: Länge der Spitze	Nuez: longitud de la punta		
		absent or short	absente ou courte	fehlend oder kurz	ausente o corta	Grandjean, Xiangling, Milotai 10	1
		medium	moyenne	mittel	mediana	Corne, Hartley, Chico, Hexuan	2
		long	longue	lang	larga	Franquette, Marbot, Payne, Serr, Victoria	3
16.	(*)	QN	VG	(+)	(d)		
		Nut: extent of pad around suture	Noix : étendue du bourrelet autour de la suture	Nuß: Ausdehnung des Wulstes um die Naht herum	Nuez: extensión del almohadillado a lo largo de la sutura		
		on upper half	sur la moitié supérieure	an der oberen Hälfte	en la mitad superior	Hartley, Marbot, Chico, Parisienne, Xiangling	1
		on upper 2/3	sur les deux tiers supérieurs	am oberen zweiten Drittel	en los dos tercios superiores	Franquette, Gustine, Payne, Liaoning 1, Liaoning 4, Pedro, Jupiter	2
		on whole length	sur la totalité de la longueur	über die gesamte Länge	a todo lo largo	Honghuadian 1	3
17.	(*)	QN	VG		(d)		
		Nut: prominence of pad on suture	Noix : importance du bourrelet de suture	Nuß: Ausprägung des Wulstes auf der Naht	Nuez: prominencia del almohadillado de la sutura		
		very weak	très faible	sehr gering	muy leve	Luguang	1
		weak	faible	gering	leve	Jinlong 2, Chuanhe 2	2
		medium	moyenne	mittel	medio	Chico, Grandjean	3
		strong	forte	stark	marcado	Franquette, Hartley, Marbot, Payne, Serr	4
		very strong	très forte	sehr stark	muy marcado	Xifu 2	5

	English		français		deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
18.	QN	VG	(+)	(d)				
	Nut: width of pad on suture in lateral view		Noix : largeur du bourrelet de suture en vue latérale		Nuß: Breite des Wulstes auf der Naht in Seitenansicht	Nuez: anchura del almohadillado de la sutura en vista lateral		
	narrow		étroit		schmal	estrecho	Chico, Grandjean, Parisienne, Xiangling	1
	medium		moyen		mittel	mediano	Hartley, Gustine	3
	broad		large		breit	ancho	Corne, Marbot, Payne, Serr	5
19.	QN	VG		(d)				
	Nut: depth of groove along pad on suture		Noix : profondeur des anfractuosités le long du bourrelet de suture		Nuß: Tiefe der Furche entlang dem Wulst auf der Naht	Nuez: profundidad del surco a lo largo del almohadillado de la sutura		
	shallow		peu profondes		flach	poco profundo	Chico, Grandjean, Parisienne, Xiangling	1
	medium		moyennes		mittel	medio	Hartley, Gustine	3
	deep		profondes		tief	profundo	Corne, Marbot, Payne, Serr	5
20.	PQ	VG	(+)	(d)				
	Nut: structure of surface of shell		Noix : structure de la surface de la coque		Nuß: Oberflächenstruktur der Schale	Nuez: estructura de la superficie de la cáscara		
	slightly grooved		peu sillonnée		kaum gefurcht	ligeramente acanalada	Liaoning 1, Liaoning 4, Luguang	1
	moderately grooved		moyennement sillonnée		mäßig gefurcht	moderadamente acanalada	Chico, Jinlong 1, Jinlong 2, Lübo, Xiangling, Fenghui, Xinzaofeng, Milotai intenzív	2
	strongly grooved		fortement sillonnée		stark gefurcht	intensamente acanalada	Hartley, Xilin 2, Tiszacsécsi 83	3
	embossed		bosselée		höckerig	protuberancias irregulares	Erbazi	4

	English		français		deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota	
21.	PQ	VG	(d)						
	Nut: color of shell		Noix : couleur de la coque		Nuß: Farbe der Schale	Nuez: color de la cáscara			
	yellow		jaune		gelb	amarillo	Xiangling	1	
	light brown		marron clair		hellbraun	marrón claro	Zhonglin 1, Milotai 10	2	
	medium brown		marron moyen		mittelbraun	marrón medio	Honghetao	3	
22.	QN	VG	(+)	(d)					
	Nut: thickness of dividing membranes		Noix : épaisseur des cloisons		Nuß: Dicke der Trennmembranen	Nuez: grosor de las membranas divisorias			
	very thin		très minces		sehr dünn	muy delgadas	Daifeng, Liaoning 1, Liaoning 4, Lübo, Shaanhe 1, Xiangling, Xilin 2, Zhonglin 1, Zhonglin 5, Fenghui, Daixiang, Luguo 2, Milotai 10	1	
	thin		minces		dünn	delgadas	Payne, Chico, Serr, Luguang	2	
	medium		moyennes		mittel	medias	Franquette, Marbot, Xinzaofeng, Honghetao	3	
	thick		épaisses		dick	gruesas	Corne	4	
	very thick		très épaisses		sehr dick	muy gruesas	Aodidaguanmao, Jilong	5	
23.	PQ	VG	(+)	(d)					
	Nut: inner pleat wall of shell		Noix : valve ridée de la coque		Nuß: Faltigkeit der Innenwand der Schale	Nuez: pared rugosa interna de la cáscara			
	papery		fine comme du papier		papierähnlich	papirácea	Daifeng, Liaoning 1, Liaoning 4, Xiangling, Zhonglin 1, Zhonglin 5, Luguang, Fenghui, Daixiang, Luguo 2	1	
	coriaceous		comme du cuir		lederartig	coriácea	Xinzaofeng	2	
	ligneous		ligneuse		holzig	leñosa	Baipihetao	3	

		English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
24. (*)	QN	MG/VG	(d)				
	Nut: thickness of shell	Noix : épaisseur de la coquille		Nuß: Dicke der Schale	Nuez: grosor de la cáscara		
	very thin	très mince		sehr dünn	muy delgada	Daifeng, Liaoning 1, Liaoning 4, Lübo, Pedro, Serr, Xiangling, Luguang, Fenghui, Luguo 2	1
	thin	mince		dünn	delgada	Payne, Chico, Jinlong 1, Jinlong 2, Serr, Xilin 2, Zhonglin 1, Zhonglin 5, Xinzaofeng, Daixiang	2
	medium	moyenne		mittel	media	Franquette, Hartley, Marbot, Chahetao, Milotai 10	3
	thick	épaisse		dick	gruesa	Corne, Shitou	4
	very thick	très épaisse		sehr dick	muy gruesa		5
25. (*)	PQ	VG	(e)				
	Kernel: color of endopleura	Cerneau : couleur du tégument		Kern: Farbe der inneren Samenhaut	Semilla: color de la endopleura		
	white	blanc		weiß	blanco	Jinmian 2	1
	yellowish white	blanc jaunâtre		gelblich weiß	blanco amarillento	Liaoning 1, Eszterhazy II	2
	yellow	jaune		gelb	amarillo	Daifeng, Milotai 10	3
	red	rouge		rot	rojo	Hongranghetao, Honghetao	4
	purple	violet		purpurn	púrpura	Chuanhe 2, Sychrov	5
	yellow brown	marron jaune		gelbbraun	marrón amarillento	Baipihetao	6
	light brown	marron clair		hellbraun	marrón claro	Shangsong 6, Alsószentiván 117	7
	medium brown	marron moyen		mittelbraun	marrón medio	Zhonglin 5	8
	dark brown	marron foncé		dunkelbraun	marrón oscuro		9

	English		français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
26.	QN	MG	(e)				
	Kernel: percentage of weight relative to total weight of nut	Cerneau : pourcentage du poids par rapport au poids total du fruit	Kern: relativer Anteil am Gesamtgewicht der Nuß	Semilla: porcentaje del peso en relación con el peso total de la nuez			
	very low	très faible	sehr gering	muy bajo	Corne	1	
	low	faible	gering	bajo	Marbot	3	
	medium	moyen	mittel	medio	Sorrento, Franquette, Hartley, Pedro	5	
	high	élevé	hoch	alto	Payne, Vina, Chase D 9, Daifeng, Jinlong 2, Liaoning 1, Liaoning 4, Lübo, Zhonglin 1, Zhonglin 5, Luguang, Fenghui, Xinzaofeng, Daixiang, Luguo 2, Milotai 10	7	
	very high	très élevé	sehr hoch	muy alto	Jinlong 1, Serr, Shaanhe 1, Xiangling, Xilin 2	9	
27	QN	VG	(+)	(e)			
	Kernel: ease of removal from shell	Cerneau : facilité d'extraction de la coquille	Kern: Leichtigkeit des Entfernens von der Schale	Semilla: facilidad con que se extrae de la cáscara			
	very easy	très aisée	sehr leicht	muy fácil	Payne, Pedro, Serr, Milotai 10	1	
	easy	aisée	leicht	fácil	Franquette, Hartley, Marbot	2	
	medium	moyenne	mittel	media	Jinlong 2, Meylannaise	3	
	difficult	difficile	schwer	difícil	Corne	4	
	very difficult	très difficile	sehr schwer	muy difícil	Aodidaguanmao, Jilong	5	

		English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
28.	QN	MG	(+)				
	Time of beginning of female flowering		Époque de début de floraison femelle	Zeitpunkt des Beginns der weiblichen Blüte	Época de inicio de la floración femenina		
	early		précoce	früh	temprana	Sorrento, Chase D 9, Lübo, Xilin 2, Luguo 2	3
	medium		moyenne	mittel	intermedia	Marbot, Daifeng, Luguang, Fenghui, Daixiang, Milotai 10	5
	late		tardive	spät	tardía	Jinlong 1, Jinlong 2, Liaoning 1, Liaoning 4, Milotai kései, Bonifác	7
29.	QN	MG	(+)				
	Time of beginning of male flowering		Époque de début de floraison mâle	Zeitpunkt des Beginns der männlichen Blüte	Época de inicio de la floración masculina		
	early		précoce	früh	temprana	Sorrento, Gustine, Chase D 9, Xiangling, Xilin 2, Luguang, Fenghui, Xinzaofeng, Luguo 2, Milotai 10	3
	medium		moyenne	mittel	intermedia	Marbot, Lübo, Shaanhe 1	5
	late		tardive	spät	tardía	Franquette, Jinlong 1, Jinlong 2, Liaoning 1, Liaoning 4, Parisienne, Zhonglin 1, Zhonglin 5	7
30. (*)	QN	MG	(c)				
	Time of male flowering compared to female flowering		Époque de début de floraison mâle par rapport à l'époque de début de floraison femelle	Zeitpunkt der Blüte der männlichen Blüte im Vergleich zur weiblichen Blüte	Época de la floración masculina en relación con la floración femenina		
	before (protandry)		avant (protandrie)	vorausgehend (Protandrie)	anterior (protandria)	Franquette, Marbot, Payne, Liaoning 1, Liaoning 4, Xiangling	1
	simultaneous (homogamy)		simultanée (homogamie)	gleichzeitig (Homogamie)	simultánea (homogamia)	Chico, Meylannaise, Xilin 2	2
	after (protogyny)		après (protogynie)	nachfolgend (Protoynie)	posterior (protoginia)	Lübo, Milotai 10	3

	English		français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
31.	QN	MG					
	Time of harvest maturity		Époque de maturité de récolte	Zeitpunkt der Erntereife	Época de madurez para la cosecha		
	early		précoce	früh	temprana	Lübo, Xiangling, Zhonglin 5, Luguang, Fenghui, Luguo 2, Milotai 10	3
	medium		moyenne	mittel	intermedia	Payne, Chico, Daifeng, Grandjean, Serr, Zhonglin 1, Xinzaofeng, Daixiang	5
	late		tardive	spät	tardía	Jinlong 1, Jinlong 2, Liaoning 1, Liaoning 4, Xilin 2, Milotai kesei	7

8. Explanations on the Table of Characteristics


8.1 *Explanations covering several characteristics*

Characteristics containing the following key in the second column of the Table of Characteristics should be examined as indicated below:

- (a) Observations on tree should be made in dormant season.
- (b) Observations on branch should be made on vegetative branches from the middle part of the canopy in dormant season.
- (c) Observations on flowers should be carried out during full-blossom period.
- (d) Observations on nuts should be made on physiological ripe nuts excluding the pericarp immediately after 25% of the pericarp cracked. Take 30 nuts randomly from each tree.
- (e) Observations on the kernel should be made when the water content is less than 8%.

8.2 *Explanations for individual characteristics*

Ad. 1: Tree: growth habit



Ad. 2: Tree: branching

Observations should relate to the number of branches with the degree of branching being indicated by the density of lateral branches and shoots, excluding fruiting shoots.

Ad. 3: Bud: shape

Observations on buds should be made on terminal buds of branches.

Ad. 4: Lateral leaflet: shape

Observations on leaflets should be made on lateral leaves from the middle part of the canopy on the sunny side.



Ad. 5: Plant: second flowering



Ad. 8: Fruit: setting type






Ad. 10: Nut: shape in lateral view

		← broadest part →		
		below middle	at middle	
width (ratio length/width)				
narrow (high)			 7 oblong	
medium (medium)		 2 ovate	 4 elliptic	 6 broad elliptic
broad (low)		 1 broad ovate	 3 triangular	 5 circular

Ad. 11: Nut: shape in ventral view

Observations should be made facing the suture.

width (ratio length/width)	← broadest part →		
	below middle	at middle	
narrow (high)	 1 broad ovate		
medium (medium)		 3 ovate	 5 broad elliptic
broad (low)		 2 triangular	 4 circular
			 6 oblanceolate

Ad. 12: Nut: shape in cross section

← broadest part → at middle		
width (ratio length/width)		
narrow (high)		 3 elliptic
medium (medium)		 4 circular
broad (low)	 1 reniform	 2 oblate


Ad. 13: Nut: shape of base in lateral view (facing the suture)




Ad. 14: Nut: shape of apex in lateral view (facing the suture, excluding tip)




Ad. 15: Nut: length of tip




Ad. 16: Nut: extent of pad around suture



Ad. 18: Nut: width of pad on suture in lateral view




Ad. 20: Nut: structure of surface of shell




a = grooved  
b = embossed

Ad. 22: Nut: thickness of dividing membranes



Ad. 23: Nut: inner pleat wall of shell



Ad. 24: Nut: thickness of shell

Thickness of the mid part of the shell should be measured and take the average value (accurate to 0.1 mm) as the thickness of shell.

Ad. 27: Kernel: ease of removal from shell

Crack the shell and remove the kernel. Assess the ease of removal according to the situation of the kernel:

- 1: whole or a half of the kernel;
- 2: a quarter of the kernel;
- 3: smashed kernel;
- 4: unable to take out the kernel normally.

Ad. 28: Time of beginning of female flowering

The time of beginning of female flowering should be observed when 5% of the female flowers are in full bloom (at full development of stigmas).

Ad. 29: Time of beginning of male flowering

The time of beginning of male flowering should be observed when 10% of the catkins are in full bloom (at dehiscence of pollen).

9. Literature

IPGRI, 1994: descriptors for walnut (*Juglans* spp.). International Plant Genetic Resource Institute, Rome, IT.

Liu, Q.Z., Zhang, L.S., 2007: Descriptors and Data Standard for walnut (*Juglans regia* L.). China Agriculture Press. Beijing, CN.

Pei, D., Lu, X.Z., 2011: Walnut germplasm resources in China. China forestry publishing house. Beijing, CN.

10. Technical Questionnaire

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
		Application date: (not to be filled in by the applicant)
<b>TECHNICAL QUESTIONNAIRE</b> to be completed in connection with an application for plant breeders' rights		
1. Subject of the Technical Questionnaire		
1.1 Botanical name	<i>Juglans regia L.</i>	
1.2 Common name	Walnut, English Walnut	
2. Applicant		
Name		
Address		
Telephone No.		
Fax No.		
E-mail address		
Breeder (if different from applicant)		
3. Proposed denomination and breeder's reference		
Proposed denomination (if available)		
Breeder's reference		

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
#4. Information on the breeding scheme and propagation of the variety		
4.1 Breeding scheme		
Variety resulting from:		
4.1.1 Crossing		
(a) controlled cross (please state parent varieties)	x	[ ]
(.....)		(.....)
female parent		male parent
(b) partially known cross (please state known parent variety(ies))	x	[ ]
(.....)		(.....)
female parent		male parent
(c) unknown cross	x	[ ]
4.1.2 Mutation		
(please state parent variety)		
4.1.3 Discovery and development		
(please state where and when discovered and how developed)		
4.1.4 Other		
(please provide details)		

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

4.2 Method of propagating the variety

4.2.1 Vegetative propagation

(a) grafting (budding)

[ ]

(b) Other (state method)

[ ]

4.2.2 Other

(Please provide details)

[ ]

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
<p>5. Characteristics of the variety to be indicated (the number in brackets refers to the corresponding characteristic in Test Guidelines; please mark the note which best corresponds).</p>		
Characteristics	Example Varieties	Note
<b>5.1 Tree: growth habit (1)</b>		
upright	Corne, Daifeng, Daixiang, Fenghui, Sorrento, Xinzaofeng, Zhonglin 1	1 [ ]
semi-upright	Alsószentiváni 117, Chuanhe 2, Franquette, Hartley, Liaoning 1, Liaoning 4, Marbot, Shaanhe 1	2 [ ]
spreading	Gustine, Jinfeng, Jinlong 1, Jinlong 2, Luguang, Milotai 10, Payne, Shangsong 6, Vina, Xilin 2, Zhonglin 5	3 [ ]
<b>5.2 Bud: shape (3)</b>		
circular	Daixiang, Jinlong 1, Luguang, Luguo 2, Milotai 10, Xiangling, Xilin 2, Xinzaofeng, Zhonglin 1	1 [ ]
semi-circular	Fenghui	2 [ ]
triangular	Chuanhe 2, Liaoning 4, Zhenzhuhetao	3 [ ]
<b>5.3 Lateral leaflet: shape (4)</b>		
very narrow elliptic		1 [ ]
very narrow elliptic to narrow elliptic		2 [ ]
narrow elliptic	Daifeng, Daixiang, Hartley, Liaoning 1, Payne, Shangsong 6, Vina	3 [ ]
narrow elliptic to medium elliptic		4 [ ]
medium elliptic	Corne, Franquette, Marbot	5 [ ]
medium elliptic to broad elliptic		6 [ ]
broad elliptic	Adam 10, Chase D 9	7 [ ]
broad elliptic to very broad elliptic		8 [ ]
very broad elliptic		9 [ ]
<b>5.4 Female flower: number per cluster (6)</b>		
1-2	Jinlong 1, Luguang, Xiangling, Xilin 2	1 [ ]
3-4	Shaanhe 1	2 [ ]
5-10		3 [ ]
11-20	Qinyou 1, Tisa	4 [ ]
more than 20	Chuanhetao	5 [ ]

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
Characteristics	Example Varieties	Note
<b>5.5 (7) Female flower:intensity of yellow color of stigma</b>		
light	Daifeng, Daixiang, Milotai 10	1 [ ]
medium	Jinlong 1, Jinlong 2, Xiangling, Xinzaofeng, Zhonglin 1, Zhonglin 5	2 [ ]
dark	Xifu 2	3 [ ]
<b>5.6 (8) Fruit: setting type</b>		
solitary	Jinlong 1, Milotai 10	1 [ ]
binate	Daifeng, Daixiang, Fenghui, Jinlong 1, Liaoning 1, Liaoning 4, Luguang, Luguo 2, Xiangling, Xilin 2, Zhonglin 5	2 [ ]
fascicled	Shaanhe 1	3 [ ]
bunchy	Chuanzihetao	4 [ ]
<b>5.7 (10) Nut: shape in lateral view</b>		
broad ovate	Marbot, Payne, Serr	1 [ ]
ovate	Gustine, Jinfeng	2 [ ]
triangular	Hartley	3 [ ]
elliptic	Corne, Daifeng, Franquette, Sorrento, Xilin 2	4 [ ]
circular	Jinlong 1, Jinlong 2, Liaoning 4, Meylannaise, Milotai 10, Xiangling, Zhonglin 5 [ ] 1, Zhonglin 5	
broad elliptic	Luguang, Parisienne	6 [ ]
oblong	Milotai bőtermő, Mumahetao, Sunland	7 [ ]
<b>5.8 (11) Nut: shape in ventral view</b>		
broad ovate	Payne, Serr, Xiangling	1 [ ]
triangular	Hartley	2 [ ]
ovate	Gustine, Jinfeng	3 [ ]
circular	Meylannaise, Milotai 10	4 [ ]
broad elliptic	Franquette	5 [ ]
oblanceolate	Yuanbao	6 [ ]
<b>5.9 (12) Nut: shape in cross section</b>		
reniform		1 [ ]
oblanceolate	Chico, Franquette, Jupiter, Liaoning 1	2 [ ]
elliptic	Corne, Hartley, Serr	3 [ ]
circular	Marbot, Milotai 10, Payne, Victoria, Xiangling	4 [ ]

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
Characteristics	Example Varieties	Note
<b>5.10 Nut: shape of base in lateral view (facing the suture) (13)</b>		
cuneate	Corne, Milotai bőtermő	1 [ ]
rounded	Chico, Franquette, Payne, Serr, Xiangling	2 [ ]
truncate	Parisienne	3 [ ]
emarginate	Hartley	4 [ ]
<b>5.11 Nut: shape of apex in lateral view (facing the suture, excluding tip) (14)</b>		
obtuse	Vina	1 [ ]
rounded	Zhonglin 1	2 [ ]
truncate	Milotai bőtermő, Zhonglin 5	3 [ ]
emarginate	Xiangling	4 [ ]
<b>5.12 Nut: length of tip (15)</b>		
absent or short	Grandjean, Milotai 10, Xiangling	1 [ ]
medium	Chico, Corne, Hartley, Hexuan	2 [ ]
long	Franquette, Marbot, Payne, Serr, Victoria	3 [ ]
<b>5.13 Nut: extent of pad around suture (16)</b>		
on upper half	Chico, Hartley, Marbot, Parisienne, Xiangling	1 [ ]
on upper 2/3	Franquette, Gustine, Jupiter, Liaoning 1, Liaoning 4, Payne, Pedro	2 [ ]
on whole length	Honghuadian 1	3 [ ]
<b>5.14 Nut: prominence of pad on suture (17)</b>		
very weak	Luguang	1 [ ]
weak	Chuanhe 2, Jinlong 2	2 [ ]
medium	Chico, Grandjean	3 [ ]
strong	Franquette, Hartley, Marbot, Payne, Serr	4 [ ]
very strong	Xifu 2	5 [ ]
<b>5.15 Nut: thickness of shell (24)</b>		
very thin	Daifeng, Fenghui, Liaoning 1, Liaoning 4, Luguang, Luguo 2, Lübo, Pedro, Serr, Xiangling	1 [ ]
thin	Chico, Daixiang, Jinlong 1, Jinlong 2, Payne, Serr, Xilin 2, Xinzaofeng, Zhonglin 1, Zhonglin 5	2 [ ]
medium	Chahetao, Franquette, Hartley, Marbot, Milotai 10	3 [ ]
thick	Corne, Shitou	4 [ ]
very thick		5 [ ]

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
Characteristics	Example Varieties	Note
<b>5.16 Kernel: color of endopleura (25)</b>		
white	Jinmian 2	1 [ ]
yellowish white	Eszterhazy II, Liaoning 1	2 [ ]
yellow	Daifeng, Milotai 10	3 [ ]
red	Honghetao, Hongranghetao	4 [ ]
purple	Chuanhe 2, Sychrov	5 [ ]
yellow brown	Baipihetao	6 [ ]
light brown	Alsószentiváni 117, Shangsong 6	7 [ ]
medium brown	Zhonglin 5	8 [ ]
dark brown		9 [ ]
<b>5.17 Time of male flowering compared to female flowering (30)</b>		
before (protandry)	Franquette, Liaoning 1, Liaoning 4, Marbot, Payne, Xiangling	1 [ ]
simultaneous (homogamy)	Chico, Meylannaise, Xilin 2	2 [ ]
after (protogyny)	Lübo, Milotai 10	3 [ ]

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:	
6. Similar varieties and differences from these varieties			
<p><i>Please use the following table and box for comments to provide information on how your candidate variety differs from the variety (or varieties) which, to the best of your knowledge, is (or are) most similar. This information may help the examination authority to conduct its examination of distinctness in a more efficient way.</i></p>			
Denomination(s) of variety(ies) similar to your candidate variety	Characteristic(s) in which your candidate variety differs from the similar variety(ies)	Describe the expression of the characteristic(s) for the <b>similar</b> variety(ies)	Describe the expression of the characteristic(s) for <b>your</b> candidate variety
<i>Example</i>	<i>Nut: thickness of shell</i>	<i>thick</i>	<i>medium</i>
Comments:			

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
#7. Additional information which may help in the examination of the variety		
7.1 In addition to the information provided in sections 5 and 6, are there any additional characteristics which may help to distinguish the variety?		
Yes [ ]	No [ ]	
(If yes, please provide details)		
7.2 Are there any special conditions for growing the variety or conducting the examination?		
Yes [ ]	No [ ]	
(If yes, please provide details)		
7.3 Other information		
<p>A representative color photograph of the variety displaying its main distinguishing feature(s), should accompany the Technical Questionnaire. The photograph will provide a visual illustration of the candidate variety which supplements the information provided in the Technical Questionnaire.</p> <p>The key points to consider when taking a photograph of the candidate variety are:</p> <ul style="list-style-type: none"><li>• Indication of the date and geographic location</li><li>• Correct labeling (breeder's reference)</li><li>• Good quality printed photograph (minimum 10 cm x 15 cm) and/or sufficient resolution electronic format version (minimum 960 x 1280 pixels)"</li></ul> <p>Further guidance on providing photographs with the Technical Questionnaire is available in document TGP/7 "Development of Test Guidelines", Guidance Note 35 (<a href="http://www.upov.int/tgp/en/">http://www.upov.int/tgp/en/</a>). [The link provided may be deleted by members of the Union when developing authorities' own test guidelines.]</p>		

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

8. Authorization for release

- (a) Does the variety require prior authorization for release under legislation concerning the protection of the environment, human and animal health?

Yes [ ] No [ ]

- (b) Has such authorization been obtained?

Yes [ ] No [ ]

If the answer to (b) is yes, please attach a copy of the authorization.

9. Information on plant material to be examined or submitted for examination

9.1 The expression of a characteristic or several characteristics of a variety may be affected by factors, such as pests and disease, chemical treatment (e.g. growth retardants or pesticides), effects of tissue culture, different rootstocks, scions taken from different growth phases of a tree, etc.

9.2 The plant material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If the plant material has undergone such treatment, full details of the treatment must be given. In this respect, please indicate below, to the best of your knowledge, if the plant material to be examined has been subjected to:

- |     |   |         |        |
|-----|---|---------|--------|
| (a) | Microorganisms (e.g. virus, bacteria, phytoplasma)    | Yes [ ] | No [ ] |
| (b) | Chemical treatment (e.g. growth retardant, pesticide) | Yes [ ] | No [ ] |
| (c) | Tissue culture  | Yes [ ] | No [ ] |
| (d) | Other factors   | Yes [ ] | No [ ] |

Please provide details for where you have indicated "yes".

.....

10. I hereby declare that, to the best of my knowledge, the information provided in this form is correct:

Applicant's name

Signature

 Date 

[End of document]