



TG/BRASS_JUN(proj.8)

ORIGINAL: English

DATE: 2020-09-11

INTERNATIONAL UNION FOR THE PROTECTION OF NEW VARIETIES OF PLANTS

Geneva

DRAFT

BROWN MUSTARD

UPOV Code(s):

BRASS_JUN

Brassica juncea (L.) Czern.**GUIDELINES****FOR THE CONDUCT OF TESTS***prepared by an expert from Japan**to be considered by the**Technical Committee at its fifty-sixth session
to be held in Geneva on October 26 and 27, 2020**Disclaimer: this document does not represent UPOV policies or guidance*

Alternative names:*

| <i>Botanical name</i> | <i>English</i> | <i>French</i> | <i>German</i> | <i>Spanish</i> |
|---------------------------------------|---|----------------|---------------|--------------------------------------|
| <i>Brassica juncea</i> (L.) Czern. | Brown mustard, Indian mustard, Oriental mustard | Moutarde brune | Sareptasenf | Mostaza de Sarepta, Mostaza india |

The purpose of these guidelines ("Test Guidelines") is to elaborate the principles contained in the General Introduction (document TG/1/3), and its associated TGP documents, into detailed practical guidance for the harmonized examination of distinctness, uniformity and stability (DUS) and, in particular, to identify appropriate characteristics for the examination of DUS and production of harmonized variety descriptions.

ASSOCIATED DOCUMENTS

These Test Guidelines should be read in conjunction with the General Introduction and its associated TGP documents.

* These names were correct at the time of the introduction of these Test Guidelines but may be revised or updated. [Readers are advised to consult the UPOV Code, which can be found on the UPOV Website (www.upov.int), for the latest information.]

| TABLE OF CONTENTS | PAGE |
|--|-----------|
| 1. SUBJECT OF THESE TEST GUIDELINES..... | <u>3</u> |
| 2. MATERIAL REQUIRED..... | <u>3</u> |
| 3. METHOD OF EXAMINATION..... | <u>3</u> |
| 3.1 Number of Growing Cycles..... | <u>3</u> |
| 3.2 Testing Place..... | <u>3</u> |
| 3.3 Conditions for Conducting the Examination..... | <u>3</u> |
| 3.4 Test Design..... | <u>3</u> |
| 3.5 Additional Tests..... | <u>3</u> |
| 4. ASSESSMENT OF DISTINCTNESS, UNIFORMITY AND STABILITY..... | <u>4</u> |
| 4.1 Distinctness..... | <u>4</u> |
| 4.2 Uniformity..... | <u>5</u> |
| 4.3 Stability..... | <u>5</u> |
| 5. GROUPING OF VARIETIES AND ORGANIZATION OF THE GROWING TRIAL..... | <u>5</u> |
| 6. INTRODUCTION TO THE TABLE OF CHARACTERISTICS..... | <u>6</u> |
| 6.1 Categories of Characteristics..... | <u>6</u> |
| 6.2 States of Expression and Corresponding Notes..... | <u>6</u> |
| 6.3 Types of Expression..... | <u>6</u> |
| 6.4 Example Varieties..... | <u>6</u> |
| 6.5 Legend..... | <u>7</u> |
| 7. TABLE OF CHARACTERISTICS/TABLEAU DES CARACTÈRES/MERKMALSTABELLE/TABLA DE CARACTERES..... | <u>8</u> |
| 8. EXPLANATIONS ON THE TABLE OF CHARACTERISTICS..... | <u>15</u> |
| 8.1 Explanations covering several characteristics..... | <u>15</u> |
| 8.2 Explanations for individual characteristics..... | <u>15</u> |
| 8.3 Key for the stages of development..... | <u>23</u> |
| 8.4 Other names of the example varieties..... | <u>24</u> |
| 9. LITERATURE..... | <u>25</u> |
| 10. TECHNICAL QUESTIONNAIRE..... | <u>26</u> |

1. Subject of these Test Guidelines

These Test Guidelines apply to all varieties of *Brassica juncea* (L.) Czern..

2. Material Required

2.1 The competent authorities decide on the quantity and quality of the plant material required for testing the variety and when and where it is to be delivered. Applicants submitting material from a State other than that in which the testing takes place must ensure that all customs formalities and phytosanitary requirements are complied with.

2.2 The material is to be supplied in the form of seed.

2.3 The minimum quantity of plant material, to be supplied by the applicant, should be:

3,000 seeds for single spaced plants

or

20,000 seeds for drilled plots.

The seed should meet the minimum requirements for germination, species and analytical purity, health and moisture content, specified by the competent authority. In cases where the seed is to be stored, the germination capacity should be as high as possible and should, be stated by the applicant.

2.4 The plant material supplied should be visibly healthy, not lacking in vigor, nor affected by any important pest or disease.

2.5 The plant material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If it has been treated, full details of the treatment must be given.

3. Method of Examination

3.1 *Number of Growing Cycles*

3.1.1 The minimum duration of tests should normally be two independent growing cycles.

3.1.2 The testing of a variety may be concluded when the competent authority can determine with certainty the outcome of the test.

3.2 *Testing Place*

Tests are normally conducted at one place. In the case of tests conducted at more than one place, guidance is provided in TGP/9 "Examining Distinctness".

3.3 *Conditions for Conducting the Examination*

3.3.1 The tests should be carried out under conditions ensuring satisfactory growth for the expression of the relevant characteristics of the variety and for the conduct of the examination.

3.3.2 The optimum stage of development for the assessment of each characteristic is indicated by a number in the Table of Characteristics. The stages of development denoted by each number are described in Chapter 8.3.

3.4 *Test Design*

3.4.1 In the case of single spaced plants, each test should be designed to result in a total of at least 60 plants which should be divided between at least 2 replicates.

3.4.2 In the case of drilled plots, each test should be designed to result in a total of at least 200 plants which should be divided between at least 2 replicates.

3.5 *Additional Tests*

Additional tests, for examining relevant characteristics, may be established.

4. Assessment of Distinctness, Uniformity and Stability

4.1 *Distinctness*

4.1.1 General Recommendations

It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding distinctness. However, the following points are provided for elaboration or emphasis in these Test Guidelines.

4.1.2 Consistent Differences

The differences observed between varieties may be so clear that more than one growing cycle is not necessary. In addition, in some circumstances, the influence of the environment is not such that more than a single growing cycle is required to provide assurance that the differences observed between varieties are sufficiently consistent. One means of ensuring that a difference in a characteristic, observed in a growing trial, is sufficiently consistent is to examine the characteristic in at least two independent growing cycles.

4.1.3 Clear Differences

Determining whether a difference between two varieties is clear depends on many factors, and should consider, in particular, the type of expression of the characteristic being examined, i.e. whether it is expressed in a qualitative, quantitative, or pseudo-qualitative manner. Therefore, it is important that users of these Test Guidelines are familiar with the recommendations contained in the General Introduction prior to making decisions regarding distinctness.

4.1.4 Number of Plants or Parts of Plants to be Examined

Unless otherwise indicated, for the purposes of distinctness, all observations on single plants should be made on 30 plants or parts of plants taken from each of 30 plants and any other observations made on all plants in the test, disregarding any off-type plants.

4.1.5 Method of Observation

The recommended method of observing the characteristic for the purposes of distinctness is indicated by the following key in the Table of Characteristics (see document TGP/9 "Examining Distinctness", Section 4 "Observation of characteristics"):

MG: single measurement of a group of plants or parts of plants

MS: measurement of a number of individual plants or parts of plants

VG: visual assessment by a single observation of a group of plants or parts of plants

VS: visual assessment by observation of individual plants or parts of plants

Type of observation: visual (V) or measurement (M)

"Visual" observation (V) is an observation made on the basis of the expert's judgment. For the purposes of this document, "visual" observation refers to the sensory observations of the experts and, therefore, also includes smell, taste and touch. Visual observation includes observations where the expert uses reference points (e.g. diagrams, example varieties, side-by-side comparison) or non-linear charts (e.g. color charts). Measurement (M) is an objective observation against a calibrated, linear scale e.g. using a ruler, weighing scales, colorimeter, dates, counts, etc.

Type of record: for a group of plants (G) or for single, individual plants (S)

For the purposes of distinctness, observations may be recorded as a single record for a group of plants or parts of plants (G), or may be recorded as records for a number of single, individual plants or parts of plants (S). In most cases, "G" provides a single record per variety and it is not possible or necessary to apply statistical methods in a plant-by-plant analysis for the assessment of distinctness.

In cases where more than one method of observing the characteristic is indicated in the Table of Characteristics (e.g. VG/MG), guidance on selecting an appropriate method is provided in document TGP/9, Section 4.2.

4.2 *Uniformity*

- 4.2.1 It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding uniformity. However, the following points are provided for elaboration or emphasis in these Test Guidelines:
- 4.2.2 These Test Guidelines have been developed for the examination of seed-propagated varieties. For varieties with other types of propagation, the recommendations in the General Introduction and document TGP/13 "Guidance for new types and species" Section 4.5 "Testing Uniformity" should be followed.
- 4.2.3 The assessment of uniformity for cross-pollinated varieties should be according to the recommendations for cross-pollinated varieties in the General Introduction.
- 4.2.4 For the assessment of uniformity of seed-propagated varieties, a population standard of 2% and an acceptance probability of at least 95% should be applied. In the case of a sample size of 60 plants, 3 off-types are allowed. In the case of a sample size of 200 plants for drilled plants, 7 off-types are allowed.

4.3 *Stability*

- 4.3.1 In practice, it is not usual to perform tests of stability that produce results as certain as those of the testing of distinctness and uniformity. However, experience has demonstrated that, for many types of variety, when a variety has been shown to be uniform, it can also be considered to be stable.
- 4.3.2 Where appropriate, or in cases of doubt, stability may be further examined by testing a new seed stock to ensure that it exhibits the same characteristics as those shown by the initial material supplied.

5. Grouping of Varieties and Organization of the Growing Trial

- 5.1 The selection of varieties of common knowledge to be grown in the trial with the candidate varieties and the way in which these varieties are divided into groups to facilitate the assessment of distinctness are aided by the use of grouping characteristics.
- 5.2 Grouping characteristics are those in which the documented states of expression, even where produced at different locations, can be used, either individually or in combination with other such characteristics: (a) to select varieties of common knowledge that can be excluded from the growing trial used for examination of distinctness; and (b) to organize the growing trial so that similar varieties are grouped together.
- 5.3 The following have been agreed as useful grouping characteristics:
- (a) Seed: color (characteristic 1)
 - (b) Leaf: type (characteristic 5)
 - (c) Only varieties with leaf: type: entire or lobed: Leaf blade: density of incisions of margin (characteristic 18)
 - (d) Only varieties with leaf: type: entire or lobed: Leaf blade: blistering (characteristic 19)
 - (e) Only varieties with leaf: type: entire: Leaf blade: width of midrib (characteristic 20)
 - (f) Plant: head formation (characteristic 21)
- 5.4 Guidance for the use of grouping characteristics, in the process of examining distinctness, is provided through the General Introduction and document TGP/9 "Examining Distinctness".

6. Introduction to the Table of Characteristics

6.1 *Categories of Characteristics*

6.1.1 Standard Test Guidelines Characteristics

Standard Test Guidelines characteristics are those which are approved by UPOV for examination of DUS and from which members of the Union can select those suitable for their particular circumstances.

6.1.2 Asterisked Characteristics

Asterisked characteristics (denoted by *) are those included in the Test Guidelines which are important for the international harmonization of variety descriptions and should always be examined for DUS and included in the variety description by all members of the Union, except when the state of expression of a preceding characteristic or regional environmental conditions render this inappropriate.

6.2 *States of Expression and Corresponding Notes*

6.2.1 States of expression are given for each characteristic to define the characteristic and to harmonize descriptions. Each state of expression is allocated a corresponding numerical note for ease of recording of data and for the production and exchange of the description.

6.2.2 In the case of qualitative and pseudo-qualitative characteristics (see Chapter 6.3), all relevant states of expression are presented in the characteristic. However, in the case of quantitative characteristics with 5 or more states, an abbreviated scale may be used to minimize the size of the Table of Characteristics. For example, in the case of a quantitative characteristic with 9 states, the presentation of states of expression in the Test Guidelines may be abbreviated as follows:

| State | Note |
|--------|------|
| small | 3 |
| medium | 5 |
| large | 7 |

However, it should be noted that all of the following 9 states of expression exist to describe varieties and should be used as appropriate:

| State | Note |
|---------------------|------|
| very small | 1 |
| very small to small | 2 |
| small | 3 |
| small to medium | 4 |
| medium | 5 |
| medium to large | 6 |
| large | 7 |
| large to very large | 8 |
| very large | 9 |

6.2.3 Further explanation of the presentation of states of expression and notes is provided in document TGP/7 "Development of Test Guidelines".

6.3 *Types of Expression*

An explanation of the types of expression of characteristics (qualitative, quantitative and pseudo-qualitative) is provided in the General Introduction.

6.4 *Example Varieties*

Where appropriate, example varieties are provided to clarify the states of expression of each characteristic.

6.5 Legend

| | | English | français | deutsch | español | Example Varieties Exemples Beispielssorten Variedades ejemplo | Note/ Nota |
|---|---|------------------------------------|------------------------------|-------------------------------|--------------------------------|--|---------------|
| 1 | 2 | 3 | 4 | 5 | 6 | 7 | |
| | | Name of characteristics in English | Nom du caractère en français | Name des Merkmals auf Deutsch | Nombre del carácter en español | | |
| | | states of expression | types d'expression | Ausprägungsstufen | tipos de expresión | | |

- 1 Characteristic number
- 2 (*) Asterisked characteristic – see Chapter 6.1.2
- 3 Type of expression
 - QL Qualitative characteristic – see Chapter 6.3
 - QN Quantitative characteristic – see Chapter 6.3
 - PQ Pseudo-qualitative characteristic – see Chapter 6.3
- 4 Method of observation (and type of plot, if applicable)
 - MG, MS, VG, VS – see Chapter 4.1.5
- 5 (+) See Explanations on the Table of Characteristics in Chapter 8.2
- 6 (a)-(b) See Explanations on the Table of Characteristics in Chapter 8.1
- 7 Growth stage key See Explanations on the Table of Characteristics in Chapter 8.3

7. Table of Characteristics/Tableau des caractères/Merkmalstabelle/Tabla de caracteres

| | English | | français | | deutsch | español | Example Varieties Exemples Beispielssorten Variedades ejemplo | Note/ Nota |
|---------------|--|--------------|--|------------|------------------------------------|---|---|---------------|
| 1. (*) | QL | VG | | | 00 | | | |
| | Seed: color | | Graine : couleur | | Samen: Farbe | Semilla: color | | |
| | yellow | | jaune | | gelb | amarillo | Kigarashina | 1 |
| | blackish brown | | brun noirâtre | | schwärzlich braun | marrón negruzco | Akaoba Takana, Esperance, Miike Takana, Terrafit, Terraplus | 2 |
| 2. | QN | VG | | | 10 | | | |
| | Hypocotyl: anthocyanin coloration | | Hypocotyle : pigmentation anthocyanique | | Hypokotyl: Anthocyanfärbung | Hipocótilo: pigmentación antociánica | | |
| | absent or weak | | nulle ou faible | | fehlend oder gering | ausente o débil | Jarangi, TTK456, Zasai FM-58 | 1 |
| | medium | | moyenne | | mittel | media | Jarami, Shinkoku Seisai | 2 |
| | strong | | forte | | stark | fuerte | Kigarashina | 3 |
| 3. | QN | MS/VG | (+) | | 10 | | | |
| | Cotyledon: length | | Cotylédon : longueur | | Keimblatt: Länge | Cotiledón: longitud | | |
| | short | | court | | kurz | corto | Junkei Yamashiona, Vittasso | 3 |
| | medium | | moyen | | mittel | medio | Katsuona, Terraplus | 5 |
| | long | | long | | lang | largo | Scala | 7 |
| 4. | QN | MS/VG | (+) | | 10 | | | |
| | Cotyledon: width | | Cotylédon : largeur | | Keimblatt: Breite | Cotiledón: anchura | | |
| | narrow | | étroit | | schmal | estrecho | Junkei Yamashiona, Vittasso | 3 |
| | medium | | moyen | | mittel | medio | Katsuona, Pacific Gold, Terraplus | 5 |
| | broad | | large | | breit | ancho | Minaret, Terminator | 7 |
| 5. (*) | PQ | VG | (+) | (a) | 19 | | | |
| | Leaf: type | | Feuille : type | | Blatt: Typ | Hoja: tipo | | |
| | entire | | entière | | ganzrandig | entera | Akaoba Takana, Kekkyu Takana, Miike Takana, Sagami Green, Shinkoku Seisai | 1 |
| | lobed | | lobée | | gelappt | lobulada | Hagarashina, Kigarashina, Terrafit | 2 |
| | divided | | découpés | | geteilt | dividida | Akariasu, Flaming Frills, Riasu Karashina, Scarlet Frills | 3 |

| | English | | français | | deutsch | español | Example Varieties Exemples Beispielssorten Variedades ejemplo | Note/ Nota |
|----------------|--------------------------------|--------------------------------------|-------------------------------------|-----------------------------------|----------------------------|---------|--|---------------|
| 6. (*) | PQ | VG | (+) | (a) | 19 | | | |
| | Leaf: shape | Feuille : forme | Blatt: Form | Hoja: forma | | | | |
| | ovate | ovale | eiförmig | oval | Serihon | | 1 | |
| | circular | circulaire | rund | circular | Kekkyu Takana | | 2 | |
| | elliptic | elliptique | elliptisch | elíptica | Akariasu | | 3 | |
| | oblong | oblongue | länglich | oblonga | Etamine, Zasai FM-58 | | 4 | |
| | obovate | obovale | verkehrt eiförmig | oboval | Esperance, Katsuona | | 5 | |
| | spatulate | spatulée | spatelförmig | espatulada | Kigarashina | | 6 | |
| 7. (*) | QN | VG | (+) | (a) | 19 | | | |
| | Leaf: attitude | Feuille : port | Blatt: Haltung | Hoja: porte | | | | |
| | erect | dressée | aufrecht | erecta | Energy, Vittasso, Wasabina | | 1 | |
| | semi-erect | demi-dressée | halbaufrecht | semierecta | Esperance, Shinkoku Seisai | | 3 | |
| | horizontal | horizontale | waagrecht | horizontal | Etamine, Miike Takana | | 5 | |
| 8. | QN | MS/VG | (+) | (a) | 19 | | | |
| | Leaf: length | Feuille : longueur | Blatt: Länge | Hoja: longitud | | | | |
| | short | courte | kurz | corta | Chirimen Hakarashina | | 3 | |
| | medium | moyenne | mittel | media | Miike Takana, Terraplus | | 5 | |
| | long | longue | lang | larga | Akaoba Takana, Vittasso | | 7 | |
| 9. | QN | MS/VG | (+) | (a) | 19 | | | |
| | Leaf: width | Feuille : largeur | Blatt: Breite | Hoja: anchura | | | | |
| | narrow | étroite | schmal | estrecha | Chirimen Hakarashina | | 3 | |
| | medium | moyenne | mittel | media | Miike Takana, Terraplus | | 5 | |
| | broad | large | breit | ancha | Katsuona, Vittasso | | 7 | |
| 10. (*) | QN | MS/VG | (+) | (a) | 19 | | | |
| | Leaf: length of petiole | Feuille : longueur du pétiole | Blatt: Länge des Blattstiels | Hoja: longitud del pecíolo | | | | |
| | absent or very short | absent ou très court | fehlend oder sehr kurz | ausente o muy corto | Serihon | | 1 | |
| | short | court | kurz | corta | Miike Takana | | 3 | |
| | medium | moyen | mittel | media | Junkei Yamashiona | | 5 | |
| | long | long | lang | largo | | | 7 | |

| | English | | français | | deutsch | español | Example Varieties Exemples Beispielssorten Variedades ejemplo | Note/ Nota |
|---------|---|-------|--|-----|---|--|--|---------------|
| 11. | QN | MS/VG | (+) | (a) | 19 | | | |
| | Leaf: width of petiole | | Feuille : largeur du pétiole | | Blatt: Breite des Blattstiels | Hoja: anchura del pecíolo | | |
| | narrow | | étroit | | schmal | estrecho | Kigarashina | 3 |
| | medium | | moyen | | mittel | medio | Katsuona | 5 |
| | broad | | large | | breit | ancho | Shinkoku Seisai | 7 |
| 12. | QN | VG | (+) | (a) | 19 | | | |
| | <u>Only varieties with leaf: type: lobed or divided: Leaf blade: size of terminal lobe</u> | | <u>Seulement les variétés avec feuille : type : lobée ou découpés : Limbe : taille du lobe terminal</u> | | <u>Nur Sorten mit Blatt: Typ: gelappt oder geteilt: Blattspreite: Größe des Endlappens</u> | <u>Solo variedades con Hoja: tipo: lobulada o dividida: Limbo: tamaño del lóbulo terminal</u> | | |
| | small | | petit | | klein | pequeño | Akariasu | 3 |
| | medium | | moyen | | mittel | medio | Kigarashina | 5 |
| | large | | large | | groß | grande | Pacific Gold, Perm Green | 7 |
| 13. (*) | QN | VG | (+) | (a) | 19 | | | |
| | Leaf blade: number of lateral lobes | | Limbe : nombre de lobes latéraux | | Blattspreite: Anzahl der Seitenlappen | Limbo: número de lóbulos laterales | | |
| | absent or very few | | nul ou très petit | | fehlend oder sehr wenige | ausentes o muy bajo | Akaoba Takana, Sagami Green | 1 |
| | few | | petit | | wenige | bajo | Minaret | 3 |
| | medium | | moyen | | mittel | medio | Esperance, Kigarashina | 5 |
| | many | | grand | | viele | alto | Akariasu, TTK456 | 7 |
| 14. | QN | VG | | (a) | 19 | | | |
| | Leaf blade: pubescence on lower side | | Limbe : pubescence sur la face inférieure | | Blattspreite: Behaarung der Unterseite | Limbo: pubescencia en envés | | |
| | absent or weak | | nulle ou faible | | fehlend oder gering | ausente o débil | Miike Takana | 1 |
| | medium | | moyenne | | mittel | media | Oba Takana | 2 |
| | strong | | forte | | stark | densa | Kigarashina | 3 |
| 15. (*) | QN | VG | (+) | (a) | 19 | | | |
| | Leaf blade: anthocyanin coloration | | Limbe : pigmentation anthocyanique | | Blattspreite: Anthocyanfärbung | Limbo: pigmentación antocianica | | |
| | absent or very weak | | nulle ou très faible | | fehlend oder sehr gering | ausente o muy débil | Kekkyu Takana, Vitamine | 1 |
| | weak | | faible | | gering | débil | | 3 |
| | medium | | moyenne | | mittel | media | Miike Takana | 5 |
| | strong | | forte | | stark | fuerte | TTK456 | 7 |

| | English | | français | | deutsch | español | Example Varieties Exemples Beispielssorten Variedades ejemplo | Note/ Nota |
|---------|---|----|--|-----|--|---|--|---------------|
| 16. | QN | VG | (a) | | 19 | | | |
| | <u>Only varieties with anthocyanin</u> coloration: absent or very weak: Leaf blade: intensity of green color | | <u>Seulement les variétés avec pigmentation anthocyanique</u> : nulle ou très faible : Limbe : intensité de la couleur verte | | <u>Nur Sorten mit Anthocyanfärbung:</u> fehlend oder sehr gering: Blattspreite: Intensität der grünen Farbe | <u>Solo variedades con pigmentación antocianica:</u> ausente o muy débil: Limbo: intensidad del color verde | | |
| | light | | claire | | hell | claro | Wasabina | 3 |
| | medium | | moyenne | | mittel | medio | Etamine, Golden Streaks, Katsuona | 5 |
| | dark | | foncée | | dunkel | oscuro | Terratop | 7 |
| 17. | QN | VG | (a) | | 19 | | | |
| | <u>Only varieties with leaf: type: entire or lobed:</u> Leaf blade: undulation of margin | | <u>Seulement les variétés avec feuille : type : entière ou lobée :</u> Limbe : ondulation du bord | | <u>Nur Sorten mit Blatt: Typ: ganzrandig oder gelappt:</u> Blattspreite: Wellung des Randes | <u>Solo variedades con Hoja: tipo: entera o lobulada:</u> Limbo: ondulación del borde | | |
| | absent or very weak | | nulle ou très faible | | fehlend oder sehr gering | ausente o muy débil | | 1 |
| | weak | | faible | | gering | débil | Akaoba Takana | 2 |
| | medium | | moyenne | | mittel | media | Katsuona | 3 |
| | strong | | forte | | stark | fuerte | Chirimen Hakarashina | 4 |
| | very strong | | très forte | | sehr stark | muy fuerte | | 5 |
| 18. (*) | QN | VG | (+) | (a) | 19 | | | |
| | <u>Only varieties with leaf: type: entire or lobed:</u> Leaf blade: density of incisions of margin | | <u>Seulement les variétés avec feuille : type : entière ou lobée :</u> Limbe : densité des incisions du bord | | <u>Nur Sorten mit Blatt: Typ: ganzrandig oder gelappt:</u> Blattspreite: Dichte der Randeinschnitte | <u>Solo variedades con Hoja: tipo: entera o lobulada:</u> Limbo: densidad de las incisiones del borde | | |
| | absent or very sparse | | nulle ou très faible | | fehlend oder sehr locker | ausentes o muy laxas | | 1 |
| | sparse | | faible | | locker | laxas | Etamine, Katsuona | 3 |
| | medium | | moyenne | | mittel | medias | Opaleska | 5 |
| | dense | | forte | | dicht | densas | Oportuna | 7 |
| 19. (*) | QN | VG | (+) | (a) | 19 | | | |
| | <u>Only varieties with leaf: type: entire or lobed:</u> Leaf blade: blistering | | <u>Seulement les variétés avec feuille : type : entière ou lobée :</u> Limbe : cloqûre | | <u>Nur Sorten mit Blatt: Typ: ganzrandig oder gelappt:</u> Blattspreite: Blasigkeit | <u>Solo variedades con Hoja: tipo: entera o lobulada:</u> Limbo: ampollado | | |
| | absent or weak | | absente ou faible | | fehlend oder schwach | ausente o débil | Etamine, Kigarashina | 1 |
| | medium | | moyenne | | mittel | medio | Akaoba Takana | 2 |
| | strong | | forte | | stark | fuerte | Katsuona | 3 |

| | English | | français | | deutsch | español | Example Varieties Exemples Beispielssorten Variedades ejemplo | Note/ Nota |
|----------------|--|--------------|---|------------|--|--|--|---------------|
| 20. (*) | QN | MS/VG | (+) | (a) | 19 | | | |
| | Only varieties with leaf: type: entire: Leaf blade: width of midrib | | Seulement les variétés avec feuille : type : entière : Limbe : largeur de la nervure médiane | | Nur Sorten mit Blatt: Typ: ganzrandig: Blattspreite: Breite der Mittelrippe | Solo variedades con Hoja: tipo: entera: Limbo: anchura del nervio central | | |
| | narrow | | étroite | | schmal | estrecho | Sagami Green | 3 |
| | medium | | moyenne | | mittel | medio | Katsuona | 5 |
| | broad | | large | | breit | ancho | Shinkoku Seisai | 7 |
| 21. (*) | QL | VG | (+) | | 19 | | | |
| | Plant: head formation | | Plante : formation d'un capitule | | Pflanze: Kopfbildung | Planta: formación de repollo | | |
| | absent | | absente | | fehlend | ausente | Kigarashina | 1 |
| | present | | présente | | vorhanden | presente | Kekkyu Takana | 9 |
| 22. | QN | MS/VG | | | 19 | | | |
| | Head: height | | Capitule : hauteur | | Kopf: Höhe | Repollo: altura | | |
| | short | | bas | | kurz | corto | | 1 |
| | medium | | moyen | | mittel | medio | Kekkyu Takana, Unzen Kekkyu Takana | 2 |
| | tall | | haut | | hoch | alto | | 3 |
| 23. | QN | MS/VG | | | 19 | | | |
| | Head: width | | Capitule : largeur | | Kopf: Breite | Repollo: anchura | | |
| | narrow | | étroit | | schmal | estrecho | | 1 |
| | medium | | moyen | | mittel | medio | Kekkyu Takana, Unzen Kekkyu Takana | 2 |
| | broad | | large | | breit | ancho | | 3 |
| 24. | QN | MS/VG | | | 19 | | | |
| | Head: number of leaves | | Capitule : nombre de feuilles | | Kopf: Anzahl Blätter | Repollo: número de hojas | | |
| | few | | petit | | wenige | bajo | | 3 |
| | medium | | moyen | | mittel | medio | Kekkyu Takana, Unzen Kekkyu Takana | 5 |
| | many | | grand | | viele | alto | | 7 |
| 25. | PQ | VG | | | 19 | | | |
| | Head: internal color | | Capitule : couleur intérieure | | Kopf: Innenfarbe | Repollo: color interno | | |
| | yellowish white | | blanc jaunâtre | | gelblich weiß | blanco amarillento | Unzen Kekkyu Takana | 1 |
| | light green | | vert clair | | hellgrün | verde claro | | 2 |
| | medium green | | vert moyen | | mittelgrün | verde medio | Kekkyu Takana | 3 |

| | English | | français | deutsch | español | Example Varieties Exemples Beispielssorten Variedades ejemplo | Note/ Nota |
|------------|-------------------------------------|--------------|-------------------------------------|------------------------------------|---|--|---------------|
| 26. | PQ | VG | (+) | 20-29 | | | |
| | Main stem: shape | | Tige principale : forme | Haupttrieb: Form | Tallo principal: forma | | |
| | narrow conical | | conique étroite | schmal kegelförmig | cónico estrecho | Kigarashina | 1 |
| | rounded | | arrondie | abgerundet | redondeado | Umino | 2 |
| | broad conical | | conique large | breit kegelförmig | cónico ancho | Zasai FM-58 | 3 |
| | branched | | ramifiée | verzweigt | ramificado | FE-K226 | 4 |
| 27. | QN | MG | | 31 | | | |
| | Time of beginning of bolting | | Époque de début de montaison | Zeitpunkt des Schossbeginns | Época del comienzo de la floración | | |
| | early | | précoce | früh | temprana | Junkei Yamashiona, Scala | 3 |
| | medium | | moyenne | mittel | media | Terraplus | 5 |
| | late | | tardive | spät | tardía | Akaoba Takana | 7 |
| 28. | QN | MG/MS | | 50 | | | |
| | Time of flowering | | Époque de floraison | Zeitpunkt der Blüte | Época de floración | | |
| | early | | précoce | früh | temprana | Terrafit | 3 |
| | medium | | moyenne | mittel | media | Minaret, Terraplus | 5 |
| | late | | tardive | spät | tardía | Brons | 7 |
| | very late | | très tardive | sehr spät | muy tardía | Vitasso | 9 |
| 29. | QN | MS/VG | (b) | 70-79 | | | |
| | Plant: height | | Plante : hauteur | Pflanze: Höhe | Planta: altura | | |
| | short | | courte | kurz | baja | Pacific Gold, Terminator | 3 |
| | medium | | moyenne | mittel | media | Terraplus | 5 |
| | tall | | haute | hoch | alta | Minaret | 7 |
| | very tall | | très haute | sehr hoch | muy alta | Vitasso | 9 |
| 30. | QN | MS/VG | (+) | (b) | 70-79 | | |
| | Siliqua: length | | Siliqua : longueur | Schote: Länge | Silicua: longitud | | |
| | short | | courte | kurz | corta | Terraplus, Vitasso | 3 |
| | medium | | moyenne | mittel | media | Pacific Gold | 5 |
| | long | | longue | lang | larga | Minaret | 7 |

| | English | | français | | deutsch | español | Example Varieties Exemples Beispielssorten Variedades ejemplo | Note/ Nota |
|------------|--|---|--|---|---------------------------|---------|--|---------------|
| 31. | QN | MS/VG | (+) | (b) | 70-79 | | | |
| | Siliqua: length of beak | Siliqua : longueur du bec | Schote: Länge der Spitze | Silicua: longitud de la punta | | | | |
| | short | court | kurz | corta | Terraplus, Vittasso | | | 3 |
| | medium | moyen | mittel | media | Terrafit | | | 5 |
| | long | long | lang | larga | | | | 7 |
| 32. | QN | MS/VG | (+) | (b) | 70-79 | | | |
| | Siliqua: width | Siliqua : largeur | Schote: Breite | Silicua: anchura | | | | |
| | narrow | étroite | schmal | estrecha | Vittasso | | | 3 |
| | medium | moyenne | mittel | media | Energy, Terrafit | | | 5 |
| | broad | large | breit | ancha | Oba Takana | | | 7 |
| 33. | QN | MS/VG | (+) | (b) | 70-79 | | | |
| | Siliqua: length of peduncle | Siliqua : longueur du pédoncule | Schote: Länge des Stiels | Silicua: longitud del pedúnculo | | | | |
| | short | court | kurz | corto | Vittasso | | | 3 |
| | medium | moyen | mittel | medio | Energy | | | 5 |
| | long | long | lang | largo | Minaret | | | 7 |
| 34. | QN | VG | (+) | | | | | |
| | Tendency to form inflorescences | Tendance à former des inflorescences | Neigung zur Bildung von Blütenständen | Tendencia a formar inflorescencias | | | | |
| | absent or very weak | nulle ou très faible | fehlend oder sehr gering | ausente o muy débil | Brons, Vittasso | | | 1 |
| | weak | faible | gering | débil | | | | 3 |
| | medium | moyenne | mittel | media | Terraplus | | | 5 |
| | strong | forte | stark | fuerte | | | | 7 |
| | very strong | très forte | sehr stark | muy fuerte | Energy, Minaret, Terrafit | | | 9 |

8. Explanations on the Table of Characteristics

8.1 *Explanations covering several characteristics*


Characteristics containing the following key in the Table of Characteristics should be examined as indicated below:

- (a) Observations should be made on the largest fully developed leaf.
- (b) Observations should only be made on varieties without head formation.

8.2 *Explanations for individual characteristics*

Ad. 3: Cotyledon: length

The observations should be made on cotyledons of 30 seedlings. If the two cotyledons differ in size, the biggest one should be measured. The length is defined as distance between the inclination at top of the cotyledon and the point where the width of the petiole is about 4 mm. The width of the cotyledon should be measured at the widest point of the cotyledons.




a = Cotyledon: width (characteristic 4)
b = Cotyledon: length (characteristic 3)







Ad. 4: Cotyledon: width

See Ad. 3

Ad. 5: Leaf: type



Ad. 6: Leaf: shape

| | ← broadest part → | | |
|----------------|--|--|--|
| | below middle | at middle | above middle |
| relative width | | | |
| narrow | |  4 oblong |  6 spatulate |
| medium |  1 ovate |  3 elliptic |  5 obovate |
| broad | |  2 circular | |

Ad. 7: Leaf: attitude



1
erect




3
semi-erect



5
horizontal

Ad. 8: Leaf: length



- a = Leaf: width (characteristic 9)
- b = Leaf: length (characteristic 8)
- c = Leaf: length of petiole (characteristic 10)
- d = Leaf: width of petiole (characteristic 11)
- e = Leaf blade: width of midrib (characteristic 20)

Ad. 9: Leaf: width

See Ad. 8

Ad. 10: Leaf: length of petiole

See Ad. 8

Ad. 11: Leaf: width of petiole


See Ad. 8

Ad. 12: Only varieties with leaf: type: lobed or divided: Leaf blade: size of terminal lobe


Parts of the leaf blade are considered as lobes if their length is at least equivalent to the width of the leaf petiole at their point of attachment and if the upper notch of the blade has at least half the length of the lobe itself.

The terminal lobe is the top lobe of the leaf, which is the No. 1 lobe in the following figure.

The lateral lobes are the lobes excluding the terminal lobe (numbers 2, 3, 4, etc. in the following figures)




lobed



divided

The size of terminal lobe should be assessed by the ratio of the terminal lobe size/the leaf size. The terminal lobe size and the leaf size are the size of the area which was surrounded by each outline of them.



a = Terminal lobe size
b = Leaf size

Ad. 13: Leaf blade: number of lateral lobes


See Ad. 12

Ad. 15: Leaf blade: anthocyanin coloration

The strongest intensity of anthocyanin should be observed (not the extension).

Ad. 18: Only varieties with leaf: type: entire or lobed: Leaf blade: density of incisions of margin

Observations should be made on the distal part of the leaves.



Ad. 19: Only varieties with leaf: type: entire or lobed: Leaf blade: blistering



Ad. 20: Only varieties with leaf: type: entire: Leaf blade: width of midrib

See Ad. 8

The width of midrib should be measured at the widest point.

Ad. 21: Plant: head formation



1
absent



9
present

Ad. 26: Main stem: shape

Observations on the shape of the main stem should be made after removing the leaves, excluding lateral stems which are located at the base of main stem.



1
narrow conical



2
rounded



3
broad conical




4
branched

Ad. 30: Silique: length

Observations on the silique should be made on the middle third of the inflorescence of the main stem.

Observations should be made on the length of the silique from attachment of peduncle to top, excluding beak.



- a = Silique: width (characteristic 32)
- b = Silique: length of beak (characteristic 31)
- c = Silique: length (characteristic 30)
- d = Silique: length of peduncle (characteristic 33)

Ad. 31: Silique: length of beak

See Ad. 30

Ad. 32: Silique: width

See Ad. 30

Ad. 33: Silique: length of peduncle

See Ad. 30

Ad. 34: Tendency to form inflorescences

Observations should be made in the year of sowing under long day conditions.

The observation of the tendency to form inflorescence (proportion of plants below bud stage, in bud stage, in flowering stage, in stage of silique formation) should be made in autumn, when the development stagnates.

Alternatively the beginning of flowering may be observed in this trial; early flowering would mean strong tendency, late flowering would mean weak tendency.

8.3 Key for the stages of development

| Key | General description |
|----------|---|
| <u>0</u> | <u>Principal growth stage 0: Germination</u> |
| 01 | Beginning of seed imbibition |
| 03 | Seed imbibition complete |
| 05 | Radicle emerged from seed |
| 07 | Hypocotyl with cotyledons emerged from seed |
| 08 | Hypocotyl with cotyledons growing towards soil surface |
| 09 | Emergence: cotyledons emerge through soil surface |
| <u>1</u> | <u>Principal growth stage 1: Leaf development</u> |
| 10 | Cotyledons completely unfolded |
| 11 | First leaf unfolded |
| 12 | 2 leaves unfolded |
| 13 | 3 leaves unfolded |
| 14 | 4 leaves unfolded |
| 15 | 5 leaves unfolded |
| 16 | 6 leaves unfolded |
| 17 | 7 leaves unfolded |
| 18 | 8 leaves unfolded |
| 19 | 9 or more leaves unfolded |
| <u>2</u> | <u>Principal growth stage 2: Formation of side shoots</u> |
| 20 | No side shoots |
| 21 | First side shoot detectable |
| 22 | 2 side shoots detectable |
| 23 | 3 side shoots detectable |
| 24 | 4 side shoots detectable |
| 25 | 5 side shoots detectable |
| 26 | 6 side shoots detectable |
| 27 | 7 side shoots detectable |
| 28 | 8 side shoots detectable |
| 29 | 9 or more side shoots detectable |
| <u>3</u> | <u>Principal growth stage 3: Stem elongation</u> |
| 30 | no internodes("rosette") |
| 31 | 1 visibly extended internode |
| 32 | 2 visibly extended internodes |
| 33 | 3 visibly extended internodes |
| 34 | 4 visibly extended internodes |
| 35 | 5 visibly extended internodes |
| 36 | 6 visibly extended internodes |
| 37 | 7 visibly extended internodes |
| 38 | 8 visibly extended internodes |
| 39 | 9 or more visibly extended internodes |
| <u>4</u> | <u>Principal growth stage 4: Inflorescence emergence</u> |
| 40 | Flower buds present, still enclosed by leaves |
| 41 | Flower buds visible from above ("green bud") |
| 42 | Flower buds free, level with the youngest leaves |
| 43 | Flower buds raised above the youngest leaves |
| 45 | Individual flower buds (main inflorescence) visible but still closed |
| 47 | Individual flower buds (secondary inflorescence) visible but still closed |
| 49 | First petals visible, flower buds still closed by ("yellow bud") |
| <u>5</u> | <u>Principal growth stage 5: Opening of flowers</u> |
| 50 | First flowers open |
| 51 | 10% of flowers on main raceme open, main raceme elongating |
| 52 | 20% of flowers on main raceme open |
| 53 | 30% of flowers on main raceme open |
| 54 | 40% of flowers on main raceme open |
| 55 | Full flowering: 50% flowers on main raceme open, older petals falling |
| 57 | Flowering declining: majority of petals fallen |
| 59 | End of flowering |

| | |
|----------|---|
| <u>6</u> | <u>Principal growth stage 6: Development of silique</u> |
| 61 | 10% of siliques have reached final size |
| 62 | 20% of siliques have reached final size |
| 63 | 30% of siliques have reached final size |
| 64 | 40% of siliques have reached final size |
| 65 | 50% of siliques have reached final size |
| 66 | 60% of siliques have reached final size |
| 67 | 70% of siliques have reached final size |
| 68 | 80% of siliques have reached final size |
| 69 | Nearly all siliques have reached final size |
| <u>7</u> | <u>Principal growth stage 7: Ripening</u> |
| 70 | seed green, filling silique cavity |
| 71 | 10% of siliques ripe, seeds dark and hard |
| 72 | 20% of siliques ripe, seeds dark and hard |
| 73 | 30% of siliques ripe, seeds dark and hard |
| 74 | 40% of siliques ripe, seeds dark and hard |
| 75 | 50% of siliques ripe, seeds dark and hard |
| 76 | 60% of siliques ripe, seeds dark and hard |
| 77 | 70% of siliques ripe, seeds dark and hard |
| 78 | 80% of siliques ripe, seeds dark and hard |
| 79 | Fully ripe: nearly all siliques ripe, seeds dark and hard |
| <u>8</u> | <u>Principal growth stage 8: Senescence</u> |
| 87 | Plant dead and dry |
| 89 | Harvested product |

8.4 *Other names of the example varieties*

| | |
|------------------------------|--------------------------|
| TTK456 ⁽¹⁾ | Chaplin ⁽²⁾ |
| Akaoba Takana ⁽³⁾ | Red Giant ⁽⁴⁾ |

(1) official denomination registered under the law in Japan in 2011.

(2) official denomination of TTK456 registered under the law in European Union in 2014.

(3) commercial name in Japan.

(4) commercial name of Akaoba Takana in European Union.

9. Literature

Fujishiro, T., 1996: Breeding processes and characteristics of a newly bred leaf mustard (*Brassica Juncea* Coss.). Kanagawa, JP

Joy Larkcom, 1991: Oriental Vegetables (The Complete guide for Garden and Kitchen). London, GB, pp. 39 to pp. 45

Meier, U., 2001: Growth stages of mono-and dicotyledonous plants. BBCH Monograph Federal Biological Research Centre for Agriculture and Forestry

Ministry of Agriculture, Forestry & Fisheries of Japan., 1994: National Test Guideline for Karashina

Phillips, R., Rix, M., 1993: Vegetables (The Pan Garden Plants Series). pp. 44

Tsukamoto, Y., 1994: The Grand Dictionary of Horticulture Volume 1. The Shogakukan Ltd., Tokyo, JP, pp. 520 to pp. 522

Takasi A., 2004: Yasai-engei-daihyakka 17. Shadanhojin Nousan-gyoson-bunkakyokai. Tokyo, JP. pp. 169 to pp. 233

10. Technical Questionnaire

| | | |
|-------------------------|-----------------|-------------------|
| TECHNICAL QUESTIONNAIRE | Page {x} of {y} | Reference Number: |
|-------------------------|-----------------|-------------------|

| | |
|--|---|
| | Application date: (not to be filled in by the applicant) |
|--|---|

TECHNICAL QUESTIONNAIRE
to be completed in connection with an application for plant breeders' rights

1. Subject of the Technical Questionnaire

1.1 Botanical name

1.2 Common name

2. Applicant

Name

Address

Telephone No.

Fax No.

E-mail address

Breeder (if different from applicant)

3. Proposed denomination and breeder's reference

Proposed denomination (if available)

Breeder's reference

| | | |
|-------------------------|-----------------|-------------------|
| TECHNICAL QUESTIONNAIRE | Page {x} of {y} | Reference Number: |
|-------------------------|-----------------|-------------------|

#4. Information on the breeding scheme and propagation of the variety

4.1 Breeding scheme

Variety resulting from:

4.1.1 Crossing

(a) controlled cross []

(b) partially known cross []

(c) unknown cross []

4.1.2 Mutation []
(please state parent variety)

4.1.3 Discovery and development []
(please state where and when discovered and how developed)

4.1.4 Other []
(Please provide details)

4.2 Method of propagating the variety

4.2.1 Seed-propagated varieties

(a) Cross-pollination

[]

(b) Other (please provide details)

[]

4.2.2 Other
(Please provide details)

[]

| | | |
|-------------------------|-----------------|-------------------|
| TECHNICAL QUESTIONNAIRE | Page {x} of {y} | Reference Number: |
|-------------------------|-----------------|-------------------|

5. Characteristics of the variety to be indicated (the number in brackets refers to the corresponding characteristic in Test Guidelines; please mark the note which best corresponds).

| Characteristics | Example Varieties | Note |
|---|---|-------|
| 5.1 Seed: color (1) | | |
| yellow | Kigarashina | 1 [] |
| blackish brown | Akaoba Takana, Esperance, Miike Takana, Terrafit, Terraplus | 2 [] |
| 5.2 Leaf: type (5) | | |
| entire | Akaoba Takana, Kekkyu Takana, Miike Takana, Sagami Green, Shinkoku Seisai | 1 [] |
| lobed | Hagarashina, Kigarashina, Terrafit | 2 [] |
| divided | Akariasu, Flaming Frills, Riasu Karashina, Scarlet Frills | 3 [] |
| 5.3 Leaf: shape (6) | | |
| ovate | Serihon | 1 [] |
| circular | Kekkyu Takana | 2 [] |
| elliptic | Akariasu | 3 [] |
| oblong | Etamine, Zasai FM-58 | 4 [] |
| obovate | Esperance, Katsuona | 5 [] |
| spatulate | Kigarashina | 6 [] |
| 5.4 Leaf: attitude (7) | | |
| erect | Energy, Vittasso, Wasabina | 1 [] |
| erect to semi-erect | | 2 [] |
| semi-erect | Esperance, Shinkoku Seisai | 3 [] |
| semi-erect to horizontal | | 4 [] |
| horizontal | Etamine, Miike Takana | 5 [] |

| | | |
|-------------------------|-----------------|-------------------|
| TECHNICAL QUESTIONNAIRE | Page {x} of {y} | Reference Number: |
|-------------------------|-----------------|-------------------|

| Characteristics | Example Varieties | Note |
|--|----------------------|-------|
| 5.5 (18) <u>Only varieties with leaf: type: entire or lobed:</u> Leaf blade: density of incisions of margin | | |
| absent or very sparse | | 1 [] |
| very sparse to sparse | | 2 [] |
| sparse | Etamine, Katsuona | 3 [] |
| sparse to medium | | 4 [] |
| medium | Opaleska | 5 [] |
| medium to dense | | 6 [] |
| dense | Oportuna | 7 [] |
| dense to very dense | | 8 [] |
| very dense | | 9 [] |
| 5.6 (19) <u>Only varieties with leaf: type: entire or lobed:</u> Leaf blade: blistering | | |
| absent or weak | Etamine, Kigarashina | 1 [] |
| medium | Akaoba Takana | 2 [] |
| strong | Katsuona | 3 [] |
| 5.7 (20) <u>Only varieties with leaf: type: entire:</u> Leaf blade: width of midrib | | |
| very narrow | | 1 [] |
| very narrow to narrow | | 2 [] |
| narrow | Sagami Green | 3 [] |
| narrow to medium | | 4 [] |
| medium | Katsuona | 5 [] |
| medium to broad | | 6 [] |
| broad | Shinkoku Seisai | 7 [] |
| broad to very broad | | 8 [] |
| very broad | | 9 [] |
| 5.8 (21) <u>Plant: head formation</u> | | |
| absent | Kigarashina | 1 [] |
| present | Kekkyu Takana | 9 [] |

| | | |
|-------------------------|-----------------|-------------------|
| TECHNICAL QUESTIONNAIRE | Page {x} of {y} | Reference Number: |
|-------------------------|-----------------|-------------------|

6. Similar varieties and differences from these varieties

Please use the following table and box for comments to provide information on how your candidate variety differs from the variety (or varieties) which, to the best of your knowledge, is (or are) most similar. This information may help the examination authority to conduct its examination of distinctness in a more efficient way.

| Denomination(s) of variety(ies) similar to your candidate variety | Characteristic(s) in which your candidate variety differs from the similar variety(ies) | Describe the expression of the characteristic(s) for the similar variety(ies) | Describe the expression of the characteristic(s) for your candidate variety |
|---|---|--|--|
| <i>Example</i> | <i>Leaf: shape</i> | <i>ovate</i> | <i>oblong</i> |
| | | | |
| | | | |
| | | | |

Comments:

| | | |
|-------------------------|-----------------|-------------------|
| TECHNICAL QUESTIONNAIRE | Page {x} of {y} | Reference Number: |
|-------------------------|-----------------|-------------------|

#7. Additional information which may help in the examination of the variety

7.1 In addition to the information provided in sections 5 and 6, are there any additional characteristics which may help to distinguish the variety?

Yes No

(If yes, please provide details)

7.2 Are there any special conditions for growing the variety or conducting the examination?

Yes No

(If yes, please provide details)

7.3 Other information

Main use:

Vegetable

Oilseed

Condiment

Green manure

other

Authorities may allow certain of this information to be provided in a confidential section of the Technical Questionnaire.

| | | |
|-------------------------|-----------------|-------------------|
| TECHNICAL QUESTIONNAIRE | Page {x} of {y} | Reference Number: |
|-------------------------|-----------------|-------------------|

8. Authorization for release

(a) Does the variety require prior authorization for release under legislation concerning the protection of the environment, human and animal health?

Yes [] No []

(b) Has such authorization been obtained?

Yes [] No []

If the answer to (b) is yes, please attach a copy of the authorization.

9. Information on plant material to be examined or submitted for examination

9.1 The expression of a characteristic or several characteristics of a variety may be affected by factors, such as pests and disease, chemical treatment (e.g. growth retardants or pesticides), effects of tissue culture, different rootstocks, scions taken from different growth phases of a tree, etc.

9.2 The plant material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If the plant material has undergone such treatment, full details of the treatment must be given. In this respect, please indicate below, to the best of your knowledge, if the plant material to be examined has been subjected to:

| | | |
|---|---------|--------|
| (a) Microorganisms (e.g. virus, bacteria, phytoplasma) | Yes [] | No [] |
| (b) Chemical treatment (e.g. growth retardant, pesticide) | Yes [] | No [] |
| (c) Tissue culture | Yes [] | No [] |
| (d) Other factors | Yes [] | No [] |

Please provide details for where you have indicated "yes".

.....

10. I hereby declare that, to the best of my knowledge, the information provided in this form is correct:

Applicant's name

Signature Date

[End of document]