



TG/30/7(proj.3)
ORIGINAL: English
DATE: 2025-07-10

INTERNATIONAL UNION FOR THE PROTECTION OF NEW VARIETIES OF PLANTS

Geneva

DRAFT

BENT GRASS *

UPOV Code(s): AGROS_CAN; AGROS_CAP;
AGROS_GIG; AGROS_STO

Agrostis canina L.; *Agrostis capillaris* L.;
Agrostis gigantea Roth; *Agrostis stolonifera* L.

GUIDELINES

FOR THE CONDUCT OF TESTS

FOR DISTINCTNESS, UNIFORMITY AND STABILITY

prepared by an expert from the Netherlands (Kingdom of the)

to be considered by the

*Technical Committee at its sixty-first session,
to be held Geneva from 2025-10-20 to 2025-10-21*

Disclaimer: this document does not represent UPOV policies or guidance

Alternative Names:*

Botanical name	English	French	German	Spanish
<i>Agrostis canina</i> L.	Velvet Bent	Agrostis des chiens	Hundsstraußgras	Agróstide canina, Agróstide de perro, Agróstide perruna
<i>Agrostis capillaris</i> L.	Browntop, Common Bent	Agrostide commune, Agrostide fine, Agrostide ténue	Gemeines Straußgras, Rotes Straußgras	Agróstide común
<i>Agrostis gigantea</i> Roth	Black Bent, Red Top	Agrostide blanche, Agrostide géante	Fioringras, Weißes Straußgras	Agróstide blanca, Pastoquilla
<i>Agrostis stolonifera</i> L.	Creeping Bent, Spreading Bent	Agrostide blanche, Agrostide stolonifère	Flechtstraußgras, Weißes Straußgras	Agróstide estolonífera

The purpose of these guidelines ("Test Guidelines") is to elaborate the principles contained in the General Introduction (document TG/1/3), and its associated TGP documents, into detailed practical guidance for the harmonized examination of distinctness, uniformity and stability (DUS) and, in particular, to identify appropriate characteristics for the examination of DUS and production of harmonized variety descriptions.

ASSOCIATED DOCUMENTS

These Test Guidelines should be read in conjunction with the General Introduction and its associated TGP documents.

* These names were correct at the time of the introduction of these Test Guidelines but may be revised or updated. [Readers are advised to consult the UPOV Code, which can be found on the UPOV Website (www.upov.int), for the latest information.]

<u>TABLE OF CONTENTS</u>	<u>PAGE</u>
1. SUBJECT OF THESE TEST GUIDELINES	3
2. MATERIAL REQUIRED	3
3. METHOD OF EXAMINATION	3
3.1 NUMBER OF GROWING CYCLES	3
3.2 TESTING PLACE	3
3.3 CONDITIONS FOR CONDUCTING THE EXAMINATION	3
3.4 TEST DESIGN	4
3.5 ADDITIONAL TESTS.....	4
4. ASSESSMENT OF DISTINCTNESS, UNIFORMITY AND STABILITY	4
4.1 DISTINCTNESS	4
4.2 UNIFORMITY	5
4.3 STABILITY.....	5
5. GROUPING OF VARIETIES AND ORGANIZATION OF THE GROWING TRIAL	5
6. INTRODUCTION TO THE TABLE OF CHARACTERISTICS.....	6
6.1 CATEGORIES OF CHARACTERISTICS	6
6.2 STATES OF EXPRESSION AND CORRESPONDING NOTES	6
6.3 TYPES OF EXPRESSION	6
6.4 EXAMPLE VARIETIES	6
6.5 LEGEND	7
7. TABLE OF CHARACTERISTICS/TABLEAU DES CARACTERES/MERKMALSTABELLE/TABLA DE CARACTERES.....	8
8. EXPLANATIONS ON THE TABLE OF CHARACTERISTICS	14
8.1 EXPLANATIONS COVERING SEVERAL CHARACTERISTICS	14
8.2 EXPLANATIONS FOR INDIVIDUAL CHARACTERISTICS	14
8.3 GROWTH STAGES FOR GRASSES	17
9. LITERATURE.....	18
10. TECHNICAL QUESTIONNAIRE	19

1. Subject of these Test Guidelines

These Test Guidelines apply to all varieties of *Agrostis canina* L., *Agrostis capillaris* L., *Agrostis gigantea* Roth and *Agrostis stolonifera* L.

2. Material Required

2.1 The competent authorities decide on the quantity and quality of the plant material required for testing the variety and when and where it is to be delivered. Applicants submitting material from a State other than that in which the testing takes place must ensure that all customs formalities and phytosanitary requirements are complied with.

2.2 The material is to be supplied in the form of seeds.

2.3 The minimum quantity of plant material, to be supplied by the applicant, should be:

400 grams of seed.

The seed should meet the minimum requirements for germination, species and analytical purity, health and moisture content, specified by the competent authority.

2.4 The plant material supplied should be visibly healthy, not lacking in vigor, nor affected by any important pest or disease.

2.5 The plant material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If it has been treated, full details of the treatment must be given.

3. Method of Examination

3.1 *Number of Growing Cycles*

3.1.1 The minimum duration of tests should normally be two independent growing cycles.

3.1.2 The two independent growing cycles should be in the form of two separate plantings.

3.1.3 The testing of a variety may be concluded when the competent authority can determine with certainty the outcome of the test.

3.2 *Testing Place*

Tests are normally conducted at one place. In the case of tests conducted at more than one place, guidance is provided in TGP/9 "Examining Distinctness".

3.3 *Conditions for Conducting the Examination*

3.3.1 The tests should be carried out under conditions ensuring satisfactory growth for the expression of the relevant characteristics of the variety and for the conduct of the examination.

3.3.2 The optimum stage of development for the assessment of each characteristic is indicated by a number in the Table of Characteristics. The stages of development denoted by each number are described in Chapter 8.3.

3.3.3 The recommended type of plot in which to observe the characteristic is indicated by the following key in the Table of Characteristics:

A: spaced plants
B: row plot
C: special test

3.4 *Test Design*

3.4.1 Each test should be designed to result in at least 60 plants which should be divided between at least two replicates. In addition, the test may include 8 meters of row plot which should be divided between at least two replicates. The density of the seed should be such that around 200 plants/meter can be expected.

3.4.2 The design of the tests should be such that plants or parts of plants may be removed for measurement or counting without prejudice to the observations which must be made up to the end of the growing cycle.

3.5 *Additional Tests*

Additional tests, for examining relevant characteristics, may be established.

4. Assessment of Distinctness, Uniformity and Stability

4.1 *Distinctness*

4.1.1 General Recommendations

It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding distinctness. However, the following points are provided for elaboration or emphasis in these Test Guidelines.

4.1.2 Consistent Differences

The differences observed between varieties may be so clear that more than one growing cycle is not necessary. In addition, in some circumstances, the influence of the environment is not such that more than a single growing cycle is required to provide assurance that the differences observed between varieties are sufficiently consistent. One means of ensuring that a difference in a characteristic, observed in a growing trial, is sufficiently consistent is to examine the characteristic in at least two independent growing cycles.

4.1.3 Clear Differences

Determining whether a difference between two varieties is clear depends on many factors, and should consider, in particular, the type of expression of the characteristic being examined, i.e. whether it is expressed in a qualitative, quantitative, or pseudo-qualitative manner. Therefore, it is important that users of these Test Guidelines are familiar with the recommendations contained in the General Introduction prior to making decisions regarding distinctness.

4.1.4 Number of Plants or Parts of Plants to be Examined

Unless otherwise indicated, for the purposes of distinctness, all observations on single plants should be made on 60 plants or parts of plants taken from each of 60 plants and any other observations made on all plants in the test, disregarding any off-type plants.

In the case of observations of parts taken from single plants, the number of parts to be taken from each of the plants should be 1.

4.1.5 Method of Observation

The recommended method of observing the characteristic for the purposes of distinctness is indicated by the following key in the Table of Characteristics (see document TGP/9 "Examining Distinctness", Section 4 "Observation of characteristics"):

MG: single measurement of a group of plants or parts of plants

MS: measurement of a number of individual plants or parts of plants

VG: visual assessment by a single observation of a group of plants or parts of plants

VS: visual assessment by observation of individual plants or parts of plants"):

Type of observation: visual (V) or measurement (M)

"Visual" observation (V) is an observation made on the basis of the expert's judgment. For the purposes of this document, "visual" observation refers to the sensory observations of the experts and, therefore, also includes

smell, taste and touch. Visual observation includes observations where the expert uses reference points (e.g. diagrams, example varieties, side-by-side comparison) or non-linear charts (e.g. color charts). Measurement (M) is an objective observation against a calibrated, linear scale e.g. using a ruler, weighing scales, colorimeter, dates, counts, etc.

Type of record: for a group of plants (G) or for single, individual plants (S)

For the purposes of distinctness, observations may be recorded as a single record for a group of plants or parts of plants (G), or may be recorded as records for a number of single, individual plants or parts of plants (S). In most cases, "G" provides a single record per variety and it is not possible or necessary to apply statistical methods in a plant-by-plant analysis for the assessment of distinctness.

In cases where more than one method of observing the characteristic is indicated in the Table of Characteristics (e.g. VG/MG), guidance on selecting an appropriate method is provided in document TGP/9, Section 4.2.

4.2 *Uniformity*

4.2.1 It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding uniformity. However, the following points are provided for elaboration or emphasis in these Test Guidelines:

4.2.2 These Test Guidelines have been developed for the examination of cross-pollinated varieties. For varieties with other types of propagation, the recommendations in the General Introduction and document TGP/13 "Guidance for new types and species" Section 4.5 "Testing Uniformity" should be followed.

4.2.3 The assessment of uniformity should be according to the recommendations for cross-pollinated varieties in the General Introduction.

4.3 *Stability*

4.3.1 In practice, it is not usual to perform tests of stability that produce results as certain as those of the testing of distinctness and uniformity. However, experience has demonstrated that, for many types of variety, when a variety has been shown to be uniform, it can also be considered to be stable.

4.3.2 Where appropriate, or in cases of doubt, stability may be further examined by testing a new seed or plant stock to ensure that it exhibits the same characteristics as those shown by the initial material supplied.

5. Grouping of Varieties and Organization of the Growing Trial

5.1 The selection of varieties of common knowledge to be grown in the trial with the candidate varieties and the way in which these varieties are divided into groups to facilitate the assessment of distinctness are aided by the use of grouping characteristics.

5.2 Grouping characteristics are those in which the documented states of expression, even where produced at different locations, can be used, either individually or in combination with other such characteristics: (a) to select varieties of common knowledge that can be excluded from the growing trial used for examination of distinctness; and (b) to organize the growing trial so that similar varieties are grouped together.

5.3 The following have been agreed as useful grouping characteristics:

- (a) Ploidy (characteristic 1)
- (b) Plant: time of inflorescence emergence (characteristic 10)

5.4 Guidance for the use of grouping characteristics, in the process of examining distinctness, is provided through the General Introduction and document TGP/9 "Examining Distinctness".

6. Introduction to the Table of Characteristics

6.1 *Categories of Characteristics*

6.1.1 Standard Test Guidelines Characteristics

Standard Test Guidelines characteristics are those which are approved by UPOV for examination of DUS and from which members of the Union can select those suitable for their particular circumstances.

6.1.2 Asterisked Characteristics

Asterisked characteristics (denoted by *) are those included in the Test Guidelines which are important for the international harmonization of variety descriptions and should always be examined for DUS and included in the variety description by all members of the Union, except when the state of expression of a preceding characteristic or regional environmental conditions render this inappropriate.

6.2 *States of Expression and Corresponding Notes*

6.2.1 States of expression are given for each characteristic to define the characteristic and to harmonize descriptions. Each state of expression is allocated a corresponding numerical note for ease of recording of data and for the production and exchange of the description.

6.2.2 All relevant states of expression are presented in the characteristic.

6.2.3 Further explanation of the presentation of states of expression and notes is provided in document TGP/7 "Development of Test Guidelines".

6.3 *Types of Expression*

An explanation of the types of expression of characteristics (qualitative, quantitative and pseudo-qualitative) is provided in the General Introduction.

6.4 *Example Varieties*

Where appropriate, example varieties are provided to clarify the states of expression of each characteristic.

The species of the example varieties are indicated as follows:

(As): *Agrostis stolonifera* L.
(Acap): *Agrostis capillaris* L.
(Ac): *Agrostis canina* L.
(Ag): *Agrostis gigantea* Roth

6.5 Legend

		English		français		deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
1	2	3	4	5	6	7			
		Name of characteristics in English		Nom du caractère en français		Name des Merkmals auf Deutsch	Nombre del carácter en español		
		states of expression		types d'expression		Ausprägungsstufen	tipos de expresión		

- 1 Characteristic number
- 2 (*) Asterisked characteristic – see Chapter 6.1.2
- 3 Type of expression
 QL Qualitative characteristic – see Chapter 6.3
 QN Quantitative characteristic – see Chapter 6.3
 PQ Pseudo-qualitative characteristic – see Chapter 6.3
- 4 Method of observation (and type of plot, if applicable)
 MG, MS, VG, VS – see Chapter 4.1.5
- 5 (+) See Explanations on the Table of Characteristics in Chapter 8.2
- 6 (a)-(b) See Explanations on the Table of Characteristics in Chapter 8.1
- 7 Growth stage key See Explanations on the Table of Characteristics in Chapter 8.3
- A, B, C See Chapter 3.3.3

7. Table of Characteristics/Tableau des caracteres/Merkmalstabelle/Tabla de caracteres

	English		français		deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
1. (*)	QL	MG C	(+)					
	Ploidy		Ploïdie		Ploidie	Ploidía		
	diploid		diploïde		diploid	diploide	Vesper (Ac)	2
	tetraploid		tétraploïde		tetraploid	tetraploide	Flagstick (As), Teetop (Acap)	4
	hexaploid		hexaploïde		hexaploid	hexaploide	Vaclav (Ag)	6
2. (*)	QN	VG B VS A	(a)		20-29			
	Plant: growth habit without vernalization		Plante : port <u>sans</u> vernalisation		Pflanze: Wuchsform <u>ohne</u> Vernalisation	Planta: hábito de crecimiento <u>sin</u> vernalización		
	erect		dressé		aufrecht	erecto		1
	erect to semi-erect		dressé à demi-dressé		aufrecht bis halbaufrecht	erecto a semierecto		2
	semi-erect		demi-dressé		halbaufrecht	semierecto		3
	semi-erect to intermediate		demi-dressé à intermédiaire		halbaufrecht bis mittel	semierecto a intermedio	Manor (Acap)	4
	intermediate		intermédiaire		mittel	intermedio	Puritan (Acap), Tyee (As)	5
	intermediate to semi- prostrate		intermédiaire à demi-étalé		mittel bis halbliegend	intermedio a semipostrado	Cobra Nova (As)	6
	semi-prostrate		demi-étalé		halbliegend	semipostrado	Jorvik (Acap)	7
	semi-prostrate to prostrate		demi-étalé à étalé		halbliegend bis liegend	semipostrado a postrado	Flagstick (As)	8
	prostrate		étalé		liegend	postrado		9
3.	QN	MS A VG B	(+)		20-29			
	Plant: natural height without vernalization		Plante : hauteur naturelle <u>sans</u> vernalisation		Pflanze: natürliche Höhe <u>ohne</u> Vernalisation	Planta: altura natural <u>sin</u> vernalización		
	very short		très basse		sehr niedrig	muy corta		1
	short		basse		niedrig	corta	Flagstick (As), Teetop (Acap)	2
	medium		moyenne		mittel	media	777 Triple Seven (As)	3
	tall		haute		hoch	alta	Manor (Acap), PC2 (As)	4
	very tall		très haute		sehr hoch	muy alta		5

	English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
4. (*)	QN	VG B VS A			20-29	
	Leaf: intensity of green color <u>without</u> vernalization	Feuille : intensité de la couleur verte <u>sans</u> vernalisation	Blatt: Intensität der Grünfärbung <u>ohne</u> Vernalisation	Hoja: intensidad del color verde <u>sin</u> vernalización		
	very light	très claire	sehr hell	muy clara		1
	very light to light	très claire à claire	sehr hell bis hell	muy clara a clara		2
	light	claire	hell	clara		3
	light to medium	claire à moyenne	hell bis mittel	clara a media	Manor (Acap)	4
	medium	moyenne	mittel	media	Match Play (As)	5
	medium to dark	moyenne à foncée	mittel bis dunkel	media a oscura	Puritan (Acap)	6
	dark	foncée	dunkel	oscura	Charles (Acap), Piranha (As)	7
	dark to very dark	foncée à très foncée	dunkel bis sehr dunkel	oscura a muy oscura		8
	very dark	très foncée	sehr dunkel	muy oscura		9
5. (*)	QN	MS A VG B VS A			20-29	
	Leaf: width <u>without</u> vernalization	Feuille : largeur <u>sans</u> vernalisation	Blatt: Breite <u>ohne</u> Vernalisation	Hoja: anchura <u>sin</u> vernalización		
	very narrow	très étroite	sehr schmal	muy estrecha		1
	narrow	étroite	schmal	estrecha	Arrowtown (Acap)	2
	medium	moyenne	mittel	media	Barking (Acap), Tyee (As)	3
	broad	large	breit	ancha	Macdonald (As), Manor (Acap)	4
	very broad	très large	sehr breit	muy ancha		5
6.	QN	VG B VS A	(+)			
	Plant: tendency to form inflorescences <u>without</u> vernalization	Plante : tendance à former des inflorescences <u>sans</u> vernalisation	Pflanze: Neigung zur Bildung von Blütenständen <u>ohne</u> Vernalisation	Planta: tendencia a formar inflorescencias <u>sin</u> vernalización		
	absent or very weak	absente ou très faible	fehlend oder sehr schwach	ausente o muy débil	Leikvin (Acap)	1
	very weak to weak	très faible à faible	sehr schwach bis schwach	muy débil a débil		2
	weak	faible	schwach	débil		3
	weak to medium	faible à moyenne	schwach bis mittel	débil a media	Sztar (As)	4
	medium	moyenne	mittel	media	Arrowtown (Acap)	5
	medium to strong	moyenne à forte	mittel bis stark	media a fuerte		6
	strong	fort	stark	fuerte		7
	strong to very strong	forte à très forte	stark bis sehr stark	fuerte a muy fuerte	Grasslands Sefton (Acap)	8
	very strong	très forte	sehr stark	muy fuerte		9

	English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
7.	QN	VG B VS A	(a)	30-39		
	Plant: growth habit <u>after</u> vernalization	Plante : port <u>après</u> vernalisation	Pflanze: Wuchsform <u>nach</u> der Vernalisation	Planta: hábito de crecimiento <u>después</u> de la vernalización		
	erect	dressé	aufrecht	erecto		1
	erect to semi-erect	dressé à demi-dressé	aufrecht bis halbaufrecht	erecto a semierecto		2
	semi-erect	demi-dressé	halbaufrecht	semierecto		3
	semi-erect to intermediate	demi-dressé à intermédiaire	halbaufrecht bis mittel	semierecto a intermedio	Heriot (Acap)	4
	intermediate	intermédiaire	mittel	intermedio	Gudrun (Acap)	5
	intermediate to semi-prostrate	intermédiaire à demi-étalé	mittel bis halbliegend	intermedio a semipostrado	Rhinegold (Acap)	6
	semi-prostrate	demi-étalé	halbliegend	semipostrado	Jorvik (Acap)	7
	semi-prostrate to prostrate	demi-étalé à étalé	halbliegend bis liegend	semipostrado a postrado		8
	prostrate	étalé	liegend	postrado		9
8.	QN	MS A VG B		30-39		
	Plant: natural height <u>after</u> vernalization	Plante : hauteur naturelle <u>après</u> vernalisation	Pflanze: natürliche Höhe <u>nach</u> der Vernalisation	Planta: altura natural <u>después</u> de la vernalización		
	very short	très basse	sehr niedrig	muy baja		1
	short	basse	niedrig	baja	Heriot (Acap)	2
	medium	moyenne	mittel	media	Tyee (As)	3
	tall	haute	hoch	alta	Grasslands Sefton (Acap)	4
	very tall	très haute	sehr hoch	muy alta		5
9.	QN	VG B VS A		30-39		
	Leaf: intensity of green color <u>after</u> vernalization	Feuille : intensité de la couleur verte <u>après</u> vernalisation	Blatt: Intensität der Grünfärbung <u>nach</u> der Vernalisation	Hoja: intensidad del color verde <u>después</u> de la vernalización		
	very light	très claire	sehr hell	muy clara		1
	very light to light	très claire à claire	sehr hell bis hell	muy clara a clara		2
	light	claire	hell	clara		3
	light to medium	claire à moyenne	hell bis mittel	clara a media	Manor (Acap)	4
	medium	moyenne	mittel	media	Musket (Acap)	5
	medium to dark	moyenne à foncée	mittel bis dunkel	media a oscura		6
	dark	foncée	dunkel	oscura	Heriot (Acap)	7
	dark to very dark	foncée à très foncée	dunkel bis sehr dunkel	oscura a muy oscura		8
	very dark	très foncée	sehr dunkel	muy oscura		9

	English		français		deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
10. (*)	QN	MG B/MS A	(+)					
	Plant: time of inflorescence emergence	Plante : époque d'épiaison	Pflanze: Zeitpunkt des Erscheinens der Blütenstände	Planta: época de emergencia de las inflorescencias				
	very early	très précoce	sehr früh	muy temprana				1
	very early to early	très précoce à précoce	sehr früh bis früh	muy temprana a temprana				2
	early	précoce	früh	temprana	Gudrun (Acap), Highland (Acap), Kingstown (Ac)			3
	early to medium	précoce à moyenne	früh bis mittel	temprana a media	Puritan (Acap)			4
	medium	moyenne	mittel	media	Galina (Acap), Macdonald (As)			5
	medium to late	moyenne à tardive	mittel bis spät	media a tardía	Barking (Acap)			6
	late	tardive	spät	tardía	Cobra Nova (As)			7
	late to very late	tardive à très tardive	spät bis sehr spät	tardía a muy tardía				8
	very late	très tardive	sehr spät	muy tardía				9
11. (*)	QN	MS A	(+)	(b)	50-56			
	Flag leaf: length	Dernière feuille : longueur	Fahnenblatt: Länge	Hoja bandera: longitud				
	very short	très courte	sehr kurz	muy corta				1
	very short to short	très courte à courte	sehr kurz bis kurz	muy corta a corta	Tyee (As)			2
	short	courte	kurz	corta	Regent (As)			3
	short to medium	courte à moyenne	kurz bis mittel	corta a media	Rhinegold (Acap)			4
	medium	moyenne	mittel	media	Teetop (Acap)			5
	medium to long	moyenne à longue	mittel bis lang	media a larga	Gudrun (Acap)			6
	long	longue	lang	larga				7
	long to very long	longue à très longue	lang bis sehr lang	larga a muy larga				8
	very long	très longue	sehr lang	muy larga				9
12. (*)	QN	MS A	(+)	(b)	50-56			
	Flag leaf: width	Dernière feuille : largeur	Fahnenblatt: Breite	Hoja bandera: anchura				
	very narrow	très étroite	sehr schmal	muy estrecha				1
	very narrow to narrow	très étroite à étroite	sehr schmal bis schmal	muy estrecha a estrecha				2
	narrow	étroite	schmal	estrecha	Regent (As), Teetop (Acap), Tyee (As)			3
	narrow to medium	étroite à moyenne	schmal bis mittel	estrecha a media	Rhinegold (Acap)			4
	medium	moyenne	mittel	media				5
	medium to broad	moyenne à large	mittel bis breit	media a ancha	Gudrun (Acap)			6
	broad	large	breit	ancha				7
	broad to very broad	large à très large	breit bis sehr breit	ancha muy ancha				8
	very broad	très large	sehr breit	muy ancha				9

	English		français		deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
13.	QN	MS A			50-56			
	Flag leaf: length/width ratio	Dernière feuille : rapport longueur/largeur		Fahnenblatt: Verhältnis Länge/Breite		Hoja bandera: relación longitud/anchura		
	very low	très bas		sehr klein		muy baja		1
	very low to low	très bas à bas		sehr klein bis klein		muy baja a baja		Tyee (As) 2
	low	bas		klein		baja		Regent (As), Rhinegold (Acap) 3
	low to medium	bas à moyen		klein bis mittel		baja a media		Gudrun (Acap) 4
	medium	moyen		mittel		media		Teetop (Acap) 5
	medium to high	moyen à élevé		mittel bis groß		media a alta		6
	high	élevé		groß		alta		7
	high to very high	élevé à très élevé		groß bis sehr groß		alta a muy alta		8
	very high	très élevé		sehr groß		muy alta		9
14. (*)	QN	MS A	(+)	(b)	60-68			
	Stem: length	Tige : longueur		Halm: Länge		Tallo: longitud		
	very short	très courte		sehr kurz		muy corta		1
	very short to short	très courte à courte		sehr kurz bis kurz		muy corta a corta		2
	short	courte		kurz		corta		Saulsbury (Acap) 3
	short to medium	courte à moyenne		kurz bis mittel		corta a media		Howden (Acap), 777 Triple Seven (As) 4
	medium	moyenne		mittel		media		Greenspeed (Acap), Kingstown (Ac) 5
	medium to long	moyenne à longue		mittel bis lang		media a larga		Cobra Nova (As), Gudrun (Acap) 6
	long	longue		lang		larga		7
	long to very long	longue à très longue		lang bis sehr lang		larga a muy larga		8
	very long	très longue		sehr lang		muy larga		Kita (Ag) 9
15.	QN	MS A	(+)	(b)	60-68			
	Stem: length of upper internode	Tige : longueur du dernier entrenœud		Halm: Länge des obersten Internodiums		Tallo: longitud del entrenudo superior		
	very short	très courte		sehr kurz		muy corta		1
	very short to short	très courte à courte		sehr kurz bis kurz		muy corta a corta		2
	short	courte		kurz		corta		PC2 (As), Saulsbury (Acap) 3
	short to medium	courte à moyenne		kurz bis mittel		corta a media		4
	medium	moyenne		mittel		media		Independence (As), Red Mountain (Acap) 5
	medium to long	moyenne à longue		mittel bis lang		media a larga		Gudrun (Acap) 6
	long	longue		lang		larga		7
	long to very long	longue à très longue		lang bis sehr lang		larga a muy larga		8
	very long	très longue		sehr lang		muy larga		9


	English		français		deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
16.	QN	MS A	(+)	(b)	60-68			
	Inflorescence: length		Inflorescence : longueur		Blütenstand: Länge	Inflorescencia: longitud		
	very short		très courte		sehr kurz	muy corta		1
	very short to short		très courte à courte		sehr kurz bis kurz	muy corta a corta		2
	short		courte		kurz	corta	Kromi (As)	3
	short to medium		courte à moyenne		kurz bis mittel	corta a media	Puritan (Acap)	4
	medium		moyenne		mittel	media	Macdonald (As), Jorvik (Acap)	5
	medium to long		moyenne à longue		mittel bis lang	media a larga	Greenspeed (Acap)	6
	long		longue		lang	larga		7
	long to very long		longue à très longue		lang bis sehr lang	larga a muy larga		8
	very long		très longue		sehr lang	muy larga		9

8. Explanations on the Table of Characteristics

8.1 *Explanations covering several characteristics*

Characteristics containing the following key in the Table of Characteristics should be examined as indicated below:

(a) Observations should be made from the attitude of the leaves of the plant as a whole. The angle formed by the imaginary line through the region of greatest leaf density and the vertical should be used.



(b) Observations should be made on the longest stem when fully expanded.

8.2 *Explanations for individual characteristics*

Ad. 1: Ploidy

Observations should be made by standard cytological methods.

Ad. 3: Plant: natural height without vernalization

Observation should be made on the average height of the foliage in the centre of the plant.

Ad. 6: Plant: tendency to form inflorescences without vernalization

The number of plants showing at least three inflorescences should be recorded for each variety. Observations should be made at one occasion on the whole trial when the varieties are considered to have reached their full expression of this characteristic.

Ad. 10: Plant: time of inflorescence emergence

Spaced plants or row plots should be observed at least twice per week.

A: Plots with spaced plants

Time of inflorescence emergence is reached when the tip of three inflorescences can be seen protruding from the flag leaf sheath (Growth Stage DC 50).

B: Row plots

Time of inflorescence emergence is reached when the average plot stage is DC 54. This date should - if necessary- be obtained by interpolation. At each observation date, the average plot stage should be expressed in one of the following growth stages:

DC 50	First spikelet of inflorescence just visible
DC 52	25% of the inflorescence emerged (across all stems)
DC 54	50% of the inflorescence emerged (across all stems)
DC 56	75% of the inflorescence emerged (across all stems)

Ad. 11: Flag leaf: length

The flag leaf is the leaf directly below the inflorescence. Length and width should be observed on the same leaf.

Observations should be made from the tip of the leaf blade to the leaf sheath.


Ad. 12: Flag leaf: width

Observations should be made at the widest point of the leaf blade.

Ad. 14: Stem: length

Observations should be made on the longest stem from ground level to the tip of the inflorescence, when the inflorescence is fully expanded.

Ad. 15: Stem: length of upper internode



Char. 15: 1 = The part of the stem above the top node up to the beginning of the inflorescence is the upper internode.

Char. 16: 2 = Length of the inflorescence.

Ad. 16: Inflorescence: length

See Ad. 15

8.3 *Growth stages for grasses*

All characteristics should be recorded at the appropriate time for the plant concerned. Growth stages of grasses are indicated by decimal codes which are derived from the decimal code for the growth stages of cereals (Zadoks, et al., 1974). This decimal code is in close conformity with the BBCH-code (Meier, 1997).

Seedling growth (seedling: one shoot)

DC 10	First leaf through coleoptile
DC 15	Five leaves unfolded
DC 19	Nine or more leaves unfolded

Tillering

DC 20	Main shoot only (beginning of tillering)
DC 23	Main shoot and 3 tillers
DC 25	Main shoot and 5 tillers
DC 29	Main shoot and 9 or more tillers

Stem elongation

DC 30	Pseudo-stem erection (formed by sheaths of leaves).
DC 31	First node detectable (early stem extension across all stems)
DC 35	Fifth node detectable (50 % extension across all stems)
DC 39	Flag leaf ligula/collar just visible (pre-boot stage)

Booting

DC 41	Flag leaf sheath extending (little enlargement of the inflorescence, early boot-stage)
DC 45	Boots swollen (late-boot stage)
DC 47	Flag leaf sheath opening
DC 49	First awns visible (in awned forms only)

Inflorescence emergence (mostly non-synchronous)

DC 50	First spikelet of inflorescence just visible
DC 52	25 % of the inflorescence emerged (across all stems)
DC 54	50 % of the inflorescence emerged (across all stems)
DC 56	75 % of the inflorescence emerged (across all stems)
DC 58	Emergence of inflorescence completed

Anthesis (mostly non-synchronous)

DC 60	Beginning of anthesis
DC 64	Anthesis half-way
DC 68	Anthesis complete

9. Literature

Meier, U., 1997. Growth stages of mono- and dicotyledonous plants: BBCH-Monograph
Blackwell Science, Berlin, Vienna, a.o., pp 622.

Zadoks, J.C., T.T. Chang and C.F. Konzak, 1974. A decimal code for the growth stages of
cereals. Weed Research 14: 415 - 421.

10. Technical Questionnaire

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

	Application date: (not to be filled in by the applicant)
--	---

TECHNICAL QUESTIONNAIRE to be completed in connection with an application for plant breeders' rights	
1. Subject of the Technical Questionnaire 1.1.1 Botanical name Agrostis canina L. ☐ 1.1.2 Common name Velvet Bent 1.2.1 Botanical name Agrostis capillaris L. ☐ 1.2.2 Common name Browntop, Common Bent 1.3.1 Botanical name Agrostis gigantea Roth ☐ 1.3.2 Common name Black Bent, Red Top 1.4.1 Botanical name Agrostis stolonifera L. ☐ 1.4.2 Common name Creeping Bent, Spreading Bent	
2. Applicant Name Address Telephone No. Fax No. E-mail address Breeder (if different from applicant)	

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

3. Proposed denomination and breeder's reference

Proposed denomination
(if available)

Breeder's reference

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

#4. Information on the breeding scheme and propagation of the variety

4.1 Breeding scheme

Variety resulting from:

4.1.1 Crossing

(a) controlled cross []

(please state parent variety)

(.....) x (.....)

female parent

male parent

(b) partially known cross []

(please state parent variety(ies))

(.....) x (.....)

female parent

male parent

(c) unknown cross []

4.1.2 Mutation

(please state parent variety)

--

4.1.3 Discovery and development

(please state where and when discovered and how developed)

--

4.1.4 Other

(Please provide details)

--

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

4.2 Method of propagating the variety

4.2.1 Seed-propagated varieties

- (a) Cross-pollination []
 (i) Population []
 (ii) Synthetic variety []
(b) Other (please provide details) []

4.2.2 Other []
(Please provide details)

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

5. Characteristics of the variety to be indicated (the number in brackets refers to the corresponding characteristic in Test Guidelines; please mark the note which best corresponds).

	Characteristics	Example Varieties	Note
5.1 (1)	Ploidy		
	diploid	Vesper (Ac)	2 []
	tetraploid	Flagstick (As), Teetop (Acap)	4 []
	hexaploid	Vaclav (Ag)	6 []
5.2 (2)	Plant: growth habit <u>without</u> vernalization		
	erect		1 []
	erect to semi-erect		2 []
	semi-erect		3 []
	semi-erect to intermediate	Manor (Acap)	4 []
	intermediate	Puritan (Acap), Tyee (As)	5 []
	intermediate to semi-prostrate	Cobra Nova (As)	6 []
	semi-prostrate	Jorvik (Acap)	7 []
	semi-prostrate to prostrate	Flagstick (As)	8 []
	prostrate		9 []
5.3 (4)	Leaf: intensity of green color <u>without</u> vernalization		
	very light		1 []
	very light to light		2 []
	light		3 []
	light to medium	Manor (Acap)	4 []
	medium	Match Play (As)	5 []
	medium to dark	Puritan (Acap)	6 []
	dark	Charles (Acap), Piranha (As)	7 []
	dark to very dark		8 []
	very dark		9 []

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

	Characteristics	Example Varieties	Note
5.4 (10)	Plant: time of inflorescence emergence		
	very early		1 []
	very early to early		2 []
	early	Gudrun (Acap), Highland (Acap), Kingstown (Ac)	3 []
	early to medium	Puritan (Acap)	4 []
	medium	Galina (Acap), Macdonald (As)	5 []
	medium to late	Barking (Acap)	6 []
	late	Cobra Nova (As)	7 []
	late to very late		8 []
	very late		9 []
5.5 (12)	Flag leaf: width		
	very narrow		1 []
	very narrow to narrow		2 []
	narrow	Regent (As), Teetop (Acap), Tyee (As)	3 []
	narrow to medium	Rhinegold (Acap)	4 []
	medium		5 []
	medium to broad	Gudrun (Acap)	6 []
	broad		7 []
	broad to very broad		8 []
	very broad		9 []
5.6 (14)	Stem: length		
	very short		1 []
	very short to short		2 []
	short	Saulsbury (Acap)	3 []
	short to medium	777 Triple Seven (As), Howden (Acap)	4 []
	medium	Greenspeed (Acap), Kingstown (Ac)	5 []
	medium to long	Cobra Nova (As), Gudrun (Acap)	6 []
	long		7 []
	long to very long		8 []
	very long	Kita (Ag)	9 []

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

6. Similar varieties and differences from these varieties

Please use the following table and box for comments to provide information on how your candidate variety differs from the variety (or varieties) which, to the best of your knowledge, is (or are) most similar. This information may help the examination authority to conduct its examination of distinctness in a more efficient way.

Denomination(s) of variety(ies) similar to your candidate variety	Characteristic(s) in which your candidate variety differs from the similar variety(ies)	Describe the expression of the characteristic(s) for the similar variety(ies)	Describe the expression of the characteristic(s) for your candidate variety
<i>Example</i>	<i>Flag leaf: length</i>	<i>short</i>	<i>medium</i>

Comments

TECHNICAL QUESTIONNAIRE

Page {x} of {y}

Reference Number:

#7. Additional information which may help in the examination of the variety

7.1 In addition to the information provided in sections 5 and 6, are there any additional characteristics which may help to distinguish the variety?

Yes ☐ No ☐

(If yes, please provide details)

7.2 Are there any special conditions for growing the variety or conducting the examination?

Yes ☐ No ☐

(If yes, please provide details)

7.3 Other information

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
----------------------------	-----------------	-------------------

8. Authorization for release

(a) Does the variety require prior authorization for release under legislation concerning the protection of the environment, human and animal health?

Yes ☐ No ☐

(b) Has such authorization been obtained?

Yes ☐ No ☐

If the answer to (b) is yes, please attach a copy of the authorization.

9. Information on plant material to be examined or submitted for examination

9.1 The expression of a characteristic or several characteristics of a variety may be affected by factors, such as pests and disease, chemical treatment (e.g. growth retardants or pesticides), effects of tissue culture, different rootstocks, scions taken from different growth phases of a tree, etc.

9.2 The plant material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If the plant material has undergone such treatment, full details of the treatment must be given. In this respect, please indicate below, to the best of your knowledge, if the plant material to be examined has been subjected to:

(a) Microorganisms (e.g. virus, bacteria, phytoplasma) Yes ☐ No ☐

(b) Chemical treatment (e.g. growth retardant, pesticide) Yes ☐ No ☐

(c) Tissue culture Yes ☐ No ☐

(d) Other factors Yes ☐ No ☐

Please provide details for where you have indicated "yes".

9.3 Has the plant material to be examined been tested for the presence of virus or other pathogens?

Yes ☐

(please provide details as specified by the Authority)

No ☐

10. I hereby declare that, to the best of my knowledge, the information provided in this form is correct:

Applicant's name

Signature

Date

[End of document]