



TG/SWEETPOT(proj.5)
ORIGINAL: English
DATE: 2009-11-13

INTERNATIONAL UNION FOR THE PROTECTION OF NEW VARIETIES OF PLANTS
 GENEVA

DRAFT

SWEET POTATO

UPOV Code: IPOMO_BAT

Ipomoea batatas (L.) Lam.

*

GUIDELINES
FOR THE CONDUCT OF TESTS
FOR DISTINCTNESS, UNIFORMITY AND STABILITY

prepared by an expert from the Republic of Korea

*to be considered by the Enlarged Editorial Committee at its meeting
 to be held in Geneva, Switzerland, on January 7, 2010*

Alternative Names:*

<i>Botanical name</i>	<i>English</i>	<i>French</i>	<i>German</i>	<i>Spanish</i>
<i>Ipomoea batatas</i> (L.) Lam.	Sweet potato	Patate douce	Batate, Süßkartoffel	Camote, Batata

The purpose of these guidelines (“Test Guidelines”) is to elaborate the principles contained in the General Introduction (document TG/1/3), and its associated TGP documents, into detailed practical guidance for the harmonized examination of distinctness, uniformity and stability (DUS) and, in particular, to identify appropriate characteristics for the examination of DUS and production of harmonized variety descriptions.

ASSOCIATED DOCUMENTS

These Test Guidelines should be read in conjunction with the General Introduction and its associated TGP documents.

* These names were correct at the time of the introduction of these Test Guidelines but may be revised or updated. [Readers are advised to consult the UPOV Code, which can be found on the UPOV Website (www.upov.int), for the latest information.]

<u>TABLE OF CONTENTS</u>	<u>PAGE</u>
1. SUBJECT OF THESE TEST GUIDELINES	3
2. MATERIAL REQUIRED	3
3. METHOD OF EXAMINATION.....	3
3.1 Number of Growing Cycles	3
3.2 Testing Place	3
3.3 Conditions for Conducting the Examination.....	3
3.4 Test Design	4
3.5 Number of Plants / Parts of Plants to be Examined.....	4
3.6 Additional Tests	4
4. ASSESSMENT OF DISTINCTNESS, UNIFORMITY AND STABILITY.....	4
4.1 Distinctness	4
4.2 Uniformity.....	5
4.3 Stability	5
5. GROUPING OF VARIETIES AND ORGANIZATION OF THE GROWING TRIAL	5
6. INTRODUCTION TO THE TABLE OF CHARACTERISTICS	6
6.1 Categories of Characteristics.....	6
6.2 States of Expression and Corresponding Notes.....	6
6.3 Types of Expression	6
6.4 Example Varieties	6
6.5 Legend.....	6
7. TABLE OF CHARACTERISTICS/TABLEAU DES CARACTÈRES/MERKMALSTABELLE/TABLA DE CARACTERES.....	7
8. EXPLANATIONS ON THE TABLE OF CHARACTERISTICS	15
8.1 Explanations covering several characteristics	15
8.2. Explanations for individual characteristics	15
9. LITERATURE	19
10. TECHNICAL QUESTIONNAIRE.....	20

1. Subject of these Test Guidelines

These Test Guidelines apply to all varieties of *Ipomoea batatas* (L.) Lam. However, additional characteristics may be needed in order to examine ornamental varieties.

2. Material Required

2.1 The competent authorities decide on the quantity and quality of the plant material required for testing the variety and when and where it is to be delivered. Applicants submitting material from a State other than that in which the testing takes place must ensure that all customs formalities and phytosanitary requirements are complied with.

2.2 The material is to be supplied in the form of storage root of medium size of the variety or in the form of cutting.

2.3 The minimum quantity of plant material, to be supplied by the applicant, should be:

50 storage roots or 150 cuttings.

2.4 The plant material supplied should be visibly healthy, not lacking in vigor, nor affected by any important pest or disease.

2.5 The plant material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If it has been treated, full details of the treatment must be given.

3. Method of Examination

3.1 *Number of Growing Cycles*

The minimum duration of tests should normally be a single growing cycle.

3.2 *Testing Place*

Tests are normally conducted at one place. In the case of tests conducted at more than one place, guidance is provided in TGP/9 "Examining Distinctness."

3.3 *Conditions for Conducting the Examination*

3.3.1 The tests should be carried out under conditions ensuring satisfactory growth for the expression of the relevant characteristics of the variety and for the conduct of the examination.

3.3.2 Type of observation

The recommended method of observing the characteristic is indicated by the following key in the second column of the Table of Characteristics:

- MG: single measurement of a group of plants or parts of plants
MS: measurement of a number of individual plants or parts of plants
VG: visual assessment by a single observation of a group of plants or parts of plants
VS: visual assessment by observation of individual plants or parts of plants

3.4 Test Design

3.4.1 Each test should be designed to result in a total of at least 50 plants, which should be divided between at least two replicates.

3.4.2 The design of the tests should be such that plants or parts of plants may be removed for measurement or counting without prejudice to the observations which must be made up to the end of the growing cycle.

3.5 Number of Plants / Parts of Plants to be Examined

Unless otherwise indicated, all observations should be made on 30 plants or parts taken from each of 30 plants.

3.6 Additional Tests

Additional tests, for examining relevant characteristics, may be established.

4. Assessment of Distinctness, Uniformity and Stability

4.1 Distinctness

4.1.1 General Recommendations

It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding distinctness. However, the following points are provided for elaboration or emphasis in these Test Guidelines.

4.1.2 Consistent Differences

The differences observed between varieties may be so clear that more than one growing cycle is not necessary. In addition, in some circumstances, the influence of the environment is not such that more than a single growing cycle is required to provide assurance that the differences observed between varieties are sufficiently consistent. One means of ensuring that a difference in a characteristic, observed in a growing trial, is sufficiently consistent is to examine the characteristic in at least two independent growing cycles.

4.1.3 Clear Differences

Determining whether a difference between two varieties is clear depends on many factors, and should consider, in particular, the type of expression of the characteristic being examined, i.e. whether it is expressed in a qualitative, quantitative, or pseudo-qualitative manner. Therefore, it is important that users of these Test Guidelines are familiar with the recommendations contained in the General Introduction prior to making decisions regarding distinctness.

4.2 *Uniformity*

4.2.1 It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding uniformity. However, the following points are provided for elaboration or emphasis in these Test Guidelines:

4.2.2 For the assessment of uniformity, a population standard of 1% and an acceptance probability of at least 95% should be applied. In the case of a sample size of 50 plants, 2 off-types are allowed.

4.3 *Stability*

4.3.1 In practice, it is not usual to perform tests of stability that produce results as certain as those of the testing of distinctness and uniformity. However, experience has demonstrated that, for many types of variety, when a variety has been shown to be uniform, it can also be considered to be stable.

4.3.2 Where appropriate, or in cases of doubt, stability may be tested, either by growing a further generation, or by testing a new plant stock to ensure that it exhibits the same characteristics as those shown by the previous material supplied.

5. Grouping of Varieties and Organization of the Growing Trial

5.1 The selection of varieties of common knowledge to be grown in the trial with the candidate varieties and the way in which these varieties are divided into groups to facilitate the assessment of distinctness are aided by the use of grouping characteristics.

5.2 Grouping characteristics are those in which the documented states of expression, even where produced at different locations, can be used, either individually or in combination with other such characteristics: (a) to select varieties of common knowledge that can be excluded from the growing trial used for examination of distinctness; and (b) to organize the growing trial so that similar varieties are grouped together.

5.3 The following have been agreed as useful grouping characteristics:

- (a) Plant: growth habit (characteristic 1)
- (b) Stem: anthocyanin coloration of tip (characteristic 6)
- (c) Leaf blade: lobes (characteristic 9)
- (d) Storage root: shape (characteristic 19)
- (e) Storage root: main color of skin (characteristic 22)
- (f) Storage root: main color of flesh (characteristic 24)

5.4 Guidance for the use of grouping characteristics, in the process of examining distinctness, is provided through the General Introduction.

6. Introduction to the Table of Characteristics

6.1 *Categories of Characteristics*

6.1.1 Standard Test Guidelines Characteristics

Standard Test Guidelines characteristics are those which are approved by UPOV for examination of DUS and from which members of the Union can select those suitable for their particular circumstances.

6.1.2 Asterisked Characteristics

Asterisked characteristics (denoted by *) are those included in the Test Guidelines which are important for the international harmonization of variety descriptions and should always be examined for DUS and included in the variety description by all members of the Union, except when the state of expression of a preceding characteristic or regional environmental conditions render this inappropriate.

6.2 *States of Expression and Corresponding Notes*

States of expression are given for each characteristic to define the characteristic and to harmonize descriptions. Each state of expression is allocated a corresponding numerical note for ease of recording of data and for the production and exchange of the description.

6.3 *Types of Expression*

An explanation of the types of expression of characteristics (qualitative, quantitative and pseudo-qualitative) is provided in the General Introduction.

6.4 *Example Varieties*

Where appropriate, example varieties are provided to clarify the states of expression of each characteristic.

6.5 *Legend*

(*) Asterisked characteristic – see Chapter 6.1.2

QL: Qualitative characteristic – see Chapter 6.3

QN: Quantitative characteristic – see Chapter 6.3

PQ: Pseudo-qualitative characteristic – see Chapter 6.3

MG, MS, VG, VS: See Chapter 3.3.2

(a)-(e) See Explanations on the Table of Characteristics in Chapter 8.1

(+) See Explanations on the Table of Characteristics in Chapter 8.2

7. Table of Characteristics/Tableau des caractères/Merkmalstabelle/Tabla de caracteresticas

English	français	deutsch	español	Example Varieties/ Exemples/ Beispielssorten/ Variedades ejemplo	Note/ Nota
1. VG Plant: growth habit	Plante : port		Pflanze: Wuchsform	Planta: porte	
(*)					
QN (a) upright	dressé	aufrecht	erecto	Sinchunmi	1
semi-upright	demi-dressé	halbaufrecht	semierecto	Younmi	3
spreading	étalé	breitwüchsig	extendido	Yulmi	5
2. MS/ Stem: length of VG primary shoots	Tige : longueur des rameaux primaires		Stiel: Länge der primären Triebe erster Ordnung	Tallo: longitud de los brotes principales	
QN (a) short	courte	kurz	cortos	Sinchunmi	3
(b) medium	moyenne	mittel	medianos	Koganesengan, Younmi	5
long	longue	lang	largos	Zami	7
3. MS/ Stem: length of VG internode	Tige : longueur de l'entre-nœud		Stiel: Länge des Internodiums	Tallo: longitud del intranudo	
QN (a) short	court	kurz	corto	Younmi	3
(c) medium	moyen	mittel	mediano	Koganesengan, Yulmi	5
long	long	lang	largo	Shinhwangmi	7
4. MS/ Stem: diameter of VG internode	Tige : diamètre de l'entre-nœud		Stiel: Durchmesser des Internodiums	Tallo: diámetro del intranudo	
QN (a) very small	très petit	sehr klein	muy pequeño	Zami	1
(c) small	petit	klein	pequeño	Sinchunmi	3
medium	moyen	mittel	mediano	Koganesengan, Yulmi	5
large	grand	groß	grande	Shinyulmi	7
very large	très grand	sehr groß	muy grande	Chinmi	9
5. VG Stem: anthocyanin coloration of internode	Tige : pigmentation anthocyane de l'entre-nœud		Stiel: Anthocyanfärbung des Internodiums	Tallo: pigmentación antociánica del intranudo	
QN (a) absent or weak	absente ou faible	fehlend oder gering	ausente o débil	Yulmi	1
(b) medium	moyenne	mittel	media	Singeonmi	2
strong	forte	stark	fuerte	Hayanmi	3

					Example Varieties/ Exemples/ Beispielssorten/ Variedades ejemplo	Note/ Nota
English	français	deutsch	español			
6. (*)	VG Stem: anthocyanin coloration of tip	Tige : pigmentation anthocyane du sommet	Stiel: Anthocyanfärbung der Spitze	Tallo: pigmentación antociánica del extremo		
QN	(a) absent or weak	absente ou faible	fehlend oder gering	ausente o débil	Yulmi	1
	(b) medium	moyenne	mittel	media	Sinjami	2
	strong	forte	stark	fuerte	Hayanmi	3
7.	VG Stem: anthocyanin coloration of node	Tige : pigmentation anthocyane du nœud	Stiel: Anthocyanfärbung des Knotens	Tallo: pigmentación antociánica del nudo		
QN	(a) absent or weak	absente ou faible	fehlend oder gering	ausente o débil	Yulmi	1
	(b) medium	moyenne	mittel	media	Norin 2	2
	strong	forte	stark	fuerte	Hayanmi, Koganesengan	3
8. (*)	VG Stem: pubescence of tip	Tige : pilosité du sommet	Stiel: Behaarung der Spitze	Tallo: pubescencia del extremo		
QN	(a) absent or sparse	absente ou faible	fehlend oder locker	ausente o laxa	Yulmi	1
	(b) medium	moyenne	mittel	media	Koganesengan	2
	dense	forte	dicht	densa	Zami	3
9. (*) (+)	VG Leaf blade: lobes	Limbe : lobes	Blattspreite: Lappen	Limbo: lóbulos		
QL	(a) absent	absents	fehlend	ausente	Gokokuimo	1
	three lobes	trois lobes	drei Lappen	tres lóbulos	Benisengan	2
	five lobes	cinq lobes	fünf Lappen	cinco lóbulos	Koganesengan, Sinchunmi	3
	seven lobes	sept lobes	sieben Lappen	siete lóbulos	Benihayato	4

		English	français	deutsch	español	Example Varieties/ Exemples/ Beispielssorten/ Variedades ejemplos	Note/ Nota
10.	VG (*) (+)	Only varieties with leaf blade lobes: absent: Leaf blade: shape	Seulement variétés avec limbes ne comportant pas de lobes : Limbe : forme	Nur Sorten mit Blattspreitenlappen: fehlend: Blattspreite: Form	Sólo variedades que no presentan lóbulos en la hoja: Limbo: forma		
PQ	(a)	cordate	cordiforme	herzförmig	cordiforme	Gokokuimo, Yulmi	1
	(d)	triangular	triangulaire	dreieckig	triangular	Beniotome	2
		reniform	réniforme	nierenförmig	reniforme	Kohkei 14	3
		circular	circulaire	kreisförmig	circular		4
11.	VG (+)	Only varieties with leaf lobes present: Leaf blade: depth of lobing	Seulement variétés avec limbes comportant des lobes : Limbe : profondeur de découpage des bords	Nur Sorten mit Blattspreitenlappen: vorhanden: Blattspreite: Tiefe der Lappung	Sólo variedades que presentan lóbulos en la hoja: Limbo: profundidad de las incisiones de los lóbulos		
QN	(a)	very shallow	très peu profonde	sehr flach	muy poco profundas		1
	(d)	shallow	peu profonde	flach	poco profundas	Benihayato, Sinchunmi	3
		moderate	moyenne	mäßig	moderadamente profundas	Koganesengan	5
		deep	profonde	tief	profundas	Tsukumoaka	7
		very deep	très profonde	sehr tief	muy profundas		9
12.	VG	Leaf blade: color (excluding anthocyanin coloration)	Limbe : couleur (sans la pigmentation anthocyanique)	Blattspreite: Farbe (ohne Anthocyanfärbung)	Limbo: color (excluyendo la pigmentación antociánica)		
PQ	(a)	yellow green	vert-jaune	gelbgrün	verde amarillento	Serolane, Suio	1
	(d)	green	vert	grün	verde	Yulmi	2
		grey green	gris-vert	graugrün	gris-verde	Hayanmi	3
13.	VG	Leaf blade: anthocyanin coloration of upper side	Limbe : pigmentation anthocyanique de la face supérieure	Blattspreite: Anthocyanfärbung der Oberseite	Limbo: pigmentación antociánica del haz		
QN	(a)	absent or weak	absente ou faible	fehlend oder gering	ausente o débil	Yulmi	1
	(d)	medium	moyenne	mittel	media	Hayanmi	2
		strong	forte	stark	fuerte		3

		English	français	deutsch	español	Example Varieties/ Exemples/ Beispielssorten/ Variedades ejemplos	Note/ Nota
14.	VG	Leaf blade: extent of anthocyanin coloration on abaxial veins	Limbe : étendue de la pigmentation anthocyane sur les nervures abaxiales	Blattspreite: Ausdehnung der Anthocyanfärbung an den abaxialen Adern	Limbo: extensión de la pigmentación antociánica en los nervios abaxiales		
(+)	QN	(a) absent or very small	absente ou très petite	fehlend oder sehr gering	ausente o muy pequeña		1
	(d)	small	petite	klein	pequeña	Koukei 14, Yulmi	3
		medium	moyenne	mittel	mediana	Beniaka, Norin 45	5
		large	grande	groß	grande	Hayanmi, Naeshirazu	7
		very large	très grande	sehr groß	muy grande	Koganesengan	9
15.	VG	Leaf blade: intensity of anthocyanin coloration on abaxial veins	Limbe : intensité de la pigmentation anthocyane sur les nervures abaxiales	Blattspreite: Intensität der Anthocyanfärbung an den abaxialen Adern	Limbo: intensidad de la pigmentación antociánica en los nervios abaxiales		
	QN	(a) very weak	très faible	sehr gering	muy débil		1
	(d)	weak	faible	gering	débil	Norin 45	3
		medium	moyenne	mittel	media	Koganesengan	5
		strong	forte	stark	fuerte		7
		very strong	très forte	sehr stark	muy fuerte		9
16.	VG	Young leaf blade: main color on upper side	Jeune limbe : couleur principale sur la face supérieure	Spreite des jungen Blättes: Farbe an der Oberseite	Limbo: color principal del haz		
PQ	yellow green	vert-jaune	gelbgrün	verde amarillento	Beniwase		1
	light green	vert clair	hellgrün	verde claro	Koganesengan		2
	medium green	vert moyen	mittelgrün	verde medio	Norin 2		3
	dark green	vert foncé	dunkelgrün	verde gris			4
	light purple	pourpre clair	hellpurpur	púrpura claro	Kyushu 14		5
	medium purple	pourpre moyen	mittelpurpur	púrpura medio			6
	purplish brown	brun-pourpre	purpurbraun	marrón violáceo	Minamiyutaka		7
	light brown	brun clair	hellbraun	marrón claro			8
	dark brown	brun foncé	dunkelbraun	marrón oscuro			9

					Example Varieties/ Exemples/ Beispielssorten/ Variedades ejemplo	Note/ Nota
English	français	deutsch	español			
17. VG Petiole: anthocyanin coloration	Pétiole : pigmentation anthocyane	Blattstiell: Anthocyanfärbung	Pecíolo: pigmentación antociánica			
QN	(a) absent or very weak	absente ou très faible	fehlend oder sehr gering	ausente o muy débil	Yulmi	1
	(d) weak	faible	gering	débil	Norin 45	3
	medium	moyenne	mittel	media	Hayanmi, Koganesengan	5
	strong	forte	stark	fuerte		7
18. VG/ MS Petiole: length	Pétiole : longueur	Blattstiell: Länge	Pecíolo: longitud			
QN	(a) very short	très court	sehr kurz	muy corto	Sinchunmi	1
	(d) short	court	kurz	corto		3
	medium	moyen	mittel	medio	Koganesengan, Yulmi	5
	long	long	lang	largo		7
	very long	très long	sehr lang	muy largo	Shinmi	9
19. VG (*) (+) Storage root: shape	Racine de réserve : forme	Speicherwurzel: Form	Raíz tuberosa: forma			
PQ	(e) ovate	ovale	eiförmig	oval		1
	elliptic	elliptique	elliptisch	elíptica		2
	obovate	obovale	verkehrt eiförmig	oboval	Geomi	3
	oblong	oblongue	rechteckig	oblonga	Serolane	4
	irregular	irrégulièr	unregelmäßig	irregular	Shinyulmi	5

					Example Varieties/ Exemples/ Beispielssorten/ Variedades ejemplos	Note/ Nota
20.	MS	Storage root: ratio length/width	Racine de réserve : rapport longueur/largeur	Speicherwurzel: Verhältnis Länge/Breite	Raíz tuberosa: relación longitud/anchura	
QN	(e)	small (moderately compressed)	petit (modérément compressé)	klein (mäßig zusammengedrückt)	pequeña (moderadamente comprimida)	Norin 2 3
		medium	moyen	mittel	media	Geonmi 5
		large (moderately elongated)	grand (modérément allongé)	groß (mäßig länglich)	grande (moderadamente alargada)	Yulmi 7
21.	MS/ VG	Storage root: VG thickness of cortex relative to overall diameter (+)	Racine de réserve : épaisseur du cortex par rapport au diamètre total	Speicherwurzel: Dicke der Rinde im Verhältnis zum Gesamtdurchmesser	Raíz tuberosa: grosor de la corteza en relación con el diámetro general	
QN	(e)	thin	fin	dünn	delgada	Yulmi 3
		medium	moyen	mittel	media	5
		thick	épais	dick	gruesa	Shingeonmi 7
22.	VG	Storage root: main color of skin (*) (+)	Racine de réserve : couleur principale de la peau	Speicherwurzel: Hauptfarbe der Schale	Raíz tuberosa: color principal de la piel	
PQ	(e)	white	blanc	weiß	blanco	Joy White 1
		light beige	beige clair	hellbeige	beige claro	Chinmi, Koganesengan 2
		yellow	jaune	gelb	amarillo	Impilo 3
		orange	orange	orange	naranja	Benihayato, Serolane 4
		brownish orange	orange brûnatre	bräunlichorange	naranja amarronado	Khano 5
		pink	rose	rosa	rosa	Yulmi 6
		red	rouge	rot	rojo	Koukei 14, Shinhwangmi 7
		purple red	rouge-pourpre	purpurrot	rojo púrpura	Beniazuma, Phala 8
		light purple	pourpre clair	hellpurpur	púrpura claro	9
		medium purple	pourpre moyen	mittelpurpur	púrpura medio	Ayamurasaki, Zami 10
		brown	brun	braun	marrón	Happymi 11

					Example Varieties/ Exemples/ Beispielssorten/ Variedades ejemplos	Note/ Nota
23.	VG (+)	Storage root: secondary color of skin	Racine de réserve : couleur secondaire de la peau	Speicherwurzel: Sekundärfarbe der Schale	Raíz tuberosa: color secundario de la piel	
PQ	(e)	absent	absente	fehlend	ausente	Koganesengan
		white	blanc	weiß	blanco	Tamayutaka
		yellow	jaune	gelb	amarillo	
		orange	orange	orange	naranja	
		pink	rose	rosa	rosa	Koukei 14
		red	rouge	rot	rojo	Nakamurasaki
		purple	pourpre	purpurn	púrpura	Benikomachi
		brown	brun	braun	marrón	Koganesengan
24.	VG (*) (+)	Storage root: main color of flesh	Racine de réserve : couleur de la chair	Speicherwurzel: Hauptfarbe des Fleisches	Raíz tuberosa: color principal de la pulpa	
PQ	(e)	white	blanc	weiß	blanco	Hayanmi, Shirosengan
		beige	beige	beige	beige	Nakamurasaki, Koukei 14
		yellow	jaune	gelb	amarillo	Benikomachi, Yulmi
		orange	orange	orange	naranja	Benihayato, Hayatoimo, Juhwangmi
		purple	pourpre	purpurn	púrpura	Ayamurasaki, Borami
25.	VG	Storage root: intensity of main color of flesh	Racine de réserve : intensité de la couleur principale de la chair	Speicherwurzel: Intensität der Hauptfarbe des Fleisches	Raíz tuberosa: intensidad del color principal de la pulpa	
QN	(e)	light	claire	hell	claro	Borami, Hayatoimo, Yulmi
		medium	moyenne	mittel	medio	Jinhongmi, Shinwangmi, Zami
		dark	foncé	dunkel	oscuro	Juhwangmi, Shinyulmi, Sinjami

					Example Varieties/ Exemples/ Beispielssorten/ Variedades ejemplar	Note/ Nota
26.	VG (+)	Storage root: secondary color of flesh	Racine de réserve : couleur secondaire de la chair	Speicherwurzel: Sekundärfarbe des Fleisches	Raíz tuberosa: color secundario de la pulpa	
PQ	(e)	white	blanc	weiß	blanco	1
		light beige	beige clair	hellbeige	beige claro	2
		yellow	jaune	gelb	amarillo	Hayatoimo
		orange	orange	orange	naranja	Toka Toka Gold
		pink	rose	rosa	rosa	5
		red	rouge	rot	rojo	6
		red-purple	rouge-pourpre	rotpurpurn	rojo púrpura	Nakamurasaki, Owairka Red
		purple	pourpre	purpurn	púrpura	8
27.	VG	Storage root: depth of eyes	Racine de réserve : profondeur des yeux	Speicherwurzel: Tiefe der Augen	Raíz tuberosa: profundidad de los ojos	
QN	(e)	shallow	peu profonde	flach	poco profundos	Beniaka
		medium	moyenne	mittel	medios	Koukei 14
		deep	profonde	tief	profundos	Kantou 80

8. Explanations on the Table of Characteristics


8.1 Explanations covering several characteristics

Characteristics containing the following key in the second column of the Table of Characteristics should be examined as indicated below:


- (a) Observations should be made after 90 days from planting.
- (b) To be observed on the main stem
- (c) Stem internodes and diameter should be observed on an internode located in the middle third of the main stem.
- (d) Observations to be made on fully developed leaves at the middle part of the main stem.
- (e) Characteristics should be observed after harvest.

8.2. Explanations for individual characteristics


Ad. 9: Leaf blade: lobes




Ad. 10: Only varieties with leaf blade lobes: absent: Leaf blade: shape




Ad. 11: Only varieties with leaf lobes present: Leaf blade: depth of lobing



Ad. 14: Leaf blade: extent of anthocyanin coloration on abaxial veins





Ad. 18: Petiole: length




Ad. 19: Storage root: shape

		< position of broadest part >		
		below middle	at middle	above middle
< lateral outline >	rounded			
		1 ovate	2 elliptic	3 obovate



Ad. 21: Storage root: thickness of cortex relative to overall diameter



Ad. 22: Storage root: main color of skin

The main color is the color which covers the largest area of skin.

Ad. 23: Storage root: secondary color of skin

The secondary color is the color with the second largest surface area of skin.

Ad. 24: Storage root: main color of flesh

The main color is the color with the largest surface area of storage root in cross section.

Ad. 26: Storage root: secondary color of flesh

The secondary color is the color with the second largest surface area of storage root in cross section.

9. Literature

NSMO, 2000: Test Guideline for Sweetpotato. National Seed Management Office/MAF, KR, p.12.

Mokpo experiment station/RDA. 2002: Production and Use of Sweetpotato. Mokpo experiment station/RDA, p. 214.

Zosimo Huaman. 1992: Morphologic Identification of Duplicates in Collections of *Ipomoea batatas*. CIP Research guide 36. CIP, p. 28.

Zosimo Huaman. 2002: Section 1.1 Systemic Botany and Morphology of the Sweetpotato plant. Sweetpotato Germplasm Management Training Manual. International Potato Center (CIP), p. 7.

Zosimo Huaman, 2006: Systmatic Botany and Morphology of the Sweetpotato Plant. Sweetpotato Germplasm Management (*Ipomoea batatas*). Training manual CIP. <http://www.cipotato.org>.

10. Technical Questionnaire

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
		Application date: (not to be filled in by the applicant)
<p style="text-align:center">TECHNICAL QUESTIONNAIRE to be completed in connection with an application for plant breeders' rights</p>		
1. Subject of the Technical Questionnaire		
1.1 Botanical name	<i>Ipomoea batatas</i> (L.) Lam.	
1.2 Common name	Sweet potato	
2. Applicant		
Name		
Address		
Telephone No.		
Fax No.		
E-mail address		
Breeder (if different from applicant)		
3. Proposed denomination and breeder's reference		
Proposed denomination (if available)		
Breeder's reference		

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

#4. Information on the breeding scheme and propagation of the variety

4.1 Breeding scheme

Variety resulting from:

4.1.1 Crossing

(a) controlled cross []
(please state parent varieties)

(b) partially known cross []
(please state known parent variety(ies))

(c) unknown cross []

4.1.2 Mutation []
(please state parent variety)

4.1.3 Discovery and development []
(please state where and when discovered
and how developed)

4.1.4 Other []
(please provide details)

4.2 Method of propagating the variety

4.2.1 Vegetative propagation

(a) cuttings []

(b) *in vitro* propagation []

(c) other (state method) []

4.2.2 Seed []

4.2.3 Other []
(please provide details)

* Authorities may allow certain of this information to be provided in a confidential section of the Technical Questionnaire.

TECHNICAL QUESTIONNAIRE		Page {x} of {y}	Reference Number:
5. Characteristics of the variety to be indicated (the number in brackets refers to the corresponding characteristic in Test Guidelines; please mark the note which best corresponds).			
Characteristics		Example Varieties	Note
5.1	Plant: growth habit		
(1)			
	upright	Sinchunmi	1[]
	semi-upright	Younmi	3[]
	spreading	Yulmi	5[]
5.2	Stem: anthocyanin coloration of tip		
(6)			
	absent or weak	Yulmi	1[]
	medium	Sinjami	2[]
	strong	Hayanmi	3[]
5.3	Leaf blade: lobes		
(9)			
	absent	Gokokuimo	1[]
	three lobes	Benisengan	2[]
	five lobes	Koganesengan, Sinchunmi	3[]
	seven lobes	Benihayato	4[]
5.4	Storage root: shape		
(19)			
	ovate		1[]
	elliptic		2[]
	obovate	Geomi	3[]
	oblong	Serolane	4[]
	irregular	Shinyulmi	5[]

TECHNICAL QUESTIONNAIRE		Page {x} of {y}	Reference Number:
Characteristics		Example Varieties	Note
5.5 Storage root: main color of skin (22)			
white		Joy White	1[]
light beige		Chinmi, Koganesengan	2[]
yellow		Impilo	3[]
orange		Benihayato, Serolane	4[]
brownish orange		Khano	5[]
pink		Yulmi	6[]
red		Koukei 14, Shinhwangmi	7[]
purple red		Beniazuma, Phala	8[]
light purple			9[]
medium purple		Ayamurasaki, Zami	10[]
brown		Happymi	11[]
5.6 Storage root: main color of flesh (24)			
white		Hayanmi, Shirosangan	1[]
beige		Nakamuraski, Koukei 14	2[]
yellow		Benikomachi, Yulmi	3[]
orange		Benihayato, Hayatoimo, Juhwangmi	4[]
purple		Ayamurasaki, Borami	5[]

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

6. Similar varieties and differences from these varieties

Please use the following table and box for comments to provide information on how your candidate variety differs from the variety (or varieties) which, to the best of your knowledge, is (or are) most similar. This information may help the examination authority to conduct its examination of distinctness in a more efficient way.

Denomination(s) of variety(ies) similar to your candidate variety	Characteristic(s) in which your candidate variety differs from the similar variety(ies)	Describe the expression of the characteristic(s) for the similar variety(ies)	Describe the expression of the characteristic(s) for your candidate variety
<i>Example</i>	<i>Plant: growth habit</i>	<i>upright</i>	<i>semi-upright</i>
Comments:			

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

#7. Additional information which may help in the examination of the variety		
7.1 In addition to the information provided in sections 5 and 6, are there any additional characteristics which may help to distinguish the variety?		
Yes	[]	No []
(If yes, please provide details)		
7.2 Are there any special conditions for growing the variety or conducting the examination?		
Yes	[]	No []
(If yes, please provide details)		
7.3 Other information		
7.3.1 What's the varietal usage?		
Food/Feed [] Ornamental []		
8. Authorization for release		
(a) Does the variety require prior authorization for release under legislation concerning the protection of the environment, human and animal health?		
Yes	[]	No []
(b) Has such authorization been obtained?		
Yes	[]	No []
If the answer to (b) is yes, please attach a copy of the authorization.		

Authorities may allow certain of this information to be provided in a confidential section of the Technical Questionnaire.

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

9. Information on plant material to be examined or submitted for examination.

9.1 The expression of a characteristic or several characteristics of a variety may be affected by factors, such as pests and disease, chemical treatment (e.g. growth retardants or pesticides), effects of tissue culture, different rootstocks, scions taken from different growth phases of a tree, etc.

9.2 The plant material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If the plant material has undergone such treatment, full details of the treatment must be given. In this respect, please indicate below, to the best of your knowledge, if the plant material to be examined has been subjected to:

- | | | |
|---|---------|--------|
| (a) Microorganisms (e.g. virus, bacteria, phytoplasma) | Yes [] | No [] |
| (b) Chemical treatment (e.g. growth retardant, pesticide) | Yes [] | No [] |
| (c) Tissue culture | Yes [] | No [] |
| (d) Other factors | Yes [] | No [] |

Please provide details for where you have indicated “yes”.

.....

10. I hereby declare that, to the best of my knowledge, the information provided in this form is correct:

Applicant's name

Signature Date

[End of document]