

UPOV

TG/SHITK(proj.4)

ORIGINAL: English

DATE: 2011-10-17

INTERNATIONAL UNION FOR THE PROTECTION OF NEW VARIETIES OF PLANTS

GENEVA

DRAFT

SHIITAKE

UPOV Code: LENTI_EDO

Lentinula edodes (Berk.) Pegler

GUIDELINES

FOR THE CONDUCT OF TESTS

FOR DISTINCTNESS, UNIFORMITY AND STABILITY

*prepared by an expert from Japan**to be considered by the**Enlarged Editorial Committee at its meeting
to be held in Geneva, on January 11 and 12, 2012*

Alternative Names:*

<i>Latin</i>	<i>English</i>	<i>French</i>	<i>German</i>	<i>Spanish</i>
<i>Lentinula edodes</i> (Berk.) Pegler <i>Lentinus elodes</i> (Berk.) Sing.	Shiitake Oak Mushroom	Shiitake	Pasaniapilz	Shiitake

The purpose of these guidelines (“Test Guidelines”) is to elaborate the principles contained in the General Introduction (document TG/1/3), and its associated TGP documents, into detailed practical guidance for the harmonized examination of distinctness, uniformity and stability (DUS) and, in particular, to identify appropriate characteristics for the examination of DUS and production of harmonized variety descriptions.

ASSOCIATED DOCUMENTS

These Test Guidelines should be read in conjunction with the General Introduction and its associated TGP documents.

* These names were correct at the time of the introduction of these Test Guidelines but may be revised or updated. [Readers are advised to consult the UPOV Code, which can be found on the UPOV Website (www.upov.int), for the latest information.]

<u>TABLE OF CONTENTS</u>	<u>PAGE</u>
1. SUBJECT OF THESE TEST GUIDELINES.....	3
2. MATERIAL REQUIRED	3
3. METHOD OF EXAMINATION.....	3
3.1 Number of Growing Cycles	3
3.2 Testing Place	3
3.3 Conditions for Conducting the Examination.....	4
3.4 Test Design	4
3.5 Additional Tests	4
4. ASSESSMENT OF DISTINCTNESS, UNIFORMITY AND STABILITY	4
4.1 Distinctness	4
4.2 Uniformity.....	5
4.3 Stability	6
5. GROUPING OF VARIETIES AND ORGANIZATION OF THE GROWING TRIAL.....	6
6. INTRODUCTION TO THE TABLE OF CHARACTERISTICS	6
6.1 Categories of Characteristics.....	6
6.2 States of Expression and Corresponding Notes.....	7
6.3 Types of Expression.....	7
6.4 Example Varieties	8
6.5 Legend.....	8
7. TABLE OF CHARACTERISTICS/TABLEAU DES CARACTÈRES/MERKMALSTABELLE/TABLA DE CARACTERES.....	9
8. EXPLANATIONS ON THE TABLE OF CHARACTERISTICS	18
8.1 Explanations covering several characteristics	18
8.2 Explanations for individual characteristics	19
8.3 Growing Types.....	26
9. LITERATURE	27
10. TECHNICAL QUESTIONNAIRE.....	28

1. Subject of these Test Guidelines

These Test Guidelines apply to all varieties of *Lentinula edodes* (Berk.) Pegler

2. Material Required

2.1 The competent authorities decide on the quantity and quality of the plant material required for testing the variety and when and where it is to be delivered. Applicants submitting material from a State other than that in which the testing takes place must ensure that all customs formalities and phytosanitary requirements are complied with.

2.2 The material is to be supplied in the form of spawn and as a pure culture on a suitable medium.

(a) Spawn should be of a quality which ensures that all relevant characteristics of the variety will be expressed. In particular, mycelium on grain should be visible to the naked eye, the grain should not be colonized to such an extent that kernels stick together. The spawn should not be older than 3 months and should have been stored under proper conditions.

(b) Pure cultures must be on slant agar tubes with an appropriate medium such as PDA (potato dextrose agar) or malt extract agar. Tubes should be covered by cotton plugs or plastic caps allowing sterile air diffusion. Cultures should be fresh, i.e. not stored for longer than 2 weeks at low temperature.

2.3 The minimum quantity of material, to be supplied by the applicant, should be:

2 liter of spawn and 3 slant tubes containing secondary mycelium by pure culture.
[see Additional information (b)]

2.4 The material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If it has been treated, full details of the treatment must be given.

3. Method of Examination

3.1 *Number of Growing Cycles*

The minimum duration of tests should normally be two independent growing cycles. The growing cycle is considered to be from spawning until the end of the first flush.

3.2 *Testing Place*

Tests are normally conducted at one place. In the case of tests conducted at more than one place, guidance is provided in TGP/9 "Examining Distinctness".

3.3 *Conditions for Conducting the Examination*

The tests should be carried out under conditions ensuring satisfactory growth for the expression of the relevant characteristics of the variety and for the conduct of the examination.

3.4 *Test Design*

3.4.1 Each test should be designed to result in a total of at least 60 bed-logs or 60 sawdust blocks, which should be divided between at least three replicates.

3.4.2 The design of the tests should be such that fruit bodies or parts of fruit bodies may be removed for measurement or counting without prejudice to the observations which must be made up to the end of the growing cycle.

3.5 *Additional Tests*

Additional tests, for examining relevant characteristics, may be established.

4. Assessment of Distinctness, Uniformity and Stability

4.1 *Distinctness*

4.1.1 General Recommendations

It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding distinctness. However, the following points are provided for elaboration or emphasis in these Test Guidelines.

4.1.2 Consistent Differences

The differences observed between varieties may be so clear that more than one growing cycle is not necessary. In addition, in some circumstances, the influence of the environment is not such that more than a single growing cycle is required to provide assurance that the differences observed between varieties are sufficiently consistent. One means of ensuring that a difference in a characteristic, observed in a growing trial, is sufficiently consistent is to examine the characteristic in at least two independent growing cycles.

4.1.3 Clear Differences

Determining whether a difference between two varieties is clear depends on many factors, and should consider, in particular, the type of expression of the characteristic being examined, i.e. whether it is expressed in a qualitative, quantitative, or pseudo-qualitative manner. Therefore, it is important that users of these Test Guidelines are familiar with the recommendations contained in the General Introduction prior to making decisions regarding distinctness.

4.1.4 Number of Fruit Bodies / Parts of Fruit Bodies to be Examined

Unless otherwise indicated, for the purposes of distinctness, all observations on single fruit bodies should be made on 60 fruit bodies or parts taken from each of 60 fruit bodies and any other observations made on all fruit bodies in the test, disregarding any off-type fruit bodies.

4.1.5 Method of Observation

The recommended method of observing the characteristic for the purposes of distinctness is indicated by the following key in the second column of the Table of Characteristics (see document TGP/9 “Examining Distinctness”, Section 4 “Observation of characteristics”):

MG: single measurement of a group of fruit bodies or parts of fruit bodies

MS: measurement of a number of individual fruit bodies or parts of fruit bodies

VG: visual assessment by a single observation of a group of fruit bodies or parts of fruit bodies

VS: visual assessment by observation of individual fruit bodies or parts of fruit bodies

Type of observation: visual (V) or measurement (M)

“Visual” observation (V) is an observation made on the basis of the expert’s judgment. For the purposes of this document, “visual” observation refers to the sensory observations of the experts and, therefore, also includes smell, taste and touch. Visual observation includes observations where the expert uses reference points (e.g. diagrams, example varieties, side-by-side comparison) or non-linear charts (e.g. color charts). Measurement (M) is an objective observation against a calibrated, linear scale e.g. using a ruler, weighing scales, colorimeter, dates, counts, etc.

Type of record: for a group of fruit bodies (G) or for single, individual fruit bodies (S)

For the purposes of distinctness, observations may be recorded as a single record for a group of fruit bodies or parts of fruit bodies (G), or may be recorded as records for a number of single, individual fruit bodies or parts of fruit bodies (S). In most cases, “G” provides a single record per variety and it is not possible or necessary to apply statistical methods in a fruit body -by- fruit body analysis for the assessment of distinctness.”

In cases where more than one method of observing the characteristic is indicated in the Table of Characteristics (e.g. VG/MG), guidance on selecting an appropriate method is provided in document TGP/9, Section 4.2.

4.2 *Uniformity*

4.2.1 It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding uniformity. However, the following points are provided for elaboration or emphasis in these Test Guidelines:

4.2.2 For the assessment of uniformity, a population standard of 1% and an acceptance probability of at least 95% should be applied. In the case of a sample size of 50 fruit bodies, 2 off-types are allowed.

4.3 *Stability*

4.3.1 In practice, it is not usual to perform tests of stability that produce results as certain as those of the testing of distinctness and uniformity. However, experience has demonstrated that, for many types of variety, when a variety has been shown to be uniform, it can also be considered to be stable.

4.3.2 Where appropriate, or in cases of doubt, stability may be further examined by testing a new stock to ensure that it exhibits the same characteristics as those shown by the initial material supplied.

5. Grouping of Varieties and Organization of the Growing Trial

5.1 The selection of varieties of common knowledge to be grown in the trial with the candidate varieties and the way in which these varieties are divided into groups to facilitate the assessment of distinctness are aided by the use of grouping characteristics.

5.2 Grouping characteristics are those in which the documented states of expression, even where produced at different locations, can be used, either individually or in combination with other such characteristics: (a) to select varieties of common knowledge that can be excluded from the growing trial used for examination of distinctness; and (b) to organize the growing trial so that similar varieties are grouped together.

5.3 The following have been agreed as useful grouping characteristics:

- (a) Cultivation type (1: bed-log cultivation type, 2: sawdust cultivation type) (TQ)
- (b) Cap: shape of vertical section (characteristic 9)
- (c) Cap: main color of apex (characteristic 11)
- (d) Cap: presence of scale (characteristic 14)
- (e) Gill: presence of gill (characteristic 18)
- (f) Stipe: shape of vertical section (characteristic 24)

5.4 Guidance for the use of grouping characteristics, in the process of examining distinctness, is provided through the General Introduction and document TGP/9 "Examining Distinctness".

6. Introduction to the Table of Characteristics

6.1 *Categories of Characteristics*

6.1.1 Standard Test Guidelines Characteristics

Standard Test Guidelines characteristics are those which are approved by UPOV for examination of DUS and from which members of the Union can select those suitable for their particular circumstances.

6.1.2 Asterisked Characteristics

Asterisked characteristics (denoted by *) are those included in the Test Guidelines which are important for the international harmonization of variety descriptions and should always be examined for DUS and included in the variety description by all members of the Union, except when the state of expression of a preceding characteristic or regional environmental conditions render this inappropriate.

6.2 States of Expression and Corresponding Notes

6.2.1 States of expression are given for each characteristic to define the characteristic and to harmonize descriptions. Each state of expression is allocated a corresponding numerical note for ease of recording of data and for the production and exchange of the description.

6.2.2 In the case of qualitative and pseudo-qualitative characteristics (see Chapter 6.3), all relevant states of expression are presented in the characteristic. However, in the case of quantitative characteristics with 5 or more states, an abbreviated scale may be used to minimize the size of the Table of Characteristics. For example, in the case of a quantitative characteristic with 9 states, the presentation of states of expression in the Test Guidelines may be abbreviated as follows:

State	Note
small	3
medium	5
large	7

However, it should be noted that all of the following 9 states of expression exist to describe varieties and should be used as appropriate:

State	Note
very small	1
very small to small	2
small	3
small to medium	4
medium	5
medium to large	6
large	7
large to very large	8
very large	9

6.2.3 Further explanation of the presentation of states of expression and notes is provided in document TGP/7 “Development of Test Guidelines”.

6.3 Types of Expression

An explanation of the types of expression of characteristics (qualitative, quantitative and pseudo-qualitative) is provided in the General Introduction.

6.4 *Example Varieties*

Where appropriate, example varieties are provided to clarify the states of expression of each characteristic. The type is indicated in brackets after the name of the example variety as follows:

- (B) bed-log cultivation type
- (S) sawdust cultivation type

6.5 *Legend*

(*) Asterisked characteristic – see Chapter 6.1.2

QL Qualitative characteristic – see Chapter 6.3

QN Quantitative characteristic – see Chapter 6.3

PQ Pseudo-qualitative characteristic – see Chapter 6.3

MG, MS, VG, VS – see Chapter 4.1.5

(a) – (c) See Explanations on the Table of Characteristics in Chapter 8.1

(+) See Explanations on the Table of Characteristics in Chapter 8.2

(B) bed-log cultivation type

(S) sawdust cultivation type

See Explanations of the growing types in Chapter 8.3

7. Table of Characteristics/Tableau des caractères/Merkmalstabelle/Tabla de caracteres

	English	français	deutsch	español	Example Varieties/ Exemples/ Beispielssorten/ Variedades ejemplo	Note/ Nota	
1.	VG	Density of hyphae on the medium	Densité de l'hyphe sur le support	Dichte der Hyphe auf dem Medium	Densidad de las hifas en el medio		
(+)							
N	(a)	sparse	lâche	locker	baja	HS607, Mori XR1	1
		intermediate	moyenne	mittel	intermedia	Morino Natsumi	2
		dense	dense	dicht	densa	KX-S005	3
2.	VG	Colony: tinting of surface on the medium	Colonie : coloration de la surface du support	Kolonie: Färbung der Oberfläche auf dem Medium	Colonia: tinte de la superficie en el medio		
(+)							
QL	(a)	absent	absente	fehlend	ausente	Kinko 115, Mori XR1, Morino Natsumi	1
		present	présente	vorhanden	presente	HS607, KX-S005	9
3.	MS	Mycelium: optimum temperature for growth	Mycélium : température optimale de culture	Myzel: optimale Wachstums-temperatur	Micelio: temperatura optima de desarrollo		
(+)							
QN	(b)	23°C	23°C	23°C	23°C	Kinko 243	3
		25°C	25°C	25°C	25°C	HS607, Kinko 115	5
		27°C	27°C	27°C	27°C	Morino Natsumi	7
4.	MS	Mycelium: growth rate at 10°C	Mycélium : vitesse de croissance à 10°C	Myzel: Wachstumsrate bei 10°C	Micelio: índice de desarrollo a 10°C		
(+)							
QN	(b)	very slow	très lente	sehr langsam	muy lento	Kinko 115	1
		slow	lente	langsam	lento	Kinoh1	2
		medium	moyenne	mittel	medio	HS607, Morino Natsumi	3
		fast	rapide	schnell	rápido	KX-S005	4
		very fast	très rapide	sehr schnell	muy rápido	Yujiro	5

	English	français	deutsch	español	Example Varieties/ Exemples/ Beispielssorten/ Variedades ejemplo	Note/ Nota
5.	MS	Mycelium: growth rate at 15°C	Mycélium : vitesse de croissance à 15°C	Myzel: Wachstumsrate bei 15°C	Micelio: índice de desarrollo a 15°C	
(+)						
QN	(b)	very slow	très lente	sehr langsam	muy lento	1
		slow	lente	langsam	lento	Kinko 115
		medium	moyenne	mittel	medio	HS607, Susono 360
		fast	rapide	schnell	rápido	Yujiro
		very fast	très rapide	sehr schnell	muy rápido	KX-S005
6.	MS	Mycelium: growth rate at 20°C	Mycélium : vitesse de croissance à 20°C	Myzel: Wachstumsrate bei 20°C	Micelio: índice de desarrollo a 20°C	
(*)						
(+)						
QN	(b)	very slow	très lente	sehr langsam	muy lento	Bridge 32, Kinno 1
		slow	lente	langsam	lento	Kinko 115
		medium	moyenne	mittel	medio	ML8, Morino Natsumi
		fast	rapide	schnell	rápido	Morino Harumitsu
		very fast	très rapide	sehr schnell	muy rápido	Akiyama A-950, Hokken 600, JMS 237
7.	MS	Mycelium: growth rate at 25°C	Mycélium : vitesse de croissance à 25°C	Myzel: Wachstumsrate bei 25°C	Micelio: índice de desarrollo a 25°C	
(+)						
QN	(b)	very slow	très lente	sehr langsam	muy lento	1
		slow	lente	langsam	lento	Kinko 115
		medium	moyenne	mittel	medio	HS73, Susono 360
		fast	rapide	schnell	rápido	Hokken600, Yujiro
		very fast	très rapide	sehr schnell	muy rápido	5

	English	français	deutsch	español	Example Varieties/ Exemples/ Beispielssorten/ Variedades ejemplo	Note/ Nota	
8. (*) (+)	MS	Mycelium: growth rate at 30°C	Mycélium : vitesse de croissance à 30°C	Myzel: Wachstumsrate bei 30°C	Micelio: índice de desarrollo a 30°C		
QN	(b)	very slow	très lente	sehr langsam	muy lento	Akiyama A-526	1
		slow	lente	langsam	lento	HS73, Morino Harumitsu	2
		medium	moyenne	mittel	medio	Kinko 115	3
		fast	rapide	schnell	rápido	Mori XR1, Susono 360	4
		very fast	très rapide	sehr schnell	muy rápido	Morino Natsumi, Yujiro	5
9. (*) (+)	VG	Cap: shape of vertical section	Chapeau : forme de la section verticale	Hut: Form im Senkrechtschnitt	Sombrero: forma de la sección vertical		
PQ	(c)	concave	concave	konkav	cóncava	JMS 7H-1	1
		flat	aplatie	flach	llana	Morino Harumitsu	2
		round	arrondie	rund	redonda	Kinko 115, Yujiro	3
		convex	convexe	konvex	convexa	KX-S005	4
10. (*) (+)	VG/ MS	Cap: diameter	Chapeau : diamètre	Hut: Durchmesser	Sombrero: diámetro		
QN	(c)	small	petit	klein	pequeño	Morino Harumitsu	3
		medium	moyen	mittel	medio	HS73, Kinko 115, Mori XR1	5
		large	grand	groß	grande	Kinko 117, Mori 505	7
11. (*)	VG	Cap: main color of apex	Chapeau : couleur principale du sommet	Hut: Hauptfarbe der Spitze	Sombrero: color principal del ápice		
PQ	(c)	white	blanc	weiß	blanco	Kinko 989	1
		yellowish brown	brun jaunâtre	gelblichbraun	marrón amarillento	Mori XR-1	2
		brown	brun	braun	marrón	Kinko 115, Susono 360	3
		reddish brown	brun rougeâtre	rötlichbraun	marrón rojizo	Akiyama A-526	4

	English	français	deutsch	español	Example Varieties/ Exemples/ Beispielssorten/ Variedades ejemplo	Note/ Nota	
12.	MS	Cap: height	Chapeau : hauteur	Hut: Höhe	Sombrero: altura		
	(+)						
QN	(c)	low	basse	niedrig	bajo	Morino Harumitsu	3
		medium	moyenne	mittel	medio	Mori XR1, Morino Natsumi, Susono 360	5
		high	haute	hoch	alto	Akiyama A-526	7
13.	VS	Cap: firmness	Chapeau : fermeté	Hut: Festigkeit	Sombrero: firmeza		
	(+)						
QN	(c)	soft	mou	weich	suave	Kinko 650	1
		medium	moyen	mittel	medio	HS607, Kinko 115, KX-S055, Susono 360	2
		solid	ferme	fest	sólido	Morino Natsumi	3
14.	VG	Cap: presence of scale	Chapeau : présence d'écailles	Hut: Vorhandensein von Schuppen	Sombrero: presencia de escamas		
	(*)						
QL	(c)	absent	absentes	fehlend	ausentes	KX-S034	1
		present	présentes	vorhanden	presentes	HS73, Kinko 115, ML8, Morino Natsumi	9
15.	VG	Cap: distribution of scale	Chapeau : répartition des écailles	Hut: Verteilung der Schuppen	Sombrero: distribución de las escamas		
	(+)						
QN	(c)	whole	sur toute la surface	überall	en toda la superficie	Kinko 115, Mori XR1	1
		periphery	sur la périphérie	am Rand	en la periferia	Morino Natsumi, Susono 360, Yujiro	2
16.	VG	Cap: size of scale	Chapeau : taille des écailles	Hut: Größe der Schuppen	Sombrero: tamaño de las escamas		
	(+)						
QN	(c)	small	petites	klein	pequeño	HS73, Mori XR1	3
		medium	moyennes	mittel	medio	Morino Natsumi, Susono 360, Yujiro	5
		large	grandes	groß	grande	Kinko 169	7

	English	français	deutsch	español	Example Varieties/ Exemples/ Beispielssorten/ Variedades ejemplo	Note/ Nota	
17.	VG	Cap: tinting of scale	Chapeau : coloration des écailles	Hut: Färbung der Schuppen	Sombrero: tinte de las escamas		
(+)							
QL	(c)	absent	absente	fehlend	ausente	JMS5K16, ML8, Morino Natsumi	1
		present	présente	vorhanden	presente	HS73, Yujiro	9
18.	VG	Cap: presence of gill	Chapeau : présence de lamelles	Hut: Vorhandensein von Lamellen	Sombrero: presencia de láminas		
(*)							
(+)							
QL	(c)	absent	absentes	fehlend	ausentes	FERM P-14310	1
		present	présentes	vorhanden	presentes	Kinko 115, Mori XR1	9
19.	VG	Gill: shape	Lamelles : forme	Lamellen: Form	Láminas: forma		
(+)							
QL	(c)	separate from stipe	séparées du pied	vom Stiel frei stehend	separadas del pie	Kinko 115, Yujiro	1
		attached to stipe	rattachées au pied	am Stiel anhaftend	pegadas al pie	Hokken 600, Mori 505	2
20.	VG	Gill: arrangement of row	Lamelles : disposition	Lamellen: Anordnung der Reihe	Láminas: disposición de las hileras		
(*)							
(+)							
QL	(c)	straight	droites	gerade	rectas	Kinko 115, KX-S055, Morino Natsumi	1
		ripple or crinkle	irrégulières ou plissées	gewellt oder gekräuselt	onduladas o arrugadas	Akiyama A-526, Mori XR1	2
21.	VG	Gill: width	Lamelles : largeur	Lamellen: Breite	Láminas: anchura		
(+)							
QN	(c)	very narrow	très étroites	sehr schmal	muy estrecha	Mori XR1	1
		narrow	étroites	schmal	estrecha	Yujiro	3
		medium	moyennes	mittel	media	Susono 360	5
		wide	larges	breit	ancha	KX-S034	7
		very wide	très larges	sehr breit	muy ancha		9

	English	français	deutsch	español	Example Varieties/ Exemples/ Beispielssorten/ Variedades ejemplo	Note/ Nota	
22.	VG	Gill: density of row	Lamelles : densité	Lamellen: Dichte der Reihe	Láminas: densidad de las hileras		
(+)							
QN	(c)	sparse	lâche	locker	baja	Kinko 169, Mori 476	1
		medium	moyenne	mittel	media	Yujiro	2
		dense	dense	dicht	densa	Kinko 115, Mori XR1, Morino Natsumi	3
23.	VG	Gill: color	Lamelles : couleur	Lamellen: Farbe	Láminas: color		
PQ	(c)	white	blanc	weiß	blanco	Kinko 115, Mori XR1, Morino Natsumi	1
		light yellow	jaune clair	hellgelb	amarillo claro	HS607, KX-S055	2
		light yellow orange	jaune orangé clair	hell gelborange	naranja amarillo claro	HS73	4
24.	VG	Stipe: shape in vertical section	Stipe : forme de la section verticale	Stiel: Form im Längsschnitt	Pie: forma en sección vertical		
(*)							
(+)							
PQ	(c)	broader toward cap	plus large vers le chapeau	breiter am Hut	ensanchada hacia el sombrero	Susono 360	1
		cylindrical	cyllindrique	zylindrisch	cilíndrica	JMS5K16, Kinko 115, Mori XR1, Morino Natsumi	2
		broader toward base	plus large vers la base	breiter an der Basis	ensanchada hacia la base	JMS 7H-1	3
25.	MS	Stipe: length	Stipe : longueur	Stiel: Länge	Pie: longitud		
(*)							
(+)							
QN	(c)	short	courte	kurz	corto	Mori XR1, Morino Natsumi, Susono 360	3
		medium	moyenne	mittel	medio	HS702, Kinko 117	5
		long	longue	lang	largo	Akiyama A-526	7
26.	MS	Stipe: diameter	Stipe : diamètre	Stiel: Durchmesser	Pie: diámetro		
(+)							
QN	(c)	small	petit	klein	pequeño	Morino Natsumi	3
		medium	moyen	mittel	medio	HS73, Susono 360	5
		large	grand	groß	grande	Kinko 115	7

	English	français	deutsch	español	Example Varieties/ Exemples/ Beispielssorten/ Variedades ejemplo	Note/ Nota	
27.	VG	Stipe: tinting	Stipe : coloration	Stiel: Färbung	Pie: tinte		
(*) (+)							
QL	(c)	absent	absente	fehlend	ausente	Mori XR1	1
		present	présente	vorhanden	presente	Kinko 115, KX-S055 Morino Natsumi	9
28.	VG	Stipe: density of fluff	Stipe : densité des peluches	Stiel: Dichte des Flaums	Pie: densidad de la pelusa		
(+)							
QN	(c)	absent or sparse	absente ou lâche	fehlend oder locker	ausente o escasa	Kinko 989	1
		medium	moyenne	mittel	media	Kinko 115, KX-S055, Morino Natsumi	2
		dense	dense	dicht	densa	KB-2010	3
29.	VG	Stipe: tinting of fluff	Stipe : coloration des peluches	Stiel: Färbung des Flaums	Pie: tinte de la pelusa		
(+)							
QL	(c)	absent	absente	fehlend	ausente	KX-S055, Mori XR1	1
		present	présente	vorhanden	presente	Kinko 115, Morino Natsumi	9
30.	VG	Stipe: firmness	Stipe : fermeté	Stiel: Festigkeit	Pie: firmeza		
(+)							
QN	(c)	soft	mou	weich	suave	HS802, Kinno 7	1
		medium	moyen	mittel	media	HS607, Mori XR1, Susono 360	2
		solid	ferme	fest	sólida	Kinko 115	3

	English	français	deutsch	español	Example Varieties/ Exemples/ Beispielssorten/ Variedades ejemplo	Note/ Nota
31.	MS	Fruit body: ratio of cap diameter / stipe length	Organe de fructification : rapport diamètre du chapeau / longueur du stipe	Fruchtkörper: Verhältnis Hutdurchmesser / Stiellänge	Cuerpo frutal: relación entre el diámetro del sombrero y la longitud del pie	
(+)						
QN	(c)	cap far shorter than stipe length	chapeau beaucoup plus court que le stipe	Hut viel kürzer als Stiellänge	sombrero mucho más corto que la longitud del pie Kinko 610	1
		cap shorter than stipe length	chapeau plus court que le stipe	Hut kürzer als Stiellänge	sombrero más corto que la longitud del pie Mori 252	3
		cap almost equal to stipe length	chapeau presque aussi long que le stipe	Hut fast gleich lang wie Stiellänge	sombrero casi igual a la longitud del pie Akiyama A-526, Susono 360	5
		cap longer than stipe length	chapeau plus long que le stipe	Hut länger als Stiellänge	sombrero más largo que la longitud del pie Morino Natsumi	7
		cap far longer than stipe length	chapeau beaucoup plus long que le stipe	Hut viel kürzer als Stiellänge	sombrero mucho más largo que la longitud del pie Morino Harumitsu	9
32.	MS	Fruit body: ratio of cap diameter / stipe diameter	Organe de fructification : rapport diamètre du chapeau / diamètre du stipe	Fruchtkörper: Verhältnis Hutdurchmesser / Stieldurchmesser	Cuerpo frutal: relación entre el diámetro del sombrero y el diámetro del pie	
QN	(c)	cap far shorter than stipe diameter	chapeau beaucoup plus étroit que le stipe	Hut viel kürzer als Stieldurchmesser	sombrero mucho más corto que el diámetro del pie	1
		cap shorter than cap diameter	chapeau plus étroit que le stipe	Hut kürzer als Hutdurchmesser	sombrero más corto que la diámetro del pie ML12	3
		cap almost equal to cap diameter	chapeau presque aussi large que le stipe	Hut fast gleich lang wie Hutdurchmesser	sombrero casi igual al diámetro del pie JHS KV92	5
		cap longer than stipe diameter	chapeau plus large que le stipe	Hut länger als Stieldurchmesser	sombrero más largo que el diámetro del pie Morino Harumitsu	7
		cap far longer than stipe diameter	chapeau beaucoup plus large que le stipe	Hut viel länger als Stieldurchmesser	sombrero mucho más largo que el diámetro del pie Mori 4T98	9

	English	français	deutsch	español	Example Varieties/ Exemples/ Beispielssorten/ Variedades ejemplo	Note/ Nota	
33.	MG	Fruit body: dry weight at harvest maturity	Organe de fructification : poids sec à maturité de récolte	Fruchtkörper: Trockengewicht bei Erntereife	Cuerpo frutal: peso seco en la época de madurez para la cosecha		
(+)							
QN	(c)	light	petit	leicht	ligero	HS73	3
		medium	moyen	mittel	medio	Akiyama A-526, Susono 360, Yujiro	5
		heavy	élevé	schwer	pesado		7
34.	VG	Fruit body: period from inoculation to fruit induction	Organe de fructification : période entre l'inoculation et l'induction du développement des sporophores	Fruchtkörper: Zeitraum von Inokulation bis Fruchtinduktion	Cuerpo frutal: período entre la inoculación y la inducción del fruto		
(*)							
(+)							
QN		short	courte	kurz	breve	A-555(B), HS73(S), Kinko 702(B)	3
		medium	moyenne	mittel	medio	HS608(B), Kinko 697(B), S-035(B),	5
		long	longue	lang	largo	HS705(S), Kinko 169(B), ML8(S) Mori-yujiro(B), S-035(S)	7
35.	VG	Fruit body: period from fruit induction to harvest	Organe de fructification : période entre l'induction du développement des sporophores et la récolte	Fruchtkörper: Zeitraum von Fruchtinduktion bis Ernte	Cuerpo frutal: período entre la inducción del fruto y la cosecha		
(*)							
(+)							
QN		short	courte	kurz	breve	A-555(B), HS-73(S), KX-S055(B), S-005(S)	3
		medium	moyenne	mittel	medio	A-526(B), HS72(S), HS705(S)	5
		long	longue	lang	largo	ML8(B), ML8(S), S-035(S)	7

8. Explanations on the Table of Characteristics

8.1 *Explanations covering several characteristics*

Characteristics containing the following key in the second column of the Table of Characteristics should be examined as indicated below:

- (a) Hyphae and Colony: should be observed by pure cultures.
Type of medium : PDA (Potato dextrose agar)
Plate : 9cm in inside diameter and 2cm in height
Conditions : in the dark at $25 \pm 2^{\circ}\text{C}$
Observations : (hyphae) developed to about 70% of the diameter of the plate
(colony) 14 days after
Number of plate : more than two
- (b) Mycelium: should be observed by pure cultures.
Type of medium : PDA
Tube/Plate : growth tube or 9cm in inside diameter and 2cm in height
Conditions : in the dark at specified temperature
Observations : 14 days after
Number of tube : more than five
- (c) Stipe, cap and gills: Unless otherwise indicated, all characteristics of the fruit bodies (the cap, the stipe and the gills) should be recorded at 80 to 90% open gills (stage 4[see Additional information (a)] handpicked mushrooms; freshly harvested).
- (d) General illustration:



A: cap B: gill C: stipe D: scale

8.2 Explanations for individual characteristics

Ad. 1: Density of hyphae on the medium

Observe the dense of hyphae of the PDA medium (see 8.1 (b)).



Ad. 2: Colony: tinting of surface on the medium




Observe the presence of tinting of colony after 14 days cultivation (see 8.1 (b)).

A: 1 absent
B: 9 present


Ad. 3: Mycelium: optimum temperature for growth

Ad. 4, 5, 6, 7, 8: Mycelium: growth rate at 10°C, 15°C, 20°C, 25°C, 30°C


Measure the length or diameter (see 8.1(b) tube/plate) that grew up on the 14th day from the 4th day of mycelium cultured at each temperature. The incubation temperature of mycelium that grows up the longest becomes an optimum temperature. Amount of mycelium growth per day at each temperature are considered to be the growth rate. These characteristics should be evaluated by drawing of mycelium growth curve (see following graph).



A: diameter of mycelium
B: length of colony



Ad. 9: Cap: shape of vertical section



Ad. 10: Cap: diameter

Ad. 12: Cap: height

Ad. 21: Gill: width

Measure the widest part of the gill.


Ad. 13: Cap: firmness

Determine by hand the hardness of the cap is compared to standard varieties

Ad. 15: Cap: distribution of scale




1
whole



2
periphery

Ad. 16: Cap: size of scale

Observe the size of scales in the part of the shoulder of Cap



3
small


5
medium

7
large

Ad. 17: Cap: tinting of scale




1
absent



9
present

Ad. 18: Cap: presence of gill




1
absent



9
present

Ad. 19: Gill: shape

Observations should be made on the gill located outside of curved stipe.



Ad. 20: Gill: arrangement of row



1
straight



2
ripple or crinkle

Ad. 22: Gill: density of row



1
sparse




2
medium




3
dense


Ad. 24: Stipe: shape in vertical section



1
broader toward cap



2
cylindrical




3
broader toward base

Ad. 25: Stipe: length

Ad. 26: Stipe: diameter


Measure the widest part of the stipe.

- 1 Cap: diameter
- 2 Cap: height
- 3 Gill: width
- 4 Stipe: length
- 5 Stipe: diameter




Ad. 27: Stipe: tinting

Ad. 29: Stipe: tinting of fluff



1
absent



9
present

A: Stipe surface
B: Fluff of stype

Ad. 28: Stipe: density of fluff



1
absent or sparse



2
medium



3
dense

Ad. 30: Stipe: firmness

Determine by hand the hardness of the stipe is compared to standard varieties.

Ad. 33: Fruit body: dry weight at harvest maturity

The fruiting body dries at 60°C until becoming a constant weight.


Ad. 34: Fruit body: Period from inoculation to fruit induction

Ad. 35: Fruit body: Period from fruit induction to harvest

The fruit induction is indispensable for the fruit body development. In the same condition, the timing of the fruit induction is different according to each variety clearly. In the fruit induction, there is water soaking treatment, water sprinkling treatment, low temperature treatment, or physical treatment etc. Any method is stimulation for the fruiting body formation. In each cultivation type (bed-log cultivation, sawdust cultivation), the fruit induction should be treated at the time when the symptom (e.g. appearance of primordium) of the fruiting body formation is shown. The harvesting time is considered to be a peak from which fruit bodies are harvested most.

Additional information:

(a) Stage of fruit body




Explanation: 1 and 2 - veil closed 4 - 80 to 90% open / gills visible
 3 - veil breaking 5 - fully open



stage 4

(b) Life cycle of *Lentinula edodes* (Berk.) Pegler



8.3 Growing Types

As explained in Chapter 3.3.1, it may be necessary for separate growing trials to be established for bed-log cultivation types and sawdust cultivation types in order to ensure the satisfactory growth of varieties of those types. The following information is provided with regard to growing conditions for different types of varieties and information which may help in deciding on the type of trial(s) which may be appropriate for a variety:

Bed-log cultivation types

Breeding is done in a rather large gene pool, in most cases much broader than the sawdust cultivation types. In general, such types of variety have the following features:

- some varieties of this types don't develop the fruit bodies in the condition of the sawdust cultivation;
- mainly used as dried shiitake mushrooms and produced in the bed-log laying yard;
- the cultivation period is very long than the sawdust cultivation types;
- firmness of cap is comparatively hard;

Sawdust cultivation types

Breeding is done in a limited gene pool. In general, such types of variety have the following features:

- some varieties of this types don't develop the fruit bodies in the condition of the bed-log cultivation;
- tolerant of higher temperatures in general;
- mainly used as fresh shiitake mushrooms ;
- concern only types produced in sheltered conditions;
- the cultivation period is comparatively short;
- firmness of cap is comparatively soft;

9. Literature

Pegler, D.N., 1975 (1976): The classification of the genus *Lentinus* Fr. (Basidiomycota),
Kavaka 3:11-20

Ministry of Agriculture, Forestry and Fisheries, 1996: National Test Guideline for Shiitake,
JP

Kirk, P.M., Cannon P.F., Minter D.W. and Stalpers J.A. (eds.), 2008: Dictionary of the Fungi
10th edition, CAB International, UK ISBN 978-0-85199-826-8

10. Technical Questionnaire

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
		Application date: (not to be filled in by the applicant)
TECHNICAL QUESTIONNAIRE to be completed in connection with an application for plant breeders' rights		
1. Subject of the Technical Questionnaire		
1.1 Botanical Name	<input type="text" value="Lentinula edodes (Berk.) Pegler"/>	
1.2 Common Name	<input type="text" value="Shiitake"/>	
2. Applicant		
Name	<input type="text"/>	
Address	<input type="text"/>	
Telephone No.	<input type="text"/>	
Fax No.	<input type="text"/>	
E-mail address	<input type="text"/>	
Breeder (if different from applicant)	<input type="text"/>	
3. Proposed denomination and breeder's reference		
Proposed denomination (if available)	<input type="text"/>	
Breeder's reference	<input type="text"/>	

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

#4. Information on the breeding scheme and propagation of the variety

4.1 Breeding scheme

Variety resulting from:

4.1.1 Crossing

4.1.2 Mutation
(please state parent variety)

4.1.3 Discovery and development
(please state where and when discovered and how developed)

4.1.4 Other
(please provide details)"

4.2 Method of propagating the variety

Authorities may allow certain of this information to be provided in a confidential section of the Technical Questionnaire.

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

5. Characteristics of the variety to be indicated (the number in brackets refers to the corresponding characteristic in Test Guidelines; please mark the note which best corresponds).

Characteristics	Example Varieties	Note
5.1 Growth type		
bed-log cultivation type	Morino Natsumi, Yujiro	1[]
sawdust cultivation type	Hokken 600, Mori XR1	2[]
5.2 Mycelium: growth rate at 20°C (6)		
very slow	Bridge 32, Kinno 1	1[]
slow	Kinko 115	2[]
medium	ML8, Morino Natsumi	3[]
fast	Morino Harumitsu	4[]
very fast	Akiyama A-950, Hokken 600, JMS 237	5[]
5.3 Mycelium: growth rate at 30°C (8)		
very slow	Akiyama A-526	1[]
slow	HS73, Morino Harumitsu	2[]
medium	Kinko 115	3[]
fast	Mori XR1, Susono 360	4[]
very fast	Morino Natsumi, Yujiro	5[]
5.4 Cap: shape of vertical section (9)		
concave	JMS 7H-1	1[]
flat	Morino Harumitsu	2[]
round	Kinko 115, Yujiro	3[]
convex	KX-S005	4[]

TECHNICAL QUESTIONNAIRE		Page {x} of {y}	Reference Number:
Characteristics		Example Varieties	Note
5.5	Cap: diameter		
(10)			
	very small		1[]
	very small to small		2[]
	small	Morino Harumitsu	3[]
	small to medium	Morino Natsumi, Yujiro	4[]
	medium	HS73, Kinko 115, Mori XR1	5[]
	medium to large	Akiyama A-526, HS607,	6[]
	large	Kinko 117, Mori 505	7[]
	large to very large	Kinko 245, Kinko 130	8[]
	very large		9[]
5.6	Cap: main color of apex		
(11)			
	white	Kinko 989	1[]
	yellowish brown	Mori XR-1	2[]
	brown	Kinko 115, Susono 360	3[]
	reddish brown	Akiyama A-526	4[]
5.7	Cap: presence of scale		
(14)			
	absent	KX-S034	1[]
	present	HS73, Kinko 115, ML8, Morino Natsumi	9[]
5.8	Cap: presence of gill		
(18)			
	absent	FERM P-14310	1[]
	present	Kinko 115, Mori XR1	9[]

TECHNICAL QUESTIONNAIRE		Page {x} of {y}	Reference Number:
Characteristics		Example Varieties	Note
5.9	Stipe: shape in vertical section		
(24)			
	broader toward cap	Susono 360	1[]
	cylindrical	JMS5K16, Kinko 115, Mori XR1, Morino Natsumi	2[]
	broader toward base	JMS 7H-1	3[]
5.10	Stipe: length		
(25)			
	very short		1[]
	very short to short	Yujiro, Morino Harumitsu	2[]
	short	Mori XR1, Morino Natsumi, Susono 360	3[]
	short to medium	Kinko 115	4[]
	medium	HS702, Kinko 117	5[]
	medium to long	HS73, KX-S005, HS607	6[]
	long	Akiyama A-526	7[]
	long to very long	HS802	8[]
	very long		9[]
5.11	Stipe: tinting		
(27)			
	absent	Mori XR1	1[]
	present	Kinko 115, KX-S055 Morino Natsumi	9[]
5.12	Zone line of colony: dual culture of mother variety		
			9[]
	absent		1[]
	present	Mori XR1	9[]

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

Characteristics	Example Varieties	Note
5.13 Zone line of colony: dual culture of father variety		
absent		1[]
present	Mori XR1	9[]
5.14 Zone line of colony: dual culture of similar variety		
absent		1[]
present	Mori XR1	9[]
5.15 Fruit body: period from inoculation to fruit induction (34)		
very short		1[]
very short to short		2[]
short	HS73(S), Kinko 702(B), A-555(B)	3[]
short to medium		4[]
medium	HS608(B), S-035(B), Kinko 697(B)	5[]
medium to long		6[]
long	Mori-yujiro(B), HS705(S), S-035(S), ML8(S)	7[]
long to very long		8[]
very long		9[]

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

Characteristics	Example Varieties	Note
5.16 Fruit body: period from fruit induction to harvest (35)		
very short		1[]
very short to short		2[]
short	KX-S055(B), A-555(B), HS-73(S), S-005(S)	3[]
short to medium		4[]
medium	A-526(B), HS72(S), HS705(S)	5[]
medium to long		6[]
long	ML8(B), ML8(S), S-035(S)	7[]
long to very long		8[]
very long		9[]

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

6. Similar varieties and differences from these varieties

Please use the following table and box for comments to provide information on how your candidate variety differs from the variety (or varieties) which, to the best of your knowledge, is (or are) most similar. This information may help the examination authority to conduct its examination of distinctness in a more efficient way.

Denomination(s) of variety(ies) similar to your candidate variety	Characteristic(s) in which your candidate variety differs from the similar variety(ies)	Describe the expression of the characteristic(s) for the similar variety(ies)	Describe the expression of the characteristic(s) for your candidate variety
<i>Example</i>	<i>Cap: shape of vertical section</i>	<i>convex</i>	<i>flat</i>

Comments:

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
<p>#7. Additional information which may help in the examination of the variety</p> <p>7.1 In addition to the information provided in sections 5 and 6, are there any additional characteristics which may help to distinguish the variety?</p> <p>Yes [] No []</p> <p>(If yes, please provide details)</p> <p>7.2 Are there any special conditions for growing the variety or conducting the examination?</p> <p>Yes [] No []</p> <p>(If yes, please provide details)</p> <p>7.3 Other information</p> <p>7.3.1 Type of cultivation</p> <p>(a) bed-log cultivation []</p> <p>(b) sawdust cultivation []</p>		
<p>8. Authorization for release</p> <p>(a) Does the variety require prior authorization for release under legislation concerning the protection of the environment, human and animal health?</p> <p>Yes [] No []</p> <p>(b) Has such authorization been obtained?</p> <p>Yes [] No []</p> <p>If the answer to (b) is yes, please attach a copy of the authorization.</p>		

Authorities may allow certain of this information to be provided in a confidential section of the Technical Questionnaire.

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

9. Information on plant material to be examined or submitted for examination.

9.1 The expression of a characteristic or several characteristics of a variety may be affected by factors, such as pests and disease, chemical treatment (e.g. growth retardants or pesticides), effects of tissue culture, different rootstocks, scions taken from different growth phases of a tree, etc.

9.2 The material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If the plant material has undergone such treatment, full details of the treatment must be given. In this respect, please indicate below, to the best of your knowledge, if the plant material to be examined has been subjected to:

- | | | |
|---|---------|--------|
| (a) Microorganisms (e.g. virus, bacteria, phytoplasma) | Yes [] | No [] |
| (b) Chemical treatment (e.g. growth retardant, pesticide) | Yes [] | No [] |
| (c) Tissue culture | Yes [] | No [] |
| (d) Other factors | Yes [] | No [] |

Please provide details for where you have indicated “yes”.

.....

10. I hereby declare that, to the best of my knowledge, the information provided in this form is correct:

Applicant's name

Signature

Date

[End of document]