



TG/BRASS_JUN(proj.6)

ORIGINAL: English

DATE: 2017-10-17

INTERNATIONAL UNION FOR THE PROTECTION OF NEW VARIETIES OF PLANTS

Geneva

DRAFT

BROWN MUSTARD

UPOV Code(s):

BRASS_JUN

*Brassica juncea (L.) Czern.***GUIDELINES****FOR THE CONDUCT OF TESTS****FOR DISTINCTNESS, UNIFORMITY AND STABILITY**

*prepared by experts from Japan
 to be considered by the
 Enlarged Editorial Committee
 at its meeting, to be held in Geneva
 from 2018-03-26 to 2018-03-27*

Disclaimer: this document does not represent UPOV policies or guidance

Alternative names:^{*}

Botanical name	English	French	German	Spanish
<i>Brassica juncea (L.) Czern.</i>	Brown mustard, Indian mustard, Oriental mustard	Moutarde brune	Sareptasenf	Mostaza de Sarepta, Mostaza india

The purpose of these guidelines (“Test Guidelines”) is to elaborate the principles contained in the General Introduction (document TG/1/3), and its associated TGP documents, into detailed practical guidance for the harmonized examination of distinctness, uniformity and stability (DUS) and, in particular, to identify appropriate characteristics for the examination of DUS and production of harmonized variety descriptions.

ASSOCIATED DOCUMENTS

These Test Guidelines should be read in conjunction with the General Introduction and its associated TGP documents.

* These names were correct at the time of the introduction of these Test Guidelines but may be revised or updated. [Readers are advised to consult the UPOV Code, which can be found on the UPOV Website (www.upov.int), for the latest information.]

TABLE OF CONTENTS	PAGE
1. SUBJECT OF THESE TEST GUIDELINES.....	4
2. MATERIAL REQUIRED.....	4
3. METHOD OF EXAMINATION.....	5
3.1 Number of Growing Cycles.....	5
3.2 Testing Place.....	5
3.3 Conditions for Conducting the Examination.....	5
3.4 Test Design.....	5
3.5 Additional Tests.....	5
4. ASSESSMENT OF DISTINCTNESS, UNIFORMITY AND STABILITY.....	6
4.1 Distinctness.....	6
4.2 Uniformity.....	7
4.3 Stability.....	7
5. GROUPING OF VARIETIES AND ORGANIZATION OF THE GROWING TRIAL.....	8
6. INTRODUCTION TO THE TABLE OF CHARACTERISTICS.....	11
6.1 Categories of Characteristics.....	11
6.2 States of Expression and Corresponding Notes.....	11
6.3 Types of Expression.....	11
6.4 Example Varieties.....	11
6.5 Legend.....	12
7. TABLE OF CHARACTERISTICS/TABLEAU DES CARACTÈRES/MERKMALSTABELLE/TABLA DE CARACTERES.....	13
8. EXPLANATIONS ON THE TABLE OF CHARACTERISTICS.....	22
8.1 Explanations covering several characteristics.....	22
8.2 Explanations for individual characteristics.....	23
9. LITERATURE.....	36
10. TECHNICAL QUESTIONNAIRE.....	37

1. Subject of these Test Guidelines

These Test Guidelines apply to all varieties of *Brassica juncea* (L.) Czern..

2. Material Required

- 2.1 The competent authorities decide on the quantity and quality of the plant material required for testing the variety and when and where it is to be delivered. Applicants submitting material from a State other than that in which the testing takes place must ensure that all customs formalities and phytosanitary requirements are complied with.
- 2.2 The material is to be supplied in the form of seed.
- 2.3 The minimum quantity of plant material, to be supplied by the applicant, should be:

3,000 seeds for single spaced plants.
20,000 seeds for drilled plants.

The seed should meet the minimum requirements for germination, species and analytical purity, health and moisture content, specified by the competent authority. In cases where the seed is to be stored, the germination capacity should be as high as possible and should, be stated by the applicant.

- 2.4 The plant material supplied should be visibly healthy, not lacking in vigor, nor affected by any important pest or disease.
- 2.5 The plant material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If it has been treated, full details of the treatment must be given.

3. Method of Examination

3.1 *Number of Growing Cycles*

The minimum duration of tests should normally be two independent growing cycles.

3.2 *Testing Place*

Tests are normally conducted at one place. In the case of tests conducted at more than one place, guidance is provided in TGP/9 "Examining Distinctness".

3.3 *Conditions for Conducting the Examination*

- 3.3.1 The tests should be carried out under conditions ensuring satisfactory growth for the expression of the relevant characteristics of the variety and for the conduct of the examination.
- 3.3.2 The optimum stage of development for the assessment of each characteristic is indicated by a number in the Table of Characteristics. The stages of development denoted by each number are described in Chapter 8.

3.4 *Test Design*

- 3.4.1 In the case of single spaced plants, each test should be designed to result in a total of at least 60 plants which should be divided between at least 2 replicates.
- 3.4.2 In the case of drilled plots, each test should be designed to result in a total of at least 220 plants which should be divided between at least 2 replicates.

3.5 *Additional Tests*

Additional tests, for examining relevant characteristics, may be established.

4. Assessment of Distinctness, Uniformity and Stability

4.1 *Distinctness*

4.1.1 General Recommendations

It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding distinctness. However, the following points are provided for elaboration or emphasis in these Test Guidelines.

4.1.2 Consistent Differences

The differences observed between varieties may be so clear that more than one growing cycle is not necessary. In addition, in some circumstances, the influence of the environment is not such that more than a single growing cycle is required to provide assurance that the differences observed between varieties are sufficiently consistent. One means of ensuring that a difference in a characteristic, observed in a growing trial, is sufficiently consistent is to examine the characteristic in at least two independent growing cycles.

4.1.3 Clear Differences

Determining whether a difference between two varieties is clear depends on many factors, and should consider, in particular, the type of expression of the characteristic being examined, i.e. whether it is expressed in a qualitative, quantitative, or pseudo-qualitative manner. Therefore, it is important that users of these Test Guidelines are familiar with the recommendations contained in the General Introduction prior to making decisions regarding distinctness.

4.1.4 Number of Plants or Parts of Plants to be Examined

Unless otherwise indicated, for the purposes of distinctness, all observations on single plants should be made on 30 plants or parts of plants taken from each of 30 plants and any other observations made on all plants in the test, disregarding any off-type plants.

4.1.5 Method of Observation

The recommended method of observing the characteristic for the purposes of distinctness is indicated by the following key in the Table of Characteristics (see document TGP/9 “Examining Distinctness”, Section 4 “Observation of characteristics”):

MG: single measurement of a group of plants or parts of plants

MS: measurement of a number of individual plants or parts of plants

VG: visual assessment by a single observation of a group of plants or parts of plants

VS: visual assessment by observation of individual plants or parts of plants

Type of observation: visual (V) or measurement (M)

“Visual” observation (V) is an observation made on the basis of the expert’s judgment. For the purposes of this document, “visual” observation refers to the sensory observations of the experts and, therefore, also includes smell, taste and touch. Visual observation includes observations where the expert uses reference points (e.g. diagrams, example varieties, side-by-side comparison) or non-linear charts (e.g. color charts). Measurement (M) is an objective observation against a calibrated, linear scale e.g. using a ruler, weighing scales, colorimeter, dates, counts, etc.

Type of record: for a group of plants (G) or for single, individual plants (S)

For the purposes of distinctness, observations may be recorded as a single record for a group of plants or parts of plants (G), or may be recorded as records for a number of single, individual plants or parts of plants (S). In most cases, “G” provides a single record per variety and it is not possible or necessary to apply statistical methods in a plant-by-plant analysis for the assessment of distinctness.

In cases where more than one method of observing the characteristic is indicated in the Table of Characteristics (e.g. VG/MG), guidance on selecting an appropriate method is provided in document TGP/9, Section 4.2.

4.2 *Uniformity*

- 4.2.1 It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding uniformity. However, the following points are provided for elaboration or emphasis in these Test Guidelines:
- 4.2.2 These Test Guidelines have been developed for the examination of seed-propagated varieties. For varieties with other types of propagation, the recommendations in the General Introduction and document TGP/13 "Guidance for new types and species" Section 4.5 "Testing Uniformity" should be followed.
- 4.2.3 The assessment of uniformity for cross-pollinated varieties should be according to the recommendations for cross-pollinated varieties in the General Introduction.
- 4.2.4 For the assessment of uniformity of seed-propagated varieties, a population standard of 2% and an acceptance probability of at least 95% should be applied. In the case of a sample size of 60 plants, 3 off-types are allowed. In the case of a sample size of 200 plants for drilled plants, 7 off-types are allowed.

4.3 *Stability*

- 4.3.1 In practice, it is not usual to perform tests of stability that produce results as certain as those of the testing of distinctness and uniformity. However, experience has demonstrated that, for many types of variety, when a variety has been shown to be uniform, it can also be considered to be stable.
- 4.3.2 Where appropriate, or in cases of doubt, stability may be further examined by testing a new seed stock to ensure that it exhibits the same characteristics as those shown by the initial material supplied.

5. Grouping of Varieties and Organization of the Growing Trial


- 5.1 The selection of varieties of common knowledge to be grown in the trial with the candidate varieties and the way in which these varieties are divided into groups to facilitate the assessment of distinctness are aided by the use of grouping characteristics.
- 5.2 Grouping characteristics are those in which the documented states of expression, even where produced at different locations, can be used, either individually or in combination with other such characteristics: (a) to select varieties of common knowledge that can be excluded from the growing trial used for examination of distinctness; and (b) to organize the growing trial so that similar varieties are grouped together.
- 5.3 The following have been agreed as useful grouping characteristics:
 - (a) Seed: color (characteristic 1)
 - (b) Leaf blade: density of incisions of margin (characteristic 17)
 - (c) Leaf blade: blistering (characteristic 18)
 - (d) Plant: head formation (characteristic 20)

In the first place, the collection should be divided according to leaf types in the following Table.
 In case of doubt to which type a variety belongs to, it should be tested in all relevant types.


Leaf: type	Example varieties	Only varieties with Leaf: type: type 1 or 2: Leaf blade: size of terminal lobe (char. 11)	Leaf blade: number of lateral lobes (char. 12)	Leaf blade: blistering (char. 18)	Only varieties with Leaf: type: type 3 and 4: Leaf blade: width of midrib (char. 19)	Plant: head formation (Char. 20)
Type 1	Hagarashina, Kigarashina, Terrafit	medium to very large	few to medium	absent or weak to medium	-	absent
Type 2	Akariasu, Flaming Frills, Riasu Karashina, Scarlet Frills	very small to small	many to very many	-	-	absent
Type 3	Akaoba Takana, Sagami Green	-	absent or very few	medium to strong	narrow	absent
Type 4	Kekkyu Takana, Miike Takana, Shinkoku Seisai	-	absent or very few	medium to strong	medium to broad	absent or present




Type 1
(lyrate)



Type 2
(division)



Type 3
(entire and narrow midrib)



Type 4
(entire and broad midrib)

- 5.4 Guidance for the use of grouping characteristics, in the process of examining distinctness, is provided through the General Introduction and document TGP/9 "Examining Distinctness".

6. Introduction to the Table of Characteristics

6.1 *Categories of Characteristics*

6.1.1 Standard Test Guidelines Characteristics

Standard Test Guidelines characteristics are those which are approved by UPOV for examination of DUS and from which members of the Union can select those suitable for their particular circumstances.

6.1.2 Asterisked Characteristics

Asterisked characteristics (denoted by *) are those included in the Test Guidelines which are important for the international harmonization of variety descriptions and should always be examined for DUS and included in the variety description by all members of the Union, except when the state of expression of a preceding characteristic or regional environmental conditions render this inappropriate.

6.2 *States of Expression and Corresponding Notes*

6.2.1 States of expression are given for each characteristic to define the characteristic and to harmonize descriptions. Each state of expression is allocated a corresponding numerical note for ease of recording of data and for the production and exchange of the description.

6.2.2 In the case of qualitative and pseudo-qualitative characteristics (see Chapter 6.3), all relevant states of expression are presented in the characteristic. However, in the case of quantitative characteristics with 5 or more states, an abbreviated scale may be used to minimize the size of the Table of Characteristics. For example, in the case of a quantitative characteristic with 9 states, the presentation of states of expression in the Test Guidelines may be abbreviated as follows:

State	Note
small	3
medium	5
large	7

However, it should be noted that all of the following 9 states of expression exist to describe varieties and should be used as appropriate:

State	Note
very small	1
very small to small	2
small	3
small to medium	4
medium	5
medium to large	6
large	7
large to very large	8
very large	9

6.2.3 Further explanation of the presentation of states of expression and notes is provided in document TGP/7 "Development of Test Guidelines".

6.3 *Types of Expression*

An explanation of the types of expression of characteristics (qualitative, quantitative and pseudo-qualitative) is provided in the General Introduction.

6.4 *Example Varieties*

Where appropriate, example varieties are provided to clarify the states of expression of each characteristic.

6.5 Legend

	English		français		deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
1	2	3	4	5	6	7		
	Name of characteristics in English	Nom du caractère en français		Name des Merkmals auf Deutsch		Nombre del carácter en español		
	states of expression	types d'expression		Ausprägungsstufen		tipos de expresión		

- 1 Characteristic number
- 2 (*) Asterisked characteristic – see Chapter 6.1.2
- 3 Type of expression

QL	Qualitative characteristic	– see Chapter 6.3
QN	Quantitative characteristic	– see Chapter 6.3
PQ	Pseudo-qualitative characteristic	– see Chapter 6.3
- 4 Method of observation (and type of plot, if applicable)

MG, MS, VG, VS	– see Chapter 4.1.5
----------------	---------------------
- 5 (+) See Explanations on the Table of Characteristics in Chapter 8.2
- 6 (a)
- 7 Growth stage key See Explanations on the Table of Characteristics in Chapter 8

7. Table of Characteristics/Tableau des caractères/Merkmalstabelle/Tabla de caracteres

	English		français		deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
1.	(*)	QL	VG			00		
	Seed: color		Graine : couleur		Samen: Farbe	Semilla: color		
	yellow		jaune		gelb	amarillo	Kigarashina	1
	blackish brown		brun noirâtre		schwärzlich braun	marrón negruzco	Akaoba Takana, Esperance, Miike Takana, Terrafit, Terraplus	2
	2.	QN	VG			10		
	Hypocotyl: anthocyanin coloration		Hypocotyle : pigmentation anthocyanique		Hypokotyl: Anthocyanfärbung	Hipocótilo: pigmentación antociánica		
	absent or weak		nulle ou faible		fehlend oder gering	ausente o débil	Jarangi, TTK456, Zasai FM-58	1
	medium		moyenne		mittel	media	Jarami, Shinkoku Seisai	2
	strong		forte		stark	fuerte	Kigarashina	3
3.	QN	MS/VG	(+)		10			
	Cotyledon: length		Cotylédon : longueur		Keimblatt: Länge	Cotiledón: longitud		
	short		court		kurz	corto	Junkei Yamashiona, Vittasso	3
	medium		moyen		mittel	medio	Katsuona, Terraplus	5
	long		long		lang	largo	Scala	7
4.	QN	MS/VG	(+)		10			
	Cotyledon: width		Cotylédon : largeur		Keimblatt: Breite	Cotiledón: anchura		
	narrow		étroit		schmal	estrecho	Junkei Yamashiona, Vittasso	3
	medium		moyen		mittel	medio	Katsuona, Pacific Gold, Terraplus	5
	broad		large		breit	ancho	Minaret, Terminator	7

	English		français		deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
5.	PQ	VG	(+)	(a)	19			
Leaf: shape	Leaf: shape		Feuille : forme		Blatt: Form	Hoja: forma		
	ovate		ovale		eiförmig	oval	Serihon	1
	circular		circulaire		rund	circular	Kekkyu Takana	2
	elliptic		elliptique		elliptisch	elíptica	Akariasu	3
	oblong		oblongue		länglich	oblonga	Etamine, Zasai FM-58	4
	obovate		obovale		verkehrt eiförmig	oboval	Esperance, Katsuona	5
	spatulate		spatulée		spatelförmig	espatulada	Kigarashina	6
6.	QN	VG	(+)	(a)	19			
Leaf: attitude	Leaf: attitude		Feuille : port		Blatt: Haltung	Hoja: porte		
	erect		dressée		aufrecht	erecta	Energy, Vittasso, Wasabina	1
	semi-erect		demi-dressée		halbaufrecht	semierecta	Esperance, Shinkoku Seisai	3
	horizontal		horizontale		waagerecht	horizontal	Etamine, Miike Takana	5
7.	QN	MS/VG	(+)	(a)	19			
Leaf: length	Leaf: length		Feuille : longueur		Blatt: Länge	Hoja: longitud		
	short		courte		kurz	corta	Chirimen Hakarashina	3
	medium		moyenne		mittel	media	Miike Takana, Terraplus	5
	long		longue		lang	larga	Akaoba Takana, Vittasso	7
8.	QN	MS/VG	(+)	(a)	19			
Leaf: width	Leaf: width		Feuille : largeur		Blatt: Breite	Hoja: anchura		
	narrow		étroite		schmal	estrecha	Chirimen Hakarashina	3
	medium		moyenne		mittel	media	Miike Takana, Terraplus	5
	broad		large		breit	ancha	Katsuona, Vittasso	7

	English		français		deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
9. (*)	QN	MS/VG	(+)	(a)	19			
	Leaf: length of petiole		Feuille : longueur du pétiole		Blatt: Länge des Blattstiels	Hoja: longitud del pecíolo		
	absent or very short		absent ou très court		fehlend oder sehr kurz	ausente o muy corto	Serihon	1
	short		court		kurz	corta	Miike Takana	3
	medium		moyen		mittel	media	Junkei Yamashiona	5
	long		long		lang	largo		7
10.	QN	MS/VG	(+)	(a)	19			
	Leaf: width of petiole		Feuille : largeur du pétiole		Blatt: Breite des Blattstiels	Hoja: anchura del pecíolo		
	narrow		étroit		schmal	estrecho	Kigarashina	3
	medium		moyen		mittel	medio	Katsuona	5
	broad		large		breit	ancho	Shinkoku Seisai	7
11.	QN	VG	(+)	(a)	19			
	<u>Only varieties with Leaf: type: type 1 or 2:</u> <u>Leaf blade: size of terminal lobe</u>		<u>Seulement les variétés avec Feuille : type : type 1 ou 2 : Limbe : taille du lobe terminal</u>		<u>Nur Sorten mit Blatt: Typ: Typ 1 oder 2:: Blattspreite: Größe des Endlappens</u>	<u>Solo variedades con Hoja: tipo: tipo 1 o 2: Limbo: tamaño del lóbulo terminal</u>		
	small		petit		klein	pequeño	Akariasu	3
	medium		moyen		mittel	medio	Kigarashina	5
	large		large		groß	grande	Pacific Gold, Perm Green	7
12. (*)	QN	VG	(+)	(a)	19			
	Leaf blade: number of lateral lobes		Limbe : nombre de lobes latéraux		Blattspreite: Anzahl der Seitenlappen	Limbo: número de lóbulos laterales		
	absent or very few		nul ou très petit		fehlend oder sehr wenige	ausentes o muy bajo		1
	few		petit		wenige	bajo	Minaret	3
	medium		moyen		mittel	medio	Esperance, Kigarashina	5
	many		grand		viele	alto	Akariasu, TTK456	7

	English		français		deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
13.	QN	VG	(a)		19			
	Leaf blade: pubescence on lower side		Limbe : pubescence sur la face inférieure		Blattspreite: Behaarung der Unterseite	Limbo: pubescencia en envés		
	absent or weak		nulle ou faible		fehlend oder gering	ausente o débil	Miike Takana	1
	medium		moyenne		mittel	media	Oba Takana	2
	strong		forte		stark	densa	Kigarashina	3
14. (*)	QN	VG	(+)	(a)	19			
	Leaf blade: anthocyanin coloration		Limbe : pigmentation anthocyanique		Blattspreite: Anthocyanfärbung	Limbo: pigmentación antociánica		
	absent or very weak		nulle ou très faible		fehlend oder sehr gering	ausente o muy débil	Kekkyu Takana, Vitamine	1
	weak		faible		gering	débil		3
	medium		moyenne		mittel	media	Miike Takana	5
	strong		forte		stark	fuerte	TTK456	7
15.	QN	VG		(a)	19			
	<u>Only varieties with anthocyanin coloration: absent or very weak:</u> Leaf blade: intensity of green color		<u>Seulement les variétés avec pigmentation anthocyanique : nulle ou très faible :</u> Limbe : intensité de la couleur verte		<u>Nur Sorten mit Anthocyanfärbung: fehlend oder sehr gering:</u> Blattspreite: Intensität der grünen Farbe	<u>Solo variedades con pigmentación antociánica: ausente o muy débil:</u> Limbo: intensidad del color verde		
	light		claire		hell	claro	Wasabina	3
	medium		moyenne		mittel	medio	Etamine, Golden Streaks, Katsuona	5
	dark		foncée		dunkel	oscuro	Terratop	7
	16.		QN	VG	(+)	(a)	19	
	Leaf blade: undulation of margin		Limbe : ondulation du bord		Blattspreite: Wellung des Randes	Limbo: ondulación del borde		
	absent or very weak		nulle ou très faible		fehlend oder sehr gering	ausente o muy débil		1
	weak		faible		gering	débil	Akaoba Takana	2
	medium		moyenne		mittel	media	Katsuona	3
	strong		forte		stark	fuerte	Chirimen Hakarashina	4
	very strong		très forte		sehr stark	muy fuerte		5

	English		français		deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
17.	(*)	QN	VG	(+)	(a)	19		
	Leaf blade: density of incisions of margin		Limbe : densité des incisions du bord		Blattspreite: Dichte der Einschnitte	Limbo: densidad de las incisiones del borde		
	absent or very sparse		nulle ou très faible		fehlend oder sehr locker	ausentes o muy laxas	1	
	sparse		faible		locker	laxas	Etamine, Katsuona	
	medium		moyenne		mittel	medias	Opaleska	
	dense		forte		dicht	densas	Oportuna	
18.	(*)	QN	VG	(+)	(a)	19		
	Leaf blade: blistering		Limbe : cloûture		Blattspreite: Blasigkeit	Limbo: ampollado		
	absent or weak		absente ou faible		fehlend oder schwach	ausente o débil	Etamine, Kigarashina	
	medium		moyenne		mittel	medio	Akaoba Takana	
	strong		forte		stark	fuerte	Katsuona	
19.		QN	MS/VG	(+)	(a)	19		
	<u>Only varieties with leaf: type: type 3 and 4: Leaf blade: width of midrib</u>		<u>Seulement les variétés avec Feuille : type : type 3 ou 4 : Limbe : largeur de la nervure médiane</u>		Nur Sorten mit Blatt: Typ: Typ 3 oder 4: Blattspreite: Breite der Mittelrippe	Solo variedades con Hoja: tipo: tipo 3 o 4; Limbo: anchura del nervio central		
	narrow		étroite		schmal	estrecho	Sagami Green	
	medium		moyenne		mittel	medio	Katsuona	
	broad		large		breit	ancho	Shinkoku Seisai	
20.	(*)	QL	VG	(+)		19		
	Plant: head formation		Plante : formation d'un capitule		Pfanze: Kopfbildung	Planta: formación de repollo		
	absent		absente		fehlend	ausente	Kigarashina	
	present		présente		vorhanden	presente	Kekkyu Takana	

	English		français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
21.	QN	MS/VG		19			
	Head: height		Capitule : hauteur	Kopf: Höhe	Repollo: altura		
	short		bas	kurz	corto		1
	medium		moyen	mittel	medio	Kekkyu Takana, Unzen Kekkyu Takana	2
	tall		haut	hoch	alto		3
22.	QN	MS/VG		19			
	Head: width		Capitule : largeur	Kopf: Breite	Repollo: anchura		
	narrow		étroit	schmal	estrecho		1
	medium		moyen	mittel	medio	Kekkyu Takana, Unzen Kekkyu Takana	2
	broad		large	breit	ancho		3
23.	QN	MS/VG		19			
	Head: number of leaves		Capitule : nombre de feuilles	Kopf: Anzahl Blätter	Repollo: número de hojas		
	few		petit	wenige	bajo		3
	medium		moyen	mittel	medio	Kekkyu Takana, Unzen Kekkyu Takana	5
	many		grand	viele	alto		7
24.	PQ	VG		19			
	Head: internal color		Capitule : couleur intérieure	Kopf: Innenfarbe	Repollo: color interno		
	yellowish white		blanc jaunâtre	gelblich weiß	blanco amarillento	Unzen Kekkyu Takana	1
	light green		vert clair	hellgrün	verde claro		2
	medium green		vert moyen	mittelgrün	verde medio	Kekkyu Takana	3

	English		français		deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
25.	PQ	VG	(+)		20-29			
	Main stem: shape		Tige principale : forme		Haupttrieb: Form	Tallo principal: forma		
	narrow conical		conique étroite		schmal kegelförmig	cónico estrecho	Kigarashina	1
	rounded		arrondie		abgerundet	redondeado	Umino	2
	broad conical		conique large		breit kegelförmig	cónico ancho	Zasai FM-58	3
	branched		ramifiée		verzweigt	ramificado	FE-K226	4
26.	QN	MG			31			
	Time of beginning of bolting		Époque de début de montaison		Zeitpunkt des Schossbeginns	Época del comienzo de la floración		
	early		précoce		früh	temprana	Junkei Yamashiona, Scala	3
	medium		moyenne		mittel	media	Terraplus	5
	late		tardive		spät	tardía	Akaoba Takana	7
	27.	QN	MG/MS		50			
	Time of flowering		Époque de floraison		Zeitpunkt der Blüte	Época de floración		
	early		précoce		früh	temprana	Terraflit	3
	medium		moyenne		mittel	media	Minaret, Terraplus	5
	late		tardive		spät	tardía	Brons	7
	very late		très tardive		sehr spät	muy tardía	Vittasso	9
28.	QN	MS/VG	(+)		65-79			
	Only varieties with head formation: absent: Plant: length		Seulement les variétés avec formation d'un capitule : absente : Plante : longueur		Nur Sorten mit Kopfbildung: fehlend: Pflanze: Länge	Solo variedades con formación de repollo: ausente: Planta: longitud		
	short		courte		kurz	baja	Pacific Gold, Terminator	3
	medium		moyenne		mittel	media	Terraplus	5
	long		longue		lang	alta	Minaret	7
	very long		très longue		sehr lang	muy alta	Vittasso	9

	English		français		deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
29.	QN	MS/VG	(+)		65-79			
	<u>Only varieties with head formation:</u> <u>absent: Silique: length</u>		<u>Seulement les variétés avec formation d'un capitule : absente : Silique : longueur</u>		Nur Sorten mit Kopfbildung: fehlend: Schote: Länge	Solo variedades con formación de repollo: ausente: Silicua: longitud		
	short		courte		kurz	corta	Terraplus, Vittasso	3
	medium		moyenne		mittel	media	Pacific Gold	5
	long		longue		lang	larga	Minaret	7
30.	QN	MS/VG	(+)		65-79			
	<u>Only varieties with head formation:</u> <u>absent: Silique: length of beak</u>		<u>Seulement les variétés avec formation d'un capitule : absente : Silique : longueur du bec</u>		Nur Sorten mit Kopfbildung: fehlend: Schote: Länge des Schnabels	Solo en variedades con formación de repollo: ausente: Silicua: longitud de la punta		
	short		court		kurz	corta	Terraplus, Vittasso	3
	medium		moyen		mittel	media	Terraflit	5
	long		long		lang	larga		7
31.	QN	MS/VG	(+)		65-79			
	<u>Only varieties with head formation:</u> <u>absent: Silique: width</u>		<u>Seulement les variétés avec formation d'un capitule : absente : Silique : largeur</u>		Nur Sorten mit Kopfbildung: fehlend: Schote: Breite	Solo en variedades con formación de repollo: ausente: Silicua: anchura		
	narrow		étroite		schmal	estrecha	Vittasso	3
	medium		moyenne		mittel	media	Energy, Terraflit	5
	broad		large		breit	ancha	Oba Takana	7
32.	QN	MS/VG	(+)		65-79			
	<u>Only varieties with head formation:</u> <u>absent: Silique: length of peduncle</u>		<u>Seulement les variétés avec formation d'un capitule : absente : Silique : longueur du pédoncule</u>		Nur Sorten mit Kopfbildung: fehlend: Schote: Länge des Blütenstandstiels	Solo en variedades con formación de repollo: ausente: Silicua: longitud del pedúnculo		
	short		court		kurz	corto	Vittasso	3
	medium		moyen		mittel	medio	Energy	5
	long		long		lang	largo	Minaret	7

	English		français		deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
33.	QN	VG	(+)					
	Tendency to form inflorescences in the year of sowing under long day conditions		Tendance à former des inflorescences dans l'année du semis en jours longs		Neigung zur Bildung von Blütenständen im Jahr der Aussaat unter Langtagsbedingungen	Tendencia a formar inflorescencias en el año de la siembra en condiciones de días largos		
	absent or very weak	nulle ou très faible		fehlend oder sehr gering		ausente o muy débil	Brons, Vittasso	1
	weak	faible		gering		débil		3
	medium	moyenne		mittel		media	Terraplus	5
	strong	forte		stark		fuerte		7
	very strong	très forte		sehr stark		muy fuerte	Energy, Minaret, Terrafit	9

8. Explanations on the Table of Characteristics

8.1 *Explanations covering several characteristics*


Characteristics containing the following key in the Table of Characteristics should be examined as indicated below:

- (a) Observations on the leaves should be made on the largest fully developed (non-senescent) leaf.

8.2 *Explanations for individual characteristics*

Ad. 3: Cotyledon: length

The measurements should be made on cotyledons of 30 seedlings. If the two cotyledons differ in size, the biggest one should be measured. The length is defined as distance between the inclination at top of the cotyledon and the point where the width of the petiole is about 4 mm. The width of the cotyledon should be measured at the widest point of the cotyledons.





a = Cotyledon: width (characteristic 4)

b = Cotyledon: length (characteristic 3)

Ad. 4: Cotyledon: width

See Ad. 3


Ad. 5: Leaf: shape

width (ratio length/width)	← broadest part →		
	below middle	at middle	above middle
narrow (high)			
		4 oblong	6 spatulate
medium (medium)			
	1 ovate	3 elliptic	5 obovate
broad (low)			
		2 circular	

Ad. 6: Leaf: attitude



Ad. 7: Leaf: length



a = Leaf: width (characteristic 8)
b = Leaf: length (characteristic 7)
c = Petiole: length (characteristic 9)
d = Petiole: width (characteristic 10)
e = Midrib: width (characteristic 19)

Ad. 8: Leaf: width

See Ad. 7

Ad. 9: Leaf: length of petiole

See Ad. 7

Ad. 10: Leaf: width of petiole


See Ad. 7

Ad. 11: Only varieties with Leaf: type: type 1 or 2: Leaf blade: size of terminal lobe


Parts of the leaf blade are considered as lobes if their length is at least equivalent to the width of the leaf petiole at their point of attachment and if the upper notch of the blade has at least half the length of the lobe itself.

The terminal lobe is the top lobe of the leaf, which is the No.1 lobe in the following figure. In the case of Type2 leaf, the shape of terminal lobe is similar to the shape of near other lobes.

The lateral lobes are the lobes excluding the terminal lobe.




Type 1



Type 2

The size of terminal lobe should be assessed by the ratio of the terminal lobe size/the leaf size. The terminal lobe size and the leaf size are the size of the area which was surrounded by each outline of them.



a = terminal lobe size
b = leaf size

Ad. 12: Leaf blade: number of lateral lobes


See Ad. 11

Ad. 14: Leaf blade: anthocyanin coloration

The strongest intensity of anthocyanin should be observed (not the extension).


Ad. 16: Leaf blade: undulation of margin

Observations should be made excluding type 2.



Ad. 17: Leaf blade: density of incisions of margin

Observations should be made on the distal part of the leaves, excluding type 2.



Ad. 18: Leaf blade: blistering

Observations should be made excluding type 2.



Ad. 19: Only varieties with leaf: type: type 3 and 4: Leaf blade: width of midrib

See Ad. 7

The width of midrib should be measured at the widest point.

Ad. 20: Plant: head formation



Ad. 25: Main stem: shape

Observations on the shape of the main stem should be made after removing the leaves, excluding lateral stems which are located at the base of main stem.



1
narrow conical



2
rounded



3
broad conical




4
branched

Ad. 28: Only varieties with head formation: absent: Plant: length

Observations should be made when the growth stagnates by observing the total plant height from soil to highest point of the plant.

Ad. 29: Only varieties with head formation: absent: Siliques: length

All observations on the siliques should be recorded in the midpart of the inflorescence of the main stem.



a = Siliques: width (characteristic 31)

b = Siliques: length of beak (characteristic 30)

c = Siliques: length (characteristic 29)

d = Siliques: length of peduncle (characteristic 32)

Observations should be made on the length of the siliques from attachment of peduncle to top, excluding beak.

Ad. 30: Only varieties with head formation: absent: Siliques: length of beak

See Ad. 29

Ad. 31: Only varieties with head formation: absent: Siliques: width

See Ad. 29

Ad. 32: Only varieties with head formation: absent: Siliques: length of peduncle

See Ad. 29

Ad. 33: Tendency to form inflorescences in the year of sowing under long day conditions

The observation of the tendency to form inflorescence (proportion of plants below bud stage, in bud stage, in flowering stage, in stage of siliques formation) should be made in autumn, when the development stagnates.

Alternatively the beginning of flowering may be observed in this trial; early flowering would mean strong tendency, late flowering would mean weak tendency.

8.3 KEY FOR THE STAGE OF DEVELOPMENT

KEY GENERAL DESCRIPTION

0 Principal growth stage 0: Germination

- 01 Beginning of seed imbibition
- 03 Seed imbibition complete
- 05 Radicle emerged from seed
- 07 Hypocotyl with cotyledons emerged from seed
- 08 Hypocotyl with cotyledons growing towards soil surface
- 09 Emergence: cotyledons emerge through soil surface

1 Principal growth stage 1: Leaf development

- 10 Cotyledons completely unfolded
- 11 First leaf unfolded
- 12 2 leaves unfolded
- 13 3 leaves unfolded
- 14 4 leaves unfolded
- 15 5 leaves unfolded
- 16 6 leaves unfolded
- 17 7 leaves unfolded
- 18 8 leaves unfolded
- 19 9 or more leaves unfolded

2 Principal growth stage 2: Formation of side shoots

- 20 No side shoots
- 21 First side shoot detectable
- 22 2 side shoots detectable
- 23 3 side shoots detectable
- 24 4 side shoots detectable
- 25 5 side shoots detectable
- 26 6 side shoots detectable
- 27 7 side shoots detectable
- 28 8 side shoots detectable
- 29 9 or more side shoots detectable

3 Principal growth stage 3: Stem elongation

- 30 no internodes("rosette")
- 31 1 visibly extended internode
- 32 2 visibly extended internodes
- 33 3 visibly extended internodes
- 34 4 visibly extended internodes
- 35 5 visibly extended internodes
- 36 6 visibly extended internodes
- 37 7 visibly extended internodes
- 38 8 visibly extended internodes
- 39 9 or more visibly extended internodes

4 Principal growth stage 4: Inflorescence emergence

- 40 Flower buds present, still enclosed by leaves
- 41 Flower buds visible from above ("green bud")
- 42 Flower buds free, level with the youngest leaves
- 43 Flower buds raised above the youngest leaves
- 45 Individual flower buds (main inflorescence) visible but still closed
- 47 Individual flower buds (secondary inflorescence) visible but still closed
- 49 First petals visible, flower buds still closed by ("yellow bud")

5 Principal growth stage 5: Opening of flowers

- 50 First flowers open
- 51 10% of flowers on main raceme open, main raceme elongating
- 52 20% of flowers on main raceme open
- 53 30% of flowers on main raceme open
- 54 40% of flowers on main raceme open
- 55 Full flowering: 50% flowers on main raceme open, older petals falling
- 57 Flowering declining: majority of petals fallen
- 59 End of flowering

- 6 Principal growth stage 6: Development of siliques
61 10% of siliques have reached final size
62 20% of siliques have reached final size
63 30% of siliques have reached final size
64 40% of siliques have reached final size
65 50% of siliques have reached final size
66 60% of siliques have reached final size
67 70% of siliques have reached final size
68 80% of siliques have reached final size
69 Nearly all siliques have reached final size
7 Principal growth stage 7: Ripening
70 seed green, filling siliques cavity
71 10% of siliques ripe, seeds dark and hard
72 20% of siliques ripe, seeds dark and hard
73 30% of siliques ripe, seeds dark and hard
74 40% of siliques ripe, seeds dark and hard
75 50% of siliques ripe, seeds dark and hard
76 60% of siliques ripe, seeds dark and hard
77 70% of siliques ripe, seeds dark and hard
78 80% of siliques ripe, seeds dark and hard
79 Fully ripe: nearly all siliques ripe, seeds dark and hard
8 Principal growth stage 8: Senescence
87 Plant dead and dry
89 Harvested product

Other Names of the Example Varieties

TTK456 ⁽¹⁾	Chaplin ⁽²⁾
Akaoba Takana ⁽³⁾	Red Giant ⁽⁴⁾

(1) official denomination registered under the law in Japan in 2011.

(2) official denomination of TTK456 registered under the law in European Union in 2014.

(3) commercial name in Japan.

(4) commercial name of Akaoba Takana in European Union.

9. Literature

Fujishiro, T., 1996: Breeding processes and characteristics of a newly bred leaf mustard (*Brassica Juncea* Coss.). Kanagawa, JP

Joy Larkcom, 1991: Oriental Vegetables (The Complete guide for Garden and Kitchen). London, GB, pp. 39 to pp. 45

Ministry of Agriculture, Forestry & Fisheries of Japan., 1994: National Test Guideline for Karashina

Phillips, R., Rix, M., 1993: Vegetables (The Pan Garden Plants Series). pp. 44

Tsukamoto, Y., 1994: The Grand Dictionary of Horticulture Volume 1. The Shogakukan Ltd., Tokyo, JP, pp. 520 to pp. 522

Takasi A., 2004: Yasai-engei-daihyakka 17. Shadanhojin Nousan-gyoson-bunkakyokai. Tokyo, JP. pp. 169 to pp. 233

Uwe Meier. Federal Biological Research Centre for Agriculture and Forestry, 2001: Growth stages of mono- and dicotyledonous plants, BBCH Monograph

10. Technical Questionnaire

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
		Application date: (not to be filled in by the applicant)
TECHNICAL QUESTIONNAIRE to be completed in connection with an application for plant breeders' rights		
1. Subject of the Technical Questionnaire		
1.1 Botanical name	<i>Brassica juncea</i> (L.) Czern.	
1.2 Common name	Brown mustard, Indian mustard, Oriental mustard	
2. Applicant		
Name	<input type="text"/>	
Address	<input type="text"/>	
Telephone No.	<input type="text"/>	
Fax No.	<input type="text"/>	
E-mail address	<input type="text"/>	
Breeder (if different from applicant)	<input type="text"/>	
3. Proposed denomination and breeder's reference		
Proposed denomination (if available)	<input type="text"/>	
Breeder's reference	<input type="text"/>	

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
#4. Information on the breeding scheme and propagation of the variety		
4.1 Breeding scheme		
Variety resulting from:		
4.1.1 Crossing		
(a) controlled cross (please state parent varieties)	[]	
(b) partially known cross (please state known parent variety(ies))	[]	
(c) unknown cross	[]	
4.1.2 Discovery and development (please state where and when discovered and how developed)		
<div style="border: 1px solid black; height: 100px;"></div>		
4.1.3 Mutation (please state parent variety)		
<div style="border: 1px solid black; height: 100px;"></div>		
4.1.4 Other (Please provide details)		
<div style="border: 1px solid black; height: 100px;"></div>		

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

4.2 Method of propagating the variety

4.2.1 Seed-propagated varieties

(a) Cross-pollination

[]

(b) Other (please provide details)

[]

4.2.2 Other

(Please provide details)

[]

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
5. Characteristics of the variety to be indicated (the number in brackets refers to the corresponding characteristic in Test Guidelines; please mark the note which best corresponds).		
Characteristics	Example Varieties	Note
5.1 Seed: color (1)		
yellow	Kigarashina	1 []
blackish brown	Akaoba Takana, Esperance, Miike Takana, Terrafit, Terraplus	2 []
5.2 Leaf: shape (5)		
ovate	Serihon	1 []
circular	Kekkyu Takana	2 []
elliptic	Akariasu	3 []
oblong	Etamine, Zasai FM-58	4 []
obovate	Esperance, Katsuona	5 []
spatulate	Kigarashina	6 []
5.3 Leaf: attitude (6)		
erect	Energy, Vittasso, Wasabina	1 []
erect to semi-erect		2 []
semi-erect	Esperance, Shinkoku Seisai	3 []
semi-erect to horizontal		4 []
horizontal	Etamine, Miike Takana	5 []
5.4 Leaf blade: undulation of margin (16)		
absent or very weak		1 []
weak	Akaoba Takana	2 []
medium	Katsuona	3 []
strong	Chirimen Hakarashina	4 []
very strong		5 []

Characteristics	Example Varieties	Note
5.5 (17) Leaf blade: density of incisions of margin		
absent or very sparse		1 []
very sparse to sparse		2 []
sparse	Etamine, Katsuona	3 []
sparse to medium		4 []
medium	Opaleska	5 []
medium to dense		6 []
dense	Oportuna	7 []
dense to very dense		8 []
very dense		9 []
5.6 (18) Leaf blade: blistering		
absent or weak	Etamine, Kigarashina	1 []
medium	Akaoba Takana	2 []
strong	Katsuona	3 []
5.7 (20) Plant: head formation		
absent	Kigarashina	1 []
present	Kekkyu Takana	9 []

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

6. Similar varieties and differences from these varieties

Please use the following table and box for comments to provide information on how your candidate variety differs from the variety (or varieties) which, to the best of your knowledge, is (or are) most similar. This information may help the examination authority to conduct its examination of distinctness in a more efficient way.

Denomination(s) of variety(ies) similar to your candidate variety	Characteristic(s) in which your candidate variety differs from the similar variety(ies)	Describe the expression of the characteristic(s) for the similar variety(ies)	Describe the expression of the characteristic(s) for your candidate variety
<i>Example</i>	<i>Leaf: shape</i>	<i>ovate</i>	<i>oblong</i>
Comments:			

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:																		
<p>#7. Additional information which may help in the examination of the variety</p> <p>7.1 In addition to the information provided in sections 5 and 6, are there any additional characteristics which may help to distinguish the variety?</p> <p>Yes <input type="checkbox"/> No <input type="checkbox"/></p> <p>(If yes, please provide details)</p> <p>7.2 Are there any special conditions for growing the variety or conducting the examination?</p> <p>Yes <input type="checkbox"/> No <input type="checkbox"/></p> <p>(If yes, please provide details)</p> <p>7.3 Other information</p> <p>(a) Main use</p> <table><tr><td>Vegetable</td><td><input type="checkbox"/></td></tr><tr><td>Oilseed</td><td><input type="checkbox"/></td></tr><tr><td>Condiment</td><td><input type="checkbox"/></td></tr><tr><td>Green manure</td><td><input type="checkbox"/></td></tr><tr><td>other</td><td><input type="checkbox"/></td></tr></table> <p>(b) Leaf type (according to Section 5.3 of the Test Guidelines)</p> <table><tr><td>type 1</td><td><input type="checkbox"/></td></tr><tr><td>type 2</td><td><input type="checkbox"/></td></tr><tr><td>type 3</td><td><input type="checkbox"/></td></tr><tr><td>type 4</td><td><input type="checkbox"/></td></tr></table>			Vegetable	<input type="checkbox"/>	Oilseed	<input type="checkbox"/>	Condiment	<input type="checkbox"/>	Green manure	<input type="checkbox"/>	other	<input type="checkbox"/>	type 1	<input type="checkbox"/>	type 2	<input type="checkbox"/>	type 3	<input type="checkbox"/>	type 4	<input type="checkbox"/>
Vegetable	<input type="checkbox"/>																			
Oilseed	<input type="checkbox"/>																			
Condiment	<input type="checkbox"/>																			
Green manure	<input type="checkbox"/>																			
other	<input type="checkbox"/>																			
type 1	<input type="checkbox"/>																			
type 2	<input type="checkbox"/>																			
type 3	<input type="checkbox"/>																			
type 4	<input type="checkbox"/>																			

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

8. Authorization for release

- (a) Does the variety require prior authorization for release under legislation concerning the protection of the environment, human and animal health?

Yes [] No []

- (b) Has such authorization been obtained?

Yes [] No []

If the answer to (b) is yes, please attach a copy of the authorization.

9. Information on plant material to be examined or submitted for examination

9.1 The expression of a characteristic or several characteristics of a variety may be affected by factors, such as pests and disease, chemical treatment (e.g. growth retardants or pesticides), effects of tissue culture, different rootstocks, scions taken from different growth phases of a tree, etc.

9.2 The plant material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If the plant material has undergone such treatment, full details of the treatment must be given. In this respect, please indicate below, to the best of your knowledge, if the plant material to be examined has been subjected to:

- (a) Microorganisms (e.g. virus, bacteria, phytoplasma) Yes [] No []
(b) Chemical treatment (e.g. growth retardant, pesticide) Yes [] No []
(c) Tissue culture Yes [] No []
(d) Other factors Yes [] No []

Please provide details for where you have indicated "yes".

.....

10. I hereby declare that, to the best of my knowledge, the information provided in this form is correct:

Applicant's name

Signature

 Date

[End of document]