



TG/16/9(proj.2)

ORIGINAL: English

DATE: 2018-04-06

INTERNATIONAL UNION FOR THE PROTECTION OF NEW VARIETIES OF PLANTS

Geneva

DRAFT

RICE

UPOV Code(s):

ORYZA_SAT

Oryza sativa L.

GUIDELINES

FOR THE CONDUCT OF TESTS

FOR DISTINCTNESS, UNIFORMITY AND STABILITY

*prepared by experts from Japan
to be considered by the
Technical Working Party for Agricultural Crops
at its forty-seventh session, to be held in Naivasha, Kenya,
from 2018-05-21 to 2018-05-25*

Disclaimer: this document does not represent UPOV policies or guidance

Alternative names:*

<i>Botanical name</i>	<i>English</i>	<i>French</i>	<i>German</i>	<i>Spanish</i>
<i>Oryza sativa</i> L.	Rice	Riz	Reis	Arroz

The purpose of these guidelines ("Test Guidelines") is to elaborate the principles contained in the General Introduction (document TG/1/3), and its associated TGP documents, into detailed practical guidance for the harmonized examination of distinctness, uniformity and stability (DUS) and, in particular, to identify appropriate characteristics for the examination of DUS and production of harmonized variety descriptions.

ASSOCIATED DOCUMENTS

These Test Guidelines should be read in conjunction with the General Introduction and its associated TGP documents.

* These names were correct at the time of the introduction of these Test Guidelines but may be revised or updated. [Readers are advised to consult the UPOV Code, which can be found on the UPOV Website (www.upov.int), for the latest information.]

TABLE OF CONTENTS	PAGE
1. SUBJECT OF THESE TEST GUIDELINES.....	4
2. MATERIAL REQUIRED.....	4
3. METHOD OF EXAMINATION.....	5
3.1 Number of Growing Cycles.....	5
3.2 Testing Place.....	5
3.3 Conditions for Conducting the Examination.....	5
3.4 Test Design.....	5
3.5 Additional Tests.....	5
4. ASSESSMENT OF DISTINCTNESS, UNIFORMITY AND STABILITY.....	6
4.1 Distinctness.....	6
4.2 Uniformity.....	7
4.3 Stability.....	8
5. GROUPING OF VARIETIES AND ORGANIZATION OF THE GROWING TRIAL.....	9
6. INTRODUCTION TO THE TABLE OF CHARACTERISTICS.....	10
6.1 Categories of Characteristics.....	10
6.2 States of Expression and Corresponding Notes.....	10
6.3 Types of Expression.....	10
6.4 Example Varieties.....	10
6.5 Legend.....	11
7. TABLE OF CHARACTERISTICS/TABLEAU DES CARACTÈRES/MERKMALSTABELLE/TABLA DE CARACTERES.....	12
8. EXPLANATIONS ON THE TABLE OF CHARACTERISTICS.....	25
8.1 Explanations covering several characteristics.....	25
8.2 Explanations for individual characteristics.....	26
9. LITERATURE.....	33
10. TECHNICAL QUESTIONNAIRE.....	34

1. Subject of these Test Guidelines

These Test Guidelines apply to all varieties of *Oryza sativa* L..

2. Material Required

2.1 The competent authorities decide on the quantity and quality of the plant material required for testing the variety and when and where it is to be delivered. Applicants submitting material from a State other than that in which the testing takes place must ensure that all customs formalities and phytosanitary requirements are complied with.

2.2 The material is to be supplied in the form of seed and panicles (if requested).

2.3 The minimum quantity of plant material, to be supplied by the applicant, should be:

Seed: 2 kg
Panicles (if requested): 120

The seed should meet the minimum requirements for germination, species and analytical purity, health and moisture content, specified by the competent authority. In cases where the seed is to be stored, the germination capacity should be as high as possible and should, be stated by the applicant.

The panicles should be well developed and should contain a sufficient number of viable seeds to establish a satisfactory row of plants for observation.

2.4 The plant material supplied should be visibly healthy, not lacking in vigor, nor affected by any important pest or disease.

2.5 The plant material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If it has been treated, full details of the treatment must be given.

3. Method of Examination

3.1 *Number of Growing Cycles*

3.1.1 The minimum duration of tests should normally be two independent growing cycles.

3.1.2 The two independent growing cycles should be in the form of two separate plantings.

3.2 *Testing Place*

Tests are normally conducted at one place. In the case of tests conducted at more than one place, guidance is provided in TGP/9 "Examining Distinctness".

3.3 *Conditions for Conducting the Examination*

3.3.1 The tests should be carried out under conditions ensuring satisfactory growth for the expression of the relevant characteristics of the variety and for the conduct of the examination.

3.3.2 The optimum stage of development for the assessment of each characteristic is indicated by a number in the Table of Characteristics. The stages of development denoted by each number are described in Chapter 8.

3.4 *Test Design*

3.4.1 Each test should be designed to result in a total of at least 1500 plants, which should be divided between at least 2 replicates.

3.4.2 The design of the tests should be such that plants or parts of plants may be removed for measurement or counting without prejudice to the observations which must be made up to the end of the growing cycle.

3.4.3 If tests on panicle rows are conducted, at least 100 panicle rows should be observed.

3.5 *Additional Tests*

Additional tests, for examining relevant characteristics, may be established.

4. Assessment of Distinctness, Uniformity and Stability

4.1 *Distinctness*

4.1.1 General Recommendations

It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding distinctness. However, the following points are provided for elaboration or emphasis in these Test Guidelines.

To assess distinctness of hybrids, the parent lines and the formula may be used according to the following recommendations:

- (i) description of parent lines according to the Test Guidelines;
- (ii) check of the originality of the parent lines in comparison with the variety collection, based on the characteristics in Chapter 7, in order to identify similar parent lines;
- (iii) check of the originality of the hybrid formula in relation to the hybrids in the variety collection, taking into account the most similar lines; and
- (iv) assessment of the distinctness at the hybrid level for varieties with a similar formula.

Further guidance is provided in documents TGP/9 "Examining Distinctness" and TGP/8 "Trial Design and Techniques Used in the Examination of Distinctness, Uniformity and Stability".

4.1.2 Consistent Differences

The differences observed between varieties may be so clear that more than one growing cycle is not necessary. In addition, in some circumstances, the influence of the environment is not such that more than a single growing cycle is required to provide assurance that the differences observed between varieties are sufficiently consistent. One means of ensuring that a difference in a characteristic, observed in a growing trial, is sufficiently consistent is to examine the characteristic in at least two independent growing cycles.

4.1.3 Clear Differences

Determining whether a difference between two varieties is clear depends on many factors, and should consider, in particular, the type of expression of the characteristic being examined, i.e. whether it is expressed in a qualitative, quantitative, or pseudo-qualitative manner. Therefore, it is important that users of these Test Guidelines are familiar with the recommendations contained in the General Introduction prior to making decisions regarding distinctness.

4.1.4 Number of Plants or Parts of Plants to be Examined

Unless otherwise indicated, for the purposes of distinctness, all observations on single plants should be made on 10 plants or parts of plants taken from each of 10 plants and any other observations made on all plants in the test, disregarding any off-type plants.

In the case of observations of parts taken from single plants, the number of parts to be taken from each of the plants should be 1.

4.1.5 Method of Observation

The recommended method of observing the characteristic for the purposes of distinctness is indicated by the following key in the Table of Characteristics (see document TGP/9 "Examining Distinctness", Section 4 "Observation of characteristics"):

MG: single measurement of a group of plants or parts of plants

MS: measurement of a number of individual plants or parts of plants

VG: visual assessment by a single observation of a group of plants or parts of plants

VS: visual assessment by observation of individual plants or parts of plants

Type of observation: visual (V) or measurement (M)

"Visual" observation (V) is an observation made on the basis of the expert's judgment. For the purposes of this document, "visual" observation refers to the sensory observations of the experts and, therefore, also includes smell, taste and touch. Visual observation includes observations where the expert uses reference points (e.g. diagrams, example varieties, side-by-side comparison) or non-linear charts (e.g. color charts). Measurement (M) is an objective observation against a calibrated, linear scale e.g. using a ruler, weighing scales, colorimeter, dates, counts, etc.

Type of record: for a group of plants (G) or for single, individual plants (S)

For the purposes of distinctness, observations may be recorded as a single record for a group of plants or parts of plants (G), or may be recorded as records for a number of single, individual plants or parts of plants (S). In most cases, "G" provides a single record per variety and it is not possible or necessary to apply statistical methods in a plant-by-plant analysis for the assessment of distinctness.

In cases where more than one method of observing the characteristic is indicated in the Table of Characteristics (e.g. VG/MG), guidance on selecting an appropriate method is provided in document TGP/9, Section 4.2.

- 4.1.6 A clear difference in a characteristic that is NOT in the Table of Characteristics, can be sufficient to establish Distinctness. A typical example would be a variety that is similar to an existing variety but the candidate variety has an additional resistance to biotic (insect, disease) or abiotic (salinity, submergence) stress or a herbicide tolerance, which is not or not yet in the Table of Characteristics.

4.2 Uniformity

- 4.2.1 It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding uniformity. However, the following points are provided for elaboration or emphasis in these Test Guidelines:
- 4.2.2 These Test Guidelines have been developed for the examination of [to be competed] varieties. For varieties with other types of propagation, the recommendations in the General Introduction and document TGP/13 "Guidance for new types and species" Section 4.5 "Testing Uniformity" should be followed.
- 4.2.3 The assessment of uniformity for hybrid varieties depends on the type of hybrid and should be according to the recommendations for hybrid varieties in the General Introduction.
- 4.2.4 Where the assessment of a hybrid variety involves the parent lines, the uniformity of the hybrid variety should, in addition to an examination of the hybrid variety itself, also be assessed by examination of the uniformity of its parent lines.
- 4.2.5 The recommended sample size for the assessment of uniformity is indicated by the following key in the table of characteristics:

A: sample size of 100 plants/parts of plants/panicle rows

B: sample size of 1500 plants

- 4.2.6 For the assessment of uniformity in a sample of 1500 plants, a population standard of 01% and an acceptance probability of at least 95% should be applied. In the case of a sample size of 1500 plants, 4 off-types are allowed.
- 4.2.7 For “A” characteristics, the assessment of uniformity can be done in 2 steps. In a first step, 20 plants or parts of plants are observed. If no off-types are observed, the variety is considered to be uniform. If more than 3 off-types are observed, the variety is considered not to be uniform. If 1 to 3 off-types are observed, an additional sample of 80 plants or parts of plants must be observed.
- 4.2.8 For the assessment of uniformity of hybrid varieties, a population standard of 10% and an acceptance probability of at least 95% should be applied. In case of characteristics indicated by B, the sample size for the assessment of uniformity may be reduced to 200 plants. In case of a sample size of 200 plants, 27 off-types are allowed. In case of a sample size of 100 ear-rows, plants or parts of plants, 15 off-types are allowed.

4.3 *Stability*

- 4.3.1 In practice, it is not usual to perform tests of stability that produce results as certain as those of the testing of distinctness and uniformity. However, experience has demonstrated that, for many types of variety, when a variety has been shown to be uniform, it can also be considered to be stable.
- 4.3.2 Where appropriate, or in cases of doubt, stability may be further examined by testing a new seed stock to ensure that it exhibits the same characteristics as those shown by the initial material supplied.
- 4.3.3 Where appropriate, or in cases of doubt, the stability of a hybrid variety may, in addition to an examination of the hybrid variety itself, also be assessed by examination of the uniformity and stability of its parent lines.

5. Grouping of Varieties and Organization of the Growing Trial

- 5.1 The selection of varieties of common knowledge to be grown in the trial with the candidate varieties and the way in which these varieties are divided into groups to facilitate the assessment of distinctness are aided by the use of grouping characteristics.
- 5.2 Grouping characteristics are those in which the documented states of expression, even where produced at different locations, can be used, either individually or in combination with other such characteristics: (a) to select varieties of common knowledge that can be excluded from the growing trial used for examination of distinctness; and (b) to organize the growing trial so that similar varieties are grouped together.
- 5.3 The following have been agreed as useful grouping characteristics:
- (a) Leaf blade: anthocyanin coloration (characteristic 6)
 - (b) Time of panicle emergence (characteristic 10)
 - (c) Stem: length (characteristic 15)
 - (d) Lemma: color of tip (characteristic 23)
 - (e) Decorticated grain: weight of 1000 seeds (characteristic 36)
 - (f) Decorticated grain: ratio length/width (characteristic 39)
 - (g) Decorticated grain: color (characteristic 40)
 - (h) Endosperm: type (characteristic 43)
- 5.4 Guidance for the use of grouping characteristics, in the process of examining distinctness, is provided through the General Introduction and document TGP/9 “Examining Distinctness”.

6. Introduction to the Table of Characteristics

6.1 *Categories of Characteristics*

6.1.1 Standard Test Guidelines Characteristics

Standard Test Guidelines characteristics are those which are approved by UPOV for examination of DUS and from which members of the Union can select those suitable for their particular circumstances.

6.1.2 Asterisked Characteristics

Asterisked characteristics (denoted by *) are those included in the Test Guidelines which are important for the international harmonization of variety descriptions and should always be examined for DUS and included in the variety description by all members of the Union, except when the state of expression of a preceding characteristic or regional environmental conditions render this inappropriate.

6.2 *States of Expression and Corresponding Notes*

6.2.1 States of expression are given for each characteristic to define the characteristic and to harmonize descriptions. Each state of expression is allocated a corresponding numerical note for ease of recording of data and for the production and exchange of the description.

6.2.2 In the case of qualitative and pseudo-qualitative characteristics (see Chapter 6.3), all relevant states of expression are presented in the characteristic. However, in the case of quantitative characteristics with 5 or more states, an abbreviated scale may be used to minimize the size of the Table of Characteristics. For example, in the case of a quantitative characteristic with 9 states, the presentation of states of expression in the Test Guidelines may be abbreviated as follows:

<i>State</i>	<i>Note</i>
small	3
medium	5
large	7

However, it should be noted that all of the following 9 states of expression exist to describe varieties and should be used as appropriate:

<i>State</i>	<i>Note</i>
very small	1
very small to small	2
small	3
small to medium	4
medium	5
medium to large	6
large	7
large to very large	8
very large	9

6.2.3 Further explanation of the presentation of states of expression and notes is provided in document TGP/7 "Development of Test Guidelines".

6.3 *Types of Expression*

An explanation of the types of expression of characteristics (qualitative, quantitative and pseudo-qualitative) is provided in the General Introduction.

6.4 *Example Varieties*

Where appropriate, example varieties are provided to clarify the states of expression of each characteristic.

6.5 Legend

		English	français	deutsch	español	Example Varieties Exemples Beispielsorten Variedades ejemplo	Note/ Nota
1	2	3	4	5	6	7	
		Name of characteristics in English	Nom du caractère en français	Name des Merkmals auf Deutsch	Nombre del carácter en español		
		states of expression	types d'expression	Ausprägungsstufen	tipos de expresión		

- 1 Characteristic number
- 2 (*) Asterisked characteristic – see Chapter 6.1.2
- 3 Type of expression
 QL Qualitative characteristic – see Chapter 6.3
 QN Quantitative characteristic – see Chapter 6.3
 PQ Pseudo-qualitative characteristic – see Chapter 6.3
- 4 Method of observation (and type of plot, if applicable)
 MG, MS, VG, VS – see Chapter 4.1.5
- 5 (+) See Explanations on the Table of Characteristics in Chapter 8.2
- 6 (a)
- 7 Growth stage key See Explanations on the Table of Characteristics in Chapter 8

A: sample size of 100 plants/parts of plants
 B: sample size of 1500 plants/parts of plants

7. Table of Characteristics/Tableau des caractères/Merkmalstabelle/Tabla de caracteres

	English		français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
1.	QN	VG A	(+)				10
	Coleoptile: anthocyanin coloration						
	absent or weak					koshihikari	1
	medium					murasakikoboshi	3
	strong					akaneasobi, satsumakuromochi	5
2. (*)	QN	VG B	(+)				40
	Plant: growth habit						
	erect					leafstar	1
	semi-erect					koshihikari, momiroman	3
	open					onari	5
	spreading						7
	prostrate						9
3.	QN	VG B	(a)				40
	Leaf sheath: anthocyanin coloration						
	absent or very weak					koshihikari	1
	weak					murasakikoboshi, sayomurasaki	3
	medium					minamiyutaka	5
	strong					beniasobi, shikibumochi	7
	very strong						9
4.	QN	VG B					40
	Basal leaf sheath: anthocyanin coloration						
	absent or very weak						1
	weak						3
	medium						5
	strong						7
	very strong						9

	English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
5.	QN	VG B	(a)	40		
	Leaf blade: intensity of green color					
	light				koihonoka	3
	medium				hinohikari, koshihikari	5
	dark				hoshiyutaka, takanari	7
6. (*)	QL	VG B	(a)	40		
	Leaf blade: anthocyanin coloration					
	absent				koshihikari	1
	present				murakikoboshi, sayomurasaki	9
7.	QN	VG B	(+)	(a)	40	
	Leaf blade: pubescence					
	absent or very weak				leafstar	1
	weak					2
	medium				koshihikari	3
	strong					4
	very strong					5
8.	PQ	VG B	(+)	(a)	40	
	Ligule: shape					
	truncate					1
	acute				murakikoboshi	2
	cleft				onari, salt star	3
9.	PQ	VG B	(a)	40		
	Ligule: color					
	white				koshihikari	1
	green					2
	purple				beniasobi, sayomurasaki	3

	English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
10. (*)	QN MG B	(+)	55			
	Time of panicle emergence					
	very early					1
	early				koshihikari	3
	medium				momiroman	5
	late				leafstar	7
	very late					9
11.	QN MS B/VG B	(+)	60-70			
	Flag leaf blade: length					
	short				ouukan 383	3
	medium				hinohikari	5
	long				tachiaoba	7
12.	QN MS B/VG B	(+)	60-70			
	Flag leaf blade: width					
	narrow				ouukan 383	1
	medium				hinohikari	3
	broad				tachiaoba	5
13.	QN VG B		60-80			
	Lemma: pubescence					
	absent or very weak				leafstar	1
	weak				murakikoboshi	2
	medium				koshihikari	3
	strong					4
	very strong					5
14. (*)	PQ VG B		65			
	Stigma: color					
	white				koshihikari	1
	light green					2
	yellow					3
	purple				ouukan 383, sayomurasaki	4
	black				murakikoboshi, shikibumochi	5

	English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
15. (*)	QN	MG B/MS B	(+)	70		
	Stem: length					
	very short					1
	short				takanari	3
	medium				hinohikari	5
	long				koshihikari	7
	very long				minamiyutaka	9
16.	QN	VG B	(+)	70		
	Stem: thickness					
	thin				murakikoboshi	3
	medium				hinohikari, koshihikari	5
	thick				hoshiyutaka, momiroman	7
17.	QN	VG B		70		
	Node: anthocyanin coloration					
	absent or weak				koshihikari	1
	medium				sayomurasaki	3
	strong				murakikoboshi	5
18.	QN	VG B		70		
	Internode: anthocyanin coloration					
	absent or weak				koshihikari	1
	medium					3
	strong				shikibumochi	5
19. (*)	QN	MS B		70		
	Panicle: number					
	few				momiroman, takanari	3
	medium				koshihikari	5
	many				ouukan 383	7

	English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
20.	(*) QN VG B					70
	Panicle: distribution of awns					
	absent				momiroman, onari	1
	1/4 upper				sari queen	2
	upper half					3
	3/4 of the total length				beniroman	4
	whole length				saikaikan 246	5
21.	QN VG B	(+)				70-80
	Awns: length					
	very short				hinohikari	1
	short				koshihikari	2
	medium				benizomemochi, leafstar	3
	long				saikaikan 246	4
	very long					5
22.	(*) QN MS B	(+)				72-90
	Panicle: length					
	short				shikibumochi	3
	medium				koshihikari, leafstar	5
	long				momiroman	7
23.	(*) PQ VG B					80-90
	Lemma: color of tip					
	white				koshihikari	1
	yellowish					2
	brown				koshinokaori, leafstar	3
	red				minamiyutaka	4
	purple				murakikoboshi, sayomurasaki	5
	black					6

	English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
24. (*)	QN	VG B	(+)	80-90		
	Flag leaf: attitude of blade					
	erect				leafstar, minamiyutaka	1
	semi-erect				momiroman, onari	3
	horizontal				murasakikoboshi, ouukan 383	5
	semi-drooping					7
	drooping					9
25.	QN	VG B		90		
	Panicle: density					
	lax					3
	medium				koshihikari	5
	dense				hoshiyutaka, takanari	7
26.	QN	VG B	(+)	90		
	Panicle: attitude					
	erect				akaneasobi	1
	semi-erect				ouukan 383	2
	semi-drooping				koshihikari	3
	drooping					4
27. (*)	QN	VG B	(+)	90		
	Panicle: attitude of branches					
	erect				habataki	1
	semi-erect				murasakikoboshi	3
	spreading					5
28.	QN	VG B		90		
	Panicle: number of secondary branches					
	absent or few					1
	medium				koshihikari	2
	many				takanari	3

	English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
29.	QN	VG B	(+)	90		
	Panicle: exertion					
	enclosed					1
	partly exerted				tachisuzuka	2
	just exerted				minamiyutaka	3
	well exerted				koshihikari	4
30. (*)	QN	MG B		90		
	Time of maturity					
	very early					1
	early				koshihikari	3
	medium				asahinoyume	5
	late				leafstar	7
	very late					9
31.	QN	VG B	(+)	92		
	Time of senescence					
	early				onari	3
	medium				salt star	5
	late				koshihikari	7
32.	PQ	VG B		92		
	Lemma: color					
	white				koshihikari	1
	yellowish				leafstar	2
	brown				beniasobi	3
	red					4
	purple				ouukan 383, satsumakuromochi	5
	black					6
33.	QN	VG B	(+)	92		
	Lemma: coloration with phenol					
	absent or very light				koshihikari, momiroman	1
	light					3
	medium				onari, salt star	5
	dark				ruriaoba	7
	very dark					9

	English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
34.	QN	VG B	(+)	92		
	Glume: length					
	short				ruriaoba	1
	medium				koshihikari	2
	long					3
35.	PQ	VG B		92		
	Glume: color					
	white				koshihikari	1
	yellowish					2
	brown					3
	red					4
	purple				beniasobi, ouukan 383	5
	black					6
36. (*)	QN	MG B	(+)	92		
	Decorticated grain: weight of 1000 seeds					
	low				beniasobi, sari queen	3
	medium				koshihikari, takanari	5
	high				momiroman	7
37. (*)	QN	MS B		92		
	Decorticated grain: length					
	short				murasakikoboshi	3
	medium				koshihikari	5
	long				hoshiyutaka, leafstar	7
38.	QN	MS B		92		
	Decorticated grain: width					
	narrow				hoshiyutaka, leafstar	1
	medium				koshihikari	3
	broad					5

	English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
39. (*)	QN MS B	(+)	92			
	Decorticated grain: ratio length/width					
	small				akaneasobi	1
	small to medium				koshihikari	2
	medium				hoshiyutaka, leafstar	3
	medium to large					4
	large					5
40. (*)	PQ VG B		92			
	Decorticated grain: color					
	white				ruriaoba	1
	light brown				koshihikari, takanari	2
	dark brown				leafstar	3
	red				benizomemochi	4
	purple					5
	black				murakikoboshi, sayomurasaki	6
41.	QN MG B	(+)	92			
	Decorticated grain: alkali digestion					
	absent or very weak				koshinokaori	1
	weak				murakikoboshi, oukan 383	3
	medium				salt star	5
	strong				koshihikari	7
	very strong					9
42. (*)	QN MG B	(+)	92			
	Decorticated grain: aroma					
	absent or weak				koshihikari	1
	medium				sari queen	2
	strong					3

	English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
43. (*)	PQ	VG B	(+)	92		
	Endosperm: type					
	glutinous				ruriaoba, sayomurasaki	1
	intermediate					2
	non-glutinous				koshihikari, takanari	3
44. (*)	QN	MG B	(+)	92		
	Endosperm: content of amylose					
	very low				ruriaoba, sayomurasaki	1
	low					3
	medium				koshihikari	5
	high				hoshiyutaka	7
	very high				koshinokaori	9

8. Explanations on the Table of Characteristics

8.1 *Explanations covering several characteristics*

Characteristics containing the following key in the Table of Characteristics should be examined as indicated below:


- (a) Unless otherwise indicated, all observations on the leaf should be made on the penultimate leaf.

8.2 *Explanations for individual characteristics*

Ad. 1: Coleoptile: anthocyanin coloration

Non-dormant grains are placed on moistened filter paper and covered with a petri-dish lid during germination. After the coleoptiles have reached a length of about 5mm in darkness they are placed in artificial light (daylight equivalent) at 750-1250 lux continuously for 3 to 4 days, at a temperature of 25 to 30 degree. The color of the coleoptiles is observed when they are fully developed at stage 09-11 (about 6 to 7 days).


Ad. 2: Plant: growth habit



Ad. 7: Leaf blade: pubescence

Observations should be made on the upper side of the blade.

Ad. 8: Ligule: shape



Ad. 10: Time of panicle emergence

Time of panicle emergence is reached when the first spikelet is visible on 50% of panicles.

Ad. 11: Flag leaf blade: length

Measurements of flag leaf blade should be made on the same leaf. Length should be measured from the tip to base of the leaf blade. Width should be measured at the widest part of the leaf blade.

Ad. 12: Flag leaf blade: width

See Ad. 11

Ad. 15: Stem: length

Measurements should be made from the base to the panicle base on the longest stem.


Ad. 16: Stem: thickness

Observations should be observed at basal internode of the longest stem.

Ad. 21: Awns: length


Measurements should be made on the longest awn in the panicle.

Ad. 22: Panicle: length




Length of panicle should be observed from panicle base to the top excluding awns.


Ad. 24: Flag leaf: attitude of blade




Ad. 26: Panicle: attitude



Ad. 27: Panicle: attitude of branches



Ad. 29: Panicle: exertion



Ad. 31: Time of senescence

The leaves below the flag leaf are observed at the time of harvest for their retention of greenness.

3 - early: Leaves are dead when the grains have become fully ripened.

5 - medium: There must be 1 leaf which retains its color.

7 - late: 2 or more leaves retain their color at maturity.

Ad. 33: Lemma: coloration with phenol

Method of testing: Place hulls from 10 grains into a petri dish of 5cm diameter, and add 5ml of 1.5% phenol solution. Cover the petri dish, and keep at room temperature (not very cold) for one day.

Ad. 34: Glume: length

Measurements should be made on longer glume.

Ad. 36: Decorticated grain: weight of 1000 seeds

Measurements should be calculated at 14% moisture.

Ad. 39: Decorticated grain: ratio length/width

1 - small: < 1.50

2 - small to medium: 1.50-1.99

3 - medium: 2.00-2.49

4 - medium to large: 2.50-2.99

5 - large: > 2.99

Ad. 41: Decorticated grain: alkali digestion

Put 10 milled complete (unbroken) rice grains in a petri dish with 1.5% solution of KOH, and keep still under room temperature of around 25 degree for about 24 hours.

1 - not digested: Rice grains are not affected.

3 - low digested: Only the margin of the grains are dissolved.

5 - intermediate: Shape of grains become unclear, but incompletely dissolved.

7 - completely digested: No margin is identified between the core part and the outer skirt.

Ad. 42: Decorticated grain: aroma

The main component of the aroma in rice is the 2-acetyl-1-pyrroline (AcPy). To vaporize this chemical, 10ml of a 1.7% solution of KOH should be added to 2g of decorticated grains. The aroma, which is similar to that in pop-corn, is released within 10 minutes. The level of expression is determined by reference to the example varieties.

Ad. 43: Endosperm: type

The three states of expression can be simply defined by reaction to KI-I solution which is prepared by mixing 0.1 % I₂ solution and 0.2 % KI solution.

- 1 - glutinous: endosperm is stained to reddish purple.
- 2 - intermediate: endosperm is stained to reddish blue purple.
- 3 - non-glutinous: endosperm is stained to dark blue purple.

Ad. 44: Endosperm: content of amylose

Method ISO 6647 should be used.

8.3 8.3 Decimal code for the growth stage codes of cereals

Germination

00	Dry seed
01	Start of imbibition
02	-
03	Imbibition complete
04	-
05	Radicle emerged from caryopsis
06	-
07	Coleoptile emerged from caryopsis
08	-
09	Leaf just at coleoptile tip

Seedling growth

10	First leaf through coleoptile
11	First leaf unfolded (1)
12	2 leaves unfolded
13	3 leaves unfolded
14	4 leaves unfolded
15	5 leaves unfolded
16	6 leaves unfolded
17	7 leaves unfolded
18	8 leaves unfolded
19	9 or more leaves unfolded

Germination

20	Main shoot only
21	Main shoot and 1 tiller
22	Main shoot and 2 tillers
23	Main shoot and 3 tillers
24	Main shoot and 4 tillers
25	Main shoot and 5 tillers
26	Main shoot and 6 tillers
27	Main shoot and 7 tillers
28	Main shoot and 8 tillers
29	Main shoot and 9 or more tillers

Stem elongation

30	Pseudo stem erection (2)
31	1st node detectable
32	2nd node detectable
33	3rd node detectable
34	4th node detectable
35	5th node detectable
36	6th node detectable
37	Flag leaf just visible
38	-
39	Flag leaf ligule/collar just visible

Booting

- 40 -
- 41 Flag leaf sheath extending
- 42 -
- 43 Boots just visibly swollen
- 44 -
- 45 Boots swollen
- 46 -
- 47 Flag leaf sheath opening
- 48 -
- 49 First awns visible

Inflorescence emergence

- 50 First spikelet of inflorescence just visible
- 51
- 52
- 53 1/4 of inflorescence emerged
- 54
- 54 1/2 of inflorescence emerged
- 55
- 56 3/4 of inflorescence emerged
- 57
- 58
- 58 Emergence of inflorescence completed
- 59

Anthesis

- 60
- 60 Beginning of anthesis
- 61
- 62 -
- 63 -
- 64
- 64 Anthesis half-way
- 65
- 66 -
- 67 -
- 68
- 68 Anthesis complete
- 69

Milk development

70	-
71	Caryopsis watery ripe
72	-
73	Early milk
74	-
75	Medium milk
76	-
77	Late milk
78	-
79	-

Dough development

80	-
81	-
82	-
83	Early dough
84	-
85	Soft dough
86	-
87	Hard dough
88	-
89	-

Ripening

90	-
91	Caryopsis hard (difficult to divide by thumbnail) (3)
92	Caryopsis hard (can no longer be dented by thumbnail) (4)
93	Caryopsis loosening in daytime
94	Over-ripe, straw dead and collapsing

Ripening (continued)

95	Seed dormant
96	Viable seed giving 50% germination
97	Seed not dormant
98	Secondary dormancy induced
99	Secondary dormancy lost

Notes on the table

- (1) Stage of seedling inoculation with rust in the greenhouse.
- (2) Only applicable to cereals with a prostrate or semi-prostrate early growth habit.
- (3) Ripeness for binder (ca. 16% water content). Chlorophyll of inflorescence largely lost.
- (4) Ripeness for combine harvester (< 16% water content).
- (5) Optimum harvest time.

9. Literature

Matsuo, T. (edit.), 1993-97: Science of the Rice Plant. Nosan Gyoson Bunka Kyokai. Tokyo, JP
Vol. 1 Morphology (1993)
Vol. 2 Physiology (1995)
Vol. 3 Genetics (1997)

Zadoks, J.C., Chang, T.T., Konzak, C.F., 1974: A Decimal code for the Growth Stages of Cereals. Weed Research. NL, 14: pp. 415 – 421.

10. Technical Questionnaire

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

	Application date: (not to be filled in by the applicant)
--	---

TECHNICAL QUESTIONNAIRE
to be completed in connection with an application for plant breeders' rights

1. Subject of the Technical Questionnaire

1.1 Botanical name

Oryza sativa L.

1.2 Common name

Rice

2. Applicant

Name

Address

Telephone No.

Fax No.

E-mail address

Breeder (if different from applicant)

3. Proposed denomination and breeder's reference

Proposed denomination
(if available)

Breeder's reference

#4. Information on the breeding scheme and propagation of the variety

4.1 Breeding scheme

Variety resulting from:

4.1.1 Crossing

(a) controlled cross
(please state parent varieties)

(.....) x (.....)

female parent

male parent

(b) partially known cross
(please state known parent variety(ies))

(.....) x (.....)

female parent

male parent

(c) unknown cross

4.1.2 Discovery and development
(please state where and when discovered and how developed)

4.1.3 Mutation
(please state parent variety)

4.1.4 Other
(Please provide details)

4.2	Method of propagating the variety	
4.2.1	Seed-propagated varieties	
(a)	Self-pollination	[]
(b)	Hybrid	[]
(c)	Other (please provide details)	[]
4.2.2	Other (Please provide details)	[]
<div style="border: 1px solid black; width: 400px; height: 20px; margin: 0 auto;"></div>		

In the case of hybrid varieties the production scheme for the hybrid should be provided on a separate sheet. This should provide details of all the parent lines required for propagating the hybrid e.g.

Single Hybrid

(.....)	x	(.....)
female parent		male parent

Three-Way Hybrid

(.....)	x	(.....)
female parent		male parent

(.....)	x	(.....)
single hybrid used as female parent		male parent

and should identify in particular:

- (a) any male sterile lines
- (b) maintenance system of male sterile lines.

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

5. Characteristics of the variety to be indicated (the number in brackets refers to the corresponding characteristic in Test Guidelines; please mark the note which best corresponds).

Characteristics	Example Varieties	Note
5.1 Leaf blade: anthocyanin coloration (6)		
absent	koshihikari	1 []
present	murakikoboshi, sayomurasaki	9 []
5.2 Time of panicle emergence (10)		
very early		1 []
early	koshihikari	3 []
medium	momiroman	5 []
late	leafstar	7 []
very late		9 []
5.3 Stem: length (15)		
very short		1 []
short	takanari	3 []
medium	hinohikari	5 []
long	koshihikari	7 []
very long	minamiyutaka	9 []
5.4 Panicle: length (22)		
short	shikibumochi	3 []
medium	koshihikari, leafstar	5 []
long	momiroman	7 []
5.5 Lemma: color of tip (23)		
white	koshihikari	1 []
yellowish		2 []
brown	koshinokaori, leafstar	3 []
red	minamiyutaka	4 []
purple	murakikoboshi, sayomurasaki	5 []
black		6 []

Characteristics	Example Varieties	Note
5.6 Decorticated grain: weight of 1000 seeds (36)		
low	beniasobi, sari queen	3 []
medium	koshihikari, takanari	5 []
high	momiroman	7 []
5.7 Decorticated grain: color (40)		
white	ruriaoba	1 []
light brown	koshihikari, takanari	2 []
dark brown	leafstar	3 []
red	benizomemochi	4 []
purple		5 []
black	murasakikoboshi, sayomurasaki	6 []
5.8 Endosperm: type (43)		
glutinous	ruriaoba, sayomurasaki	1 []
intermediate		2 []
non-glutinous	koshihikari, takanari	3 []

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

6. Similar varieties and differences from these varieties

Please use the following table and box for comments to provide information on how your candidate variety differs from the variety (or varieties) which, to the best of your knowledge, is (or are) most similar. This information may help the examination authority to conduct its examination of distinctness in a more efficient way.

Denomination(s) of variety(ies) similar to your candidate variety	Characteristic(s) in which your candidate variety differs from the similar variety(ies)	Describe the expression of the characteristic(s) for the similar variety(ies)	Describe the expression of the characteristic(s) for your candidate variety
<i>Example</i>	<i>Panicle: length</i>	<i>long</i>	<i>short to medium</i>
Comments:			

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

#7.	Additional information which may help in the examination of the variety		
7.1	In addition to the information provided in sections 5 and 6, are there any additional characteristics which may help to distinguish the variety?		
	Yes	[]	No []
	(If yes, please provide details)		
7.2	Are there any special conditions for growing the variety or conducting the examination?		
	Yes	[]	No []
	(If yes, please provide details)		
7.3	Other information		

8. Authorization for release

(a) Does the variety require prior authorization for release under legislation concerning the protection of the environment, human and animal health?

Yes [] No []

(b) Has such authorization been obtained?

Yes [] No []

If the answer to (b) is yes, please attach a copy of the authorization.

9. Information on plant material to be examined or submitted for examination

9.1 The expression of a characteristic or several characteristics of a variety may be affected by factors, such as pests and disease, chemical treatment (e.g. growth retardants or pesticides), effects of tissue culture, different rootstocks, scions taken from different growth phases of a tree, etc.

9.2 The plant material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If the plant material has undergone such treatment, full details of the treatment must be given. In this respect, please indicate below, to the best of your knowledge, if the plant material to be examined has been subjected to:

(a) Microorganisms (e.g. virus, bacteria, phytoplasma)	Yes []	No []
(b) Chemical treatment (e.g. growth retardant, pesticide)	Yes []	No []
(c) Tissue culture	Yes []	No []
(d) Other factors	Yes []	No []

Please provide details for where you have indicated "yes".

.....

10. I hereby declare that, to the best of my knowledge, the information provided in this form is correct:

Applicant's name

Signature Date

[End of document]