TECHNICAL WORKING PARTY FOR AGRICULTURAL CROPS

Forty-Eighth Session

PREPARATORY WORKSHOP

Naivasha, Kenya, May 22, 2018

UPQV

International Union for the Protection of New Varieties of Plants

PROGRAM

- 1. Introduction to UPOV and the role of UPOV Technical Working Parties (TWPs)
- 2. Overview of the General Introduction (document TG/1/3 and TGP documents)
 - Characteristics as the Basis for DUS Examination and Selection of Characteristics
 - Molecular techniques
- 3. Guidance on drafting Test Guidelines (document TGP/7)
 - a) Subject of the Test Guidelines, Material Required and Method of Examination;
 - b) Method of Observation (MS, MG, VS, VG);
 - c) Types of Expression (QL, PQ, QN), notes and distinctness;
 - d) Shape and Color Characteristics;
 - e) Example Varieties;
 - f) The process for developing UPOV Test Guidelines, including: TG Template; Additional Standard Wording; and Guidance Notes;
- 4. Agenda for the TWP Session

1. INTRODUCTION TO UPOV AND THE ROLE OF UPOV TECHNICAL WORKING PARTIES (TWPS)


UPOV: INDEPENDENT INTERGOVERNMENTAL ORGANIZATION


The International Convention for the Protection of New Varieties of Plants established in 1961


The International Union for the Protection of New Varieties of Plants


Union internationale pour la protection des obtentions végétales











Role of the BMT

The BMT is a group open to DUS experts, biochemical and molecular specialists and plant breeders, whose role is to:

- (i) Review general developments in biochemical and molecular techniques;
- (ii) Maintain an awareness of relevant applications of biochemical and molecular techniques in plant breeding;
- (iii) Consider the possible application of biochemical and molecular techniques in DUS testing and report its considerations to the TC;
- (iv) If appropriate, establish guidelines for biochemical and molecular methodologies and their harmonization [...];
- (v) Consider initiatives from TWPs, for the establishment of crop specific subgroups [...];
- (vi) Develop guidelines regarding the management and harmonization of databases of biochemical and molecular information, in conjunction with the TWC;
- (vii) Receive reports from Crop Subgroups and the BMT Review Group;
- (viii) Provide a forum for discussion on the use of biochemical and molecular techniques in the consideration of essential derivation and variety identification.

Questions

2. OVERVIEW OF THE GENERAL INTRODUCTION (document TG/1/3 and TGP documents)

- a) Characteristics as the Basis for DUS

 Examination
 - **b) Selection of Characteristics**

2. OVERVIEW OF THE GENERAL INTRODUCTION (document TG/1/3 and TGP documents)

- a) Characteristics as the Basis for DUS

 Examination
 - b) Selection of Characteristics

THE CONDITIONS FOR GRANTING A BREEDER'S RIGHT

Criteria to be satisfied

- NOVELTY
- **DISTINCTNESS**
- **U**NIFORMITY
- **S**TABILITY

"DUS"

THE CONDITIONS FOR GRANTING A BREEDER'S RIGHT

Other conditions

- VARIETY DENOMINATION
- FORMALITIES
- PAYMENT OF FEES

NO OTHER CONDITIONS!

Guidance for DUS Examination

facilitates:


BEST PRACTICE (based on experience)

- => good decisions
- => good definition of the object of protection
 (strong protection)
- => efficiency in method of examination (learn from the best)

HARMONIZATION

- => efficiency
 - mutual acceptance of DUS reports (minimize cost of examination for individual authorities)
 - mutual recognition of variety descriptions (all parties speak the same "language")
 - simple and cheap system for applicants (minimize cost for breeders)

UPOV provides guidance by: The "General Introduction" (TG/1/3) General technical principles Organization of DUS Testing Associated "TGP" Documents (e.g. statistical methods) = version 3



2. OVERVIEW OF THE GENERAL INTRODUCTION (document TG/1/3 and TGP documents)

- a) Characteristics as the Basis for DUS

 Examination
 - b) Selection of Characteristics

"CHARACTERISTICS"

- may have direct commercial relevance
 - Flower color (ornamental)
 - Fruit color
- but commercial relevance NOT required
 - Leaf shape

Selection of Characteristics

The basic requirements that a characteristic should fulfill before it is used for DUS testing or producing a variety description are that its expression (TG/1/3: Section 4.2.1):

- (a) **results from a given genotype** or combination of genotypes;
- (b) is sufficiently consistent and repeatable in a particular environment:
- (c) exhibits sufficient **variation between varieties** to be able to establish distinctness:
- (d) is capable of precise definition and recognition;
- (e) allows uniformity requirements to be fulfilled;
- (f) allows **stability requirements** to be fulfilled, meaning that it produces consistent and repeatable results after repeated propagation or, where appropriate, at the end of each cycle of propagation.

Selection of Characteristics

- Yield ???
- Straw strength ???


Etc.

Criteria	Fruit: color	Leaf: shape
(a) results from a given genotype or combination of genotypes	Yes	Yes
(b) sufficiently consistent and repeatable in a particular environment	Yes	Yes
(c) exhibits sufficient variation between varieties to be able to establish distinctness	Yes	Yes
(d) is capable of precise definition and recognition	Yes	Yes
(e) allows uniformity requirements to be fulfilled	Yes	Yes
(f) allows stability requirements to be fulfilled	Yes	Yes
Commercial value	Yes	No

Selection of Characte	31 13 tic	5	
Criteria	Fruit: color	Leaf: shape	Yie
(a) results from a given genotype or combination of genotypes	Yes	Yes	Ye
(b) sufficiently consistent and repeatable in a particular environment	Yes	Yes	(No
(c) exhibits sufficient variation between varieties to be able to establish distinctness	Yes	Yes	??'
(d) is capable of precise definition and recognition	Yes	Yes	(No
(e) allows uniformity requirements to be fulfilled	Yes	Yes	??′
(f) allows stability requirements to be fulfilled	Yes	Yes	??′
Commercial value	Yes	No	Yes
ACCEPTABILITY	Yes	Yes	No


Criteria	Disease Resistance
(a) results from a given genotype or combination of genotypes	*Knowledge of nature of genetic control of resistance is important
(b) sufficiently consistent and repeatable in a particular environment	*Standardize conditions (greenhouse / laboratory) & methodology *Standardize inoculum *Ring-test
(c) exhibits sufficient variation between varieties to be able to establish distinctness	*Susceptible / Resistant OR varying degrees resistance?
(d) is capable of precise definition and recognition	*Define and recognize races and strains
(e) allows uniformity requirements to be fulfilled	see above
(f) allows stability requirements to be fulfilled	see above
	Difficult and expensive


Questions



IMPORTANCE OF HARMONIZED APPROACH WITHIN UPOV

- ⇒ To facilitate cooperation in DUS testing e.g. purchase of DUS reports
- ⇒ To establish internationally recognized variety descriptions (effective protection)





STATUS OF UPOV DOCUMENTS CONCERNING MOLECULAR TECHNIQUES

Document reference	Title
UPOV/INF/17/1	Guidelines for DNA Profiling: Molecular Marker Selection and Database Construction ("BMT Guidelines") (2010)

Document reference	Title
TGP/15	Guidance on the Use of Biochemical and Molecular Markers in the Examination of Distinctness, Uniformity and Stability (DUS)
UPOV/INF/18/1	Possible Use of Molecular Markers in the Examination of Distinctness, Uniformity and Stability (2011)

UPOV/INF/17/1 (INFormation document)

"Guidelines for DNA Profiling: Molecular Marker Selection and Database Construction ("BMT Guidelines")"

The purpose of this document (BMT Guidelines) is to provide guidance for developing harmonized methodologies with the aim of generating high quality molecular data for a range of applications. The BMT Guidelines are also intended to address the construction of databases containing molecular profiles of plant varieties [...]

UPOV/INF/18/1 (INFormation document)

"Possible Use of Molecular Markers in the Examination of Distinctness, Uniformity and Stability"

The purpose of this document is to provide guidance on the possible use of biochemical and molecular markers in the examination of Distinctness, Uniformity and Stability (DUS). [...]


UPOV/INF/18 POSSIBLE APPLICATION MODELS

MODELS WITH A POSITIVE ASSESSMENT

- Characteristic-specific molecular markers
- Combining phenotypic and molecular distances in the management of variety collections
- Calibrated molecular distances in the management of variety collections

MODELS WITHOUT A POSITIVE ASSESSMENT

Use of molecular marker characteristics



TGP/15/1 (Technical Guidelines Protocol)

"Guidance on the Use of Biochemical and Molecular Markers in the Examination of Distinctness, Uniformity and Stability (DUS)"

The purpose of this document is to provide guidance on the use of biochemical and molecular markers in the examination of Distinctness, Uniformity and Stability (DUS) on the basis of the models in document UPOV/INF/18 that have received a positive assessment and for which accepted examples have been provided.

→ Adopted by the Council of UPOV in October, 2013.

Model 1: Characteristic-specific molecular markers


Example: gene specific marker for herbicide tolerance introduced by genetic modification

On the basis that:

[...]

- there is <u>verification of the reliability</u> of the link between the marker and the characteristic;
- <u>different markers for the same characteristic</u> are different methods for examining the same characteristic;


[...]





Combining phenotypic and molecular distances in the management of variety collections

"A combination of phenotypic differences and molecular distances can be used to identify within the variety collection, those varieties which need to be compared with candidate varieties in order to improve the selection of "Distinct plus" varieties, on the following basis:

- (a) there is <u>reliable information that the molecular distances are sufficiently related to phenotypic differences</u>, such that
- (b) the method selects varieties in the variety collection which are similar to the candidate varieties; and
- (c) the method does not create an increased risk of not selecting a variety in the variety collection which needs to be compared to the candidate varieties in the field.







OVERVIEW

Molecular techniques

- UPOV guidance on molecular techniques
- Coordination with other international organizations
- Possible future developments
- Summary (FAQs)

43







OECD/UPOV/ISTA JOINT WORKSHOP ON MOLECULAR TECHNIQUES

WORKSHOP 1



(UPOV BMT/14 - Seoul, Republic of Korea, November 12, 2014)

Agreed:

 would be useful to develop a joint document explaining the principal features (e.g. DUS, variety identification, variety purity, etc.) of the systems of OECD, UPOV and ISTA.

WORKSHOP 2



(OECD Seed Schemes-Paris, June 8, 2016)

OECD Annual Meeting approved:

 develop a joint document explaining the principal features (e.g. DUS, variety identification, variety purity, etc.) of the systems of OECD, UPOV, AOSA and ISTA







WORKSHOP 1



Agreed:

 to propose an inventory by UPOV, OECD and ISTA of the use of molecular marker techniques, by crop, with a view to developing a document containing that information, in a similar format to UPOV document UPOV/INF/16 "Exchangeable Software".

WORKSHOP 2



OECD Annual Meeting approved:

to carry out a joint inventory by UPOV, OECD, AOSA and ISTA of the
use of molecular marker techniques, by crop, with a view to
developing a document containing that information. The OECD will
contribute to the document by sharing the ongoing list of molecular
techniques used by NDAs and continuously collected by the
Secretariat.







WORKSHOP 1



Agreed:

- to propose to invite UPOV, OECD and ISTA to develop lists of possible joint initiatives in relation to molecular techniques. It was noted that, in the case of UPOV, the list could be drafted by the BMT at its fifteenth session, subject to approval by the Technical Committee.
 - The UPOV TC, at its fifty-second session, agreed that the BMT should include the development of a list of terminology (definitions) used by OECD, UPOV and ISTA in the list of joint initiatives
 - The UPOV BMT, at its fifteenth session, agreed that possible future collaboration between UPOV, OECD and ISTA might include the harmonization of terms and methodologies used for different crops and the possible development of standards, after agreement by these organizations.







WORKSHOP 2



OECD Annual Meeting approved:

3. To develop a list of terms and their definitions as used by OECD, UPOV, AOSA and ISTA and to make an attempt to harmonise these.







WORKSHOP 1



Agreed:

4. would be useful for mutual understanding, to repeat the joint workshop at relevant meetings of the OECD and ISTA

WORKSHOP 2



OECD Annual Meeting approved:

4. to consider organising another similar workshop [in OECD] in three years time







UPOV BMT/15



- OECD, ISTA and UPOV had different objectives and cooperation between the organizations in the use of molecular techniques would need to reflect that. However, the BMT agreed that it would be important to explore circumstances in which the same techniques and information could be used. In the first instance, it agreed that it would be more effective to explore such possibilities on the basis of real situations rather than at a theoretical and institutional level.
- welcomed the proposal by the Netherlands to organize a practical workshop in 2017, with support from UPOV, OECD and ISTA, to explore how molecular techniques might be applied in an efficient way for UPOV, OECD and ISTA purposes.







WORKSHOP 2



OECD Annual Meeting approved:

 endorsed the proposal of the Netherlands to organise a practical workshop in 2017, with support of the OECD, UPOV and ISTA, to explore how molecular techniques might be applied in an efficient way for UPOV, OECD and ISTA purposes.

OVERVIEW

Molecular techniques

- Introduction to UPOV
- UPOV guidance on molecular techniques
- Coordination with other international organizations
- Possible future developments
- Summary (FAQs)

51

Working Group on Biochemical and Molecular Techniques, and DNA-Profiling in Particular

The BMT is a group open to DUS experts, biochemical and molecular specialists and plant breeders, whose role is to:

- Review general developments in biochemical and molecular techniques;
- Maintain an awareness of relevant applications of biochemical and molecular techniques in plant breeding;
- Consider the possible application of biochemical and molecular techniques in DUS testing and report its considerations to the TC;
- If appropriate, establish **guidelines** for biochemical and molecular **methodologies** and their harmonization;
- Develop guidelines regarding the management and harmonization of databases of biochemical and molecular information, in conjunction with the TWC;
- Provide a **forum for discussion** on the use of biochemical and molecular techniques in the consideration of **essential derivation** and **variety identification**.

Working Group on Biochemical and Molecular Techniques, and DNA-Profiling in Particular

The BMT is a group open to DUS experts, biochemical and molecular specialists and plant breeders, whose role is to:

- Review general developments in biochemical and molecular techniques;
- Maintain an awareness of relevant applications of biochemical and molecular techniques in plant breeding;
- Consider the possible application of biochemical and molecular techniques in DUS testing and report its considerations to the TC;
- If appropriate, establish guidelines for biochemical and molecular methodologies and their harmonization;
- Develop guidelines regarding the management and harmonization of databases of biochemical and molecular information, in conjunction with the TWC;
- Provide a **forum for discussion** on the use of biochemical and molecular techniques in the consideration of **essential derivation** and **variety identification**.

53


Presentation at the Fifty-First session of the Technical Committee Geneva, March 2015

http://www.upov.int/edocs/mdocs/upov/en/tc 51/tc 51 presentation 2.pdf



Marker-Assisted Selection of 'Similar Variety' in DUS Testing

March 2015.



OVERVIEW

Molecular techniques

- Introduction to UPOV
- UPOV guidance on molecular techniques
- Coordination with other international organizations
- Possible future developments
- Summary (FAQs)

Is it possible to obtain protection of a variety on the basis of its DNA-profile?

- For a variety to be protected, it needs to be clearly distinguishable from all existing varieties on the basis of characteristics that are physically expressed, e.g. plant height, time of flowering, fruit color, disease resistance etc.
- The DNA-profile is not the basis for obtaining the protection of a variety, although this information may be used as supporting information.
- A more detailed explanation is provided in the FAQ <u>Does</u> <u>UPOV allow molecular techniques (DNA profiles) in the</u> <u>examination of Distinctness</u>, <u>Uniformity and Stability ("DUS")?</u>

57

Question: Does UPOV allow molecular techniques (DNA profiles) in the DUS examination?

- It is important to note that, in some cases, varieties may have a
 different DNA profile but be phenotypically identical, whilst, in
 other cases, varieties which have a large phenotypic difference
 may have the same DNA profile for a particular set of molecular
 markers (e.g. some mutations).
- In relation to the use of molecular markers that are not related to phenotypic differences, the concern is that it might be possible to use a limitless number of markers to find differences between varieties at the genetic level that are not reflected in phenotypic characteristics.

On the above basis, UPOV has agreed the following uses in relation to DUS examination:

Question: Does UPOV allow molecular techniques (DNA profiles) in the DUS examination? (Cont'd)

- (a) Molecular markers can be used as a method of examining DUS characteristics that satisfy the criteria for characteristics set out in the General Introduction if there is a reliable link between the marker and the characteristic.
- (b) A combination of phenotypic differences and molecular distances can be used to improve the selection of varieties to be compared in the growing trial if the molecular distances are sufficiently related to phenotypic differences and the method does not create an increased risk of not selecting a variety in the variety collection which should be compared to candidate varieties in the DUS growing trial.

59

Questions

TGP/7: "Development of Test Guidelines"

Additional Information and guidance on Asterisked, grouping and TQ characteristics

Standard Test Guidelines Characteristic

Function	Criteria
1. Characteristics that are accepted by UPOV for examination of DUS and from which members of the Union can select those suitable for their particular	1. Must satisfy the criteria for use of any characteristic for DUS as set out in Chapter 4, section 4.2.
circumstances.	2.Must have been used to develop a variety description by at least one member of the Union .
	3. Where there is a long list of such characteristics and, where considered appropriate, there may be an indication of the extent of use of each characteristic.

Asterisked Characteristic

14 (*)	PQ	VG	(+)	(c)			
	distri	blade: bution of ndary color					
	none					Edward Goucher	1
	on ma	argin only				Wevo2	2
		inal zone				Keylib	3
	centra	al zone					4
	irregu	ılar				Francis Mason	5

Asterisked Characteristic

Function	Criteria
1.Characteristics that are important for the international harmonization of variety	1. Must be a characteristic included in the Test Guidelines.
descriptions.	2. Should always be examined for DUS and included in the variety description by all members of the Union
	EXCEPT when the state of expression of a preceding characteristic or regional environmental conditions render this inappropriate.
	3. Must be useful for function 1.
	4. Particular care should be taken before selection of disease resistance characteristics.

Grouping Characteristic

- Grouping of Varieties and Organization of the Growing Trial
- The selection of varieties of common knowledge to be grown in the trial with the candidate varieties and the way in which these varieties are divided into groups to facilitate the assessment of 5.1 distinctness are aided by the use of grouping characteristics.
- Grouping characteristics are those in which the documented states of expression, even where produced at different locations, can be used, either individually or in combination with other such characteristics: (a) to select varieties of common knowledge that can be excluded from the growing trial used for examination of distinctness; and (b) to organize the growing trial so that similar varieties are grouped together.
- The following have been agreed as useful grouping characteristics:
 - Plant: growth habit (characteristic 1)
 - Plant: height in relation to width (characteristic 2)

 - Young shoot: anthocyanin coloration (characteristic 5) Leaf blade: main color on upper side (characteristic 12) green yellow


green grey green purple green


Leaf blade: secondary color (characteristic 13)

white pinkish white yellow yellow red

Sepal: color (characteristic 20) pinkish white

Apple: Fruit color





TECHNICA	L QUESTIONNAIRE	Page {x} of {y}	Reference Number:
			Application date: (not to be filled in by the applicant)
	to be completed in	TECHNICAL QUESTIO connection with an application	NNAIRE ation for plant breeders' rights
1. Sub	ect of the Technical Questio	nnaire	
1.1	Genus	Plectranthus L'Hér	
1.2	Species		[]
(ple	ase complete)		
1.3	Hybrid		[]
	Species (please complete)		

TEG	CHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:	
5. con	Characteristics of the variety		ne number in brackets refers t ark the note which best correspo	
	Characteristics		Example Varieties	Note
5.5 (37)	Fruit: hue of over color – with bloom	n removed		
	orange red		Cox's Orange Pippin, Egremont Russet	1[]
	pink red		Cripps Pink, Delorgue	2[]
	red		Akane, Galaxy, Red Elstar, Regal Prince	3[]
	purple red		Red Jonaprince, Spartan	4[]
	brown red		Fiesta, Joburn, Lord Burghley	5[]
5.6 (39)	Fruit: pattern of over color			
	only solid flush		Red Jonaprince, Richared Delicious	1[]
	solid flush with weakly defined stripes		Galaxy	2[]
	solid flush with strongly defined stripe	s	Jonagored	3[]
	weakly defined flush with strongly def	fined stripes	Gravensteiner	4[]
	only stripes (no flush)		Helios	5[]
	flushed and mottled		Elstar	6[]
	flushed, striped and mottled		Jonagold	7[]

Grouping Characteristic

Function	Criteria		
acteristics in which the	1.(a) Qualitative characteristics or		
documented states of	(b) Quantitative or pseudo-qualitative		
expression , even where recorded	characteristics which provide useful		
at different locations, can be	discrimination between the varieties of commor		
used either individually or in	knowledge from documented states of		
combination with other such characteristics:	expression recorded at different locations.		
to select varieties of common knowledge that can be	2.Must be useful for functions 1 and 2.		
excluded from the growing trial used for examination of	3. Should be an asterisked characteristic and/or included in the Technical		
distinctness, and/or	Questionnaire or application form.		
to organize the growing trial so that similar varieties are grouped together			
	acteristics in which the documented states of expression, even where recorded at different locations, can be used either individually or in combination with other such characteristics: to select varieties of common knowledge that can be excluded from the growing trial used for examination of distinctness, and/or to organize the growing trial so		

Relationship between functions

- (a) **GROUPING CHARACTERISTICS** selected from the Table of Characteristics should, in general, **receive an asterisk** in the Table of Characteristics and be **included in the Technical Questionnaire**.
- (b) TQ CHARACTERISTICS selected from the Table of Characteristics should, in general, receive an asterisk in the Table of Characteristics and be used as grouping characteristics. TQ characteristics are not restricted to those characteristics used as grouping characteristics;
- (c) ASTERISKED CHARACTERISTICS are not restricted to those characteristics selected as grouping or TQ characteristics.

Questions

3. GUIDANCE ON DRAFTING TEST GUIDELINES (Document TGP/7)

UPOV provides guidance by:

- The "General Introduction" (TG/1/3)
 - General technical principles
 - Organization of DUS Testing
 - Associated "TGP" Documents (e.g. statistical methods)

AND

- "Test Guidelines"
 - Species/Crop-specific recommendations developed by crop experts
 - TGP/7 "Development of Test Guidelines" adopted

3. GUIDANCE ON DRAFTING TEST GUIDELINES

a) Subject of the Test Guidelines, Material Required and Method of Examination

Example

- 1. Subject of these Test Guidelines
- These Test Guidelines apply to all varieties of *Theobroma cacao* L.
- 2. Material Required
- 2.2 The material is to be supplied in the form of seed or plants.
- 2.3 The minimum quantity of plant material, to be supplied by the applicant, should be:
- seed-propagated varieties: 20 fresh seeds

vegetatively propagated varieties: 5 plants

38

76

Example

Method of Examination

- 3.1 Number of Growing Cycles
- 3.1.1 The minimum duration of tests should normally be two independent growing cycles.
- In particular, it is essential that the trees produce a satisfactory crop of fruit in each of the two growing cycles.
- 3.1.2 The growing cycle is considered to be the duration of a single growing season, beginning with vegetative growth, followed by flowering and fruit harvest.
- 3.4 Test Design
- 3.4.1 Each test should be designed to result in a total of at least 10 plants in the case of seed-propagated plants or, in the case of vegetatively propagated varieties, in a total of at least 5 plants.

Example

Method of Examination

Assessment of Distinctness

4.1.4 Number of Plants / Parts of Plants to be Examined Seed-propagated varieties: [...] all observations on single plants should be made on 10 plants or parts taken from each of 10 plants and any other observations made on all plants in the test, disregarding any off-type plants.

Vegetatively propagated varieties: [...] all observations should be made on 5 plants or parts taken from each of 5 plants, disregarding any off-type plants.

78

Example

Method of Examination

4.2 Uniformity

- 4.2.2 Vegetatively propagated varieties
- For the assessment of uniformity of vegetatively propagated varieties, a population standard of 1% and an acceptance probability of 95% should be applied. In the case of a sample size of 5 plants, no off-types are allowed.
- 4.2.3 Seed propagated varieties
- The assessment of uniformity for seed-propagated varieties should be according to the recommendations for cross-pollinated varieties in the General Introduction.

10 Chapters of UPOV Test Guidelines

- 1. Subject of the Test Guidelines
- 2. Material Required
- 3. Methods of Examination
- 4. Assessment of Distinctness, Uniformity and Stability
- 5. Grouping of Varieties and Organization of the Growing Trial
- 6. Introduction to the Table of Characteristics
- 7. Table of Characteristics
- 8. Explanation on the Table of Characteristics
- 9. Literature
- 10. Technical Questionnaire


TGP/7: "Development of Test Guidelines"

Section 3. Guidance for Drafting Test Guidelines

- •The TG Template
- Additional Standard Wording for the TG Template
- •Guidance Notes for the TG Template

Web-Based TG Template

Web-Based TG Template






Demonstration

3. GUIDANCE ON DRAFTING TEST GUIDELINES

b) Method of observation (MS, MG, VS, VG)



Method of Observation

M: Measurement:

an objective observation against a calibrated, linear scale e.g. using a ruler, weighing scales, colorimeter, dates, counts, etc.);

V: Visual observation:

includes observations where the expert uses reference points (e.g. diagrams, example varieties, side-by-side comparison) or non-linear charts (e.g. color charts).

"Visual" observation refers to the sensory observations of the expert and, therefore, also includes smell, taste and touch.

161	P/9 "Examining Distinctness" Type of expression of characteristic			
Method of propagation QL PQ QN of the variety (QUAL itative) (PSEUDO qualitative) (QUANT itative)				
Vegetatively propagated, self-pollinated	Notes (VG)	Notes (VG) Side-by-side (VG)	Notes (VG/MG/MS) Side-by-side (VG) Statistics (MG/MS)	
Cross-pollinated	Notes (VG) Statistics (VS*)	Notes (VG) Side-by-side (VG) Statistics (VS*)	Statistics ([MG]/MS/VS) Side-by-side (VG) Notes (VG/MG/MS)	
Hybrids	Notes (VG) Statistics (VS*)	Notes (VG) Side-by-side (VG) Statistics (VS*)	**	

TGP/9 "Examining Distinctne V= Visual observation			
	Туре о	f expression of characte	ristic
Method of propagatior of the variety	QL (QUAL itative)	PQ (PSEUDO qualitative)	QN (QUANT itative)
Vegetatively propagated, Self-pollinated	Notes (VG)	Notes (VG) Side-by-side (VG)	Notes (VG/MG/MS) Side-by-side (VG) Statistics (MG/MS)
Cross-pollinated	Notes (VG) Statistics (VS*)	Notes (VG) Side-by-side (VG) Statistics (VS*)	Statistics ([MG]/MS/VS) Side-by-side (VG) Notes (VG/MG/MS)
Hybrids	Notes (VG) Statistics (VS*)	Notes (VG) Side-by-side (VG) Statistics (VS*)	**


TGP/9 "Examining Distinctness" V= Visual observation or M= Measurement				
	Турє	e of expression of char	acteristic	
Method of propagation of the variety	QL (QUAL itative)	PQ (PSEUDO qualitative	QN (QUANT itative)	
Vegetatively propagated, self-pollinated	Notes (VG)	Notes (VG) Side-by-side (VG)	Notes (VG/MG/MS) Side-by-side (VG) Statistics (MG/MS)	
Cross-pollinated	Notes (VG) Statistics (VS*)	Notes (VG) Side-by-side (VG) Statistics (VS*)	Statistics ([MG]/MS/VS) Side-by-side (VG) Notes (VG/MG/MS)	
Hybrids	Notes (VG) Statistics (VS*)	Notes (VG) Side-by-side (VG) Statistics (VS*)	**	


Type of Record (for the purposes of distinctness)

G: single record for a variety, or a GROUP of plants or parts of plants;

In most cases, "G" provides a single record per variety and it is not possible or necessary to apply statistical methods in a plant-by-plant analysis for the assessment of distinctness.

<u>S</u>: records for a number of **SINGLE**, individual **plants** or parts of plants ...





Questions	
EXERCISE	

3. GUIDANCE ON DRAFTING TEST GUIDELINES

c) Types of Expression (QL, PQ, QN), notes and distinctness;

TYPE OF EXPRESSION OF CHARACTERISTICS (QL, QN, PQ)

Types of Expression

QL: QUALITATIVE

QN: QUANTITATIVE


PQ: PSEUDO-QUALITATIVE


Table of Characteristics/Tableau des caractères/Merkmalstabelle/Tabla de caracteres Example Varieties Exemples Beispielssorten Variedades ejemplo English français deutsch español Note/ 1. (*) PQ Plant: growth habit Plante : port Pflanze: Wuchsform Planta: hábito de crecimiento Edward Goucher erguido semi-upright Minaud semi dressé halbaufrecht semierguido 2 rounded Golden Panache 3 breitwüchsig extendido 4 2. (* QN VG Pflanze: Höhe im Verhältnis zur Breite nant: height in Plante : hauteur par Planta: altura en rapport à la largeur relation to width relación con la taller than broad plus haute que large höher als breit más alta que ancha Edward Goucher, 1 Sherwood as tall as broad aussi haute que large gleich hoch wie breit tan alta como ancha Golden Panache 2 broader than tall plus large que haute más ancha que alta Rupestri 3 breiter als hoch QN VG 3.


QUALITATIVE Characteristics

"Qualitative characteristics" are those that are **expressed in discontinuous states** (e.g. sex of plant: dioecious female (1), dioecious male (2), monoecious unisexual (3), monoecious hermaphrodite (4)).

These states are self-explanatory and independently meaningful. All states are necessary to describe the full range of the characteristic, and every form of expression can be described by a single state. The order of states is not important. As a rule, the **characteristics are not influenced by environment**.









QUANTITATIVE Characteristics

"Quantitative characteristics" are those where the expression covers the full range of variation from one extreme to the other. The **expression can be recorded on a one-dimensional, continuous or discrete, linear scale**. The range of expression is divided into a number of states for the purpose of description (e.g. length of stem: very short (1), short (3), medium (5), long (7), very long (9)). The division seeks to provide, as far as is practical, an even distribution across the scale. The Test Guidelines do not specify the difference needed for distinctness. The states of expression should, however, be meaningful for DUS assessment.











PSEUDO-QUALITATIVE Characteristics

In the case of "pseudo-qualitative characteristics," the **range of expression is at least partly continuous, but varies in more than one dimension** (e.g. shape: ovate (1), elliptic (2), circular (3), obovate (4)) and cannot be adequately described by just defining two ends of a linear range. In a similar way to qualitative (discontinuous) characteristics — hence the term "pseudo-qualitative" — each individual state of expression needs to be identified to adequately describe the range of the characteristic.









EXERCISE


NOTES and DISTINCTNESS according to TYPE OF EXPRESSION (QL, PQ, QN)

Types of Expression

QL: QUALITATIVE

QN: QUANTITATIVE

PQ: PSEUDO-QUALITATIVE



	Qua	alitative C	Characteri	stics	
		(specia	ıl cases)		
Char both of the No. Wethord of No.	français	deutsch	español	Example Varieties/ Exemples/ Beispielssorten/ Variedades ejemplo	Note/ Nota
1. MS Plant: ploidy (*) C					
QL diploid tetraploid					4
3. VG Stem: anthoc (*) coloration	yanin				
QL absent present				Gumpoong Chunpoong, Gopoong	9

Qualitative Characteristics: **distinctness**

In qualitative characteristics, the difference between two varieties may be considered clear if one or more characteristics have expressions that fall into **two different states in the Test Guidelines**. Varieties should not be considered distinct for a qualitative characteristic if they have the same state of expression.

(e.g. sex of plant: dioecious female (1), dioecious male (2), monoecious unisexual (3), monoecious hermaphrodite (4)).

Types of Expression


QL: QUALITATIVE

QN: QUANTITATIVE


PQ: PSEUDO-QUALITATIVE

PSEUDO-QUALITATIVE Characteristics

In the case of "pseudo-qualitative characteristics," the **range of expression is at least partly continuous, but varies in more than one dimension** (e.g. shape: ovate (1), elliptic (2), circular (3), obovate (4)) and cannot be adequately described by just defining two ends of a linear range. In a similar way to qualitative (discontinuous) characteristics — hence the term "pseudo-qualitative" — each individual state of expression needs to be identified to adequately describe the range of the characteristic.



		(ty	ypical ex	amples)	
24. (+)	Flower: color of the center	Fleur: couleur du centre	Farbe der Mitte	Flor: color del centro	
PQ	green	vert	grün	verde	1
	yellow	jaune	gelb	amarillo	2
	orange	orange	orange	naranja	3
	pink	rose	rosa	rosa	4
	red	rouge	rot	гојо	5
	purple	pourpre	purpurn	púrpura	6



Types of Expression

QL: QUALITATIVE

QN: QUANTITATIVE


PQ: PSEUDO-QUALITATIVE


QUANTITATIVE Characteristics

"Quantitative characteristics" are those where the expression covers the full range of variation from one extreme to the other. The **expression can be recorded on a one-dimensional, continuous or discrete, linear scale**. The range of expression is divided into a number of states for the purpose of description (e.g. length of stem: very short (1), short (3), medium (5), long (7), very long (9)). The division seeks to provide, as far as is practical, an even distribution across the scale. The Test Guidelines do not specify the difference needed for distinctness. The states of expression should, however, be meaningful for DUS assessment.

Quantitative Characteristics: **distinctness**

Quantitative characteristics are considered for distinctness according to the method of observation and the features of propagation of the variety concerned...





Quantitative Characteristics (1-9)

weak/strong short/long small/large

Note	State
1	very weak
	(or: absent or very weak)
2	very weak to weak
3	weak
4	weak to medium
5	medium
6	medium to strong
7	strong
8	strong to very strong
9	very strong

Note	State
1	very small
	(or: absent or very small)
2	very small to small
3	small
4	small to medium
5	medium
6	medium to large
7	large
8	large to very large
9	very large

Quantitative Characteristics (1-9)

Standard Range Version 1	Standard Range Version 2		
1 very weak	1 very weak		
(or: absent or very weak)	(or: absent or very weak)		
3 weak	3 weak		
5 medium	5 medium		
7 strong	7 strong		
9 very strong	-		

Standard Range Version 3	Standard Range Version 4	
-	-	
3 weak	3 weak	
5 medium	5 medium	
7 strong	7 strong	
9 very strong		

Quantitative Characteristics (1-9)

State	Example 1	Example 1 Example 2 E		Example 4	
	Size relative to:	Angle:	Position:	Length in relation to:	
1	much smaller	very acute	at base	equal	
3	moderately smaller	moderately acute	one quarter from base	slightly shorter	
5	same size	right angle	in middle	moderately shorter	
7	moderately larger	moderately obtuse	one quarter from apex end	much shorter	
9	much larger	very obtuse	at apex	very much shorter	

Quantitative Characteristics (at least 3 notes)

Ex	Example 2		
1	e.g. absent or weak		
	(absent or weakly expressed)		
2	moderate (or medium)		
	(moderately expressed)		
3	strong		
l	(strongly expressed)		

State	Example 1			
	Stem: attitude			
1	erect			
3	semi-erect			
5	prostrate			

NOTES

Versus

SIDE-BY-SIDE COMPARISON

(Quantitative characteristics)

TGP/9 "Examining Distinctness"

- 5.2 Approaches for assessing distinctness
- 5.2.1 Introduction
- 5.2.1.1 Approaches for assessment of distinctness based on the growing trial can be summarized as follows:
 - (a) **Side-by-side visual comparison** in the growing trial (see Section 5.2.2);
 - (b) Assessment by Notes / single variety records ("Notes"): the assessment of distinctness is based on the recorded state of expression of the characteristics of the variety

(see Section 5.2.3);

(c) Statistical analysis of growing trial data:

Quantitative Characteristics: distinctness


The General Introduction explains that, in the case of visually observed quantitative characteristics:

"5.5.2.2.2 A direct comparison between two similar varieties is always recommended, since direct pairwise comparisons are the most reliable. In each comparison, a difference between two varieties is acceptable as soon as it can be assessed visually and could be measured, although such measurement might be impractical or require unreasonable effort."

TGP/9 "Examining Distinctness"

5.2.3.1.2 Where the requirements for distinctness assessment by Notes / single variety records are met it would usually also be possible to make a side-by-side visual comparison. However, in the case of assessment by Notes / single variety records, such proximity is not required, which is a particular advantage where the growing trial contains a large number of varieties and where there are limited possibilities for ensuring that all similar varieties are grouped together in the growing trial. ...

On the other hand, because the varieties are not the subject of a side-by-side visual comparison, the difference required between varieties as a basis for distinctness is, with the exception of qualitative characteristics (see below), somewhat greater.




Quantitative Characteristics: **distinctness**


Quantitative characteristics are considered for distinctness according to the method of observation and the features of propagation of the variety concerned.

Test Guidelines (TGP/7)

Difference of **two Notes to represent a clear difference if** the **comparison** between two varieties is performed **at the level of Notes**:

WHY?





Quantitative Characteristics: **distinctness**

Quantitative characteristics are considered for distinctness according to the method of observation and the features of propagation of the variety concerned.

Test Guidelines (TGP/7)


Difference of **two Notes to represent a clear difference if** the **comparison** between two varieties is performed **at the level of Notes**:

Quantitative Characteristics: **distinctness**

35.	QN	MG/VG	(+)					
	Time of flo	of beginning wering						
	early						3	
	medi	um				Minaud	5	
	late					Golden Panache	7	

1 to 9 scale: Notes 1 and 3, Notes 2 and 4, Notes 3 and 5 etc. represent a clear difference

Quantitative Characteristics: **distinctness** TG/233/1 Diascia/Diascie, 2007-03-28 Example Varieties/ Exemples/ Note/ English français Deutsch español Beispielssorten/ Variedades ejemplo Stem: anthocyanin Tige: pigmentation Tallo: pigmentación anthocyanique sous inflorescence coloration below Anthocyanfärbung antociánica por debajo de la inflorescence unter dem Blütenstand absent or weak absente ou faible fehlend oder gering ausente o débil Heccharm medium moyenne mitte1 media Hecrace 1 to 3 scale: only Notes 1 and 3 represent a clear difference



Questions


3. GUIDANCE ON DRAFTING TEST GUIDELINES


d) Shape and Color Characteristics

TGP/14: Shape

Characteristics related to shape, could use the following components:

- Shape: e.g. ovate (1), elliptic (2), circular (3), obovate
 (4)...
- Ratio length/ width (from low to high)
- · Position of broadest part
- Shape of base
- Shape of apex
- · Lateral outline





shape	very compressed	moderately compressed	slightly compressed	medium	slightly elongated	moderately elongated	very elongated	
ratio length/width	very low	low	low to medium		medium to high		very high	
Parallel set						•	0	
oblong	12		10				(1)	
Rounded set				_		^	^	٨
ovate	\bigcirc	\bigcirc	4			\bigcirc		0
-Weste			1				\bigcap	Λ
elliptic	8	7	46	V 5	\vee	Ψ	Ψ	Ψ
abanata.								0
obovate	~	4	4	V	\bigvee		V	Y
Angular set						Λ	A	Λ
triangular	\Rightarrow	4	4	4	\triangle	4	4	4
			\wedge		\bigwedge	Λ	Λ	٨
trullate	\Leftrightarrow	\Leftrightarrow	\Diamond	\Diamond	\Diamond	\Diamond	Ψ	1
			\wedge			\wedge	Λ	/
rhombic	\Leftrightarrow	\Diamond	\Diamond	\forall	\forall	4	\Diamond	1
	~						\triangle	^
obtrullate	~	1	1	\forall	\bigvee	V	Y	Y
	>							n
obtriangular	'	1	V	V	\bigvee	V	Y	V

TGP/14: Shape



Alternative 1:

Alternative 2:



ratio length/width:

Shape: broad obovate



ratio length/width: medium

Shape: medium obovate



ratio length/width: high 3

Shape: narrow obovate



broadest part towards base



elliptic



broadest part towards apex 3 obovate

TGP/14: Shape




Alternative 1


- position of broadest part (QN):

 e.g. strongly towards base (1); moderately towards base (3); at middle (5); moderately towards apex (7); strongly towards apex (9)
- ratio length/width (QN):
 e.g., xerx low (1); low (3); medium (5); high (7); very high (9); (b)

Alternative 2 General shape (PC): triangular (1); ovate (2); circular (3); elliptic (4); oblong (5); linear (6); obovate (7); obtainceolate (6); apailutate (6); obtainspular (10) (Note: Where the overall shapes presented as a single pseudo-qualitative characteristic, the order of states should be: primary order broadest part above middle; seconday order, broad to hair one (low to high rate tenghishidh). The proadest part above middle (2); obtain to hair one (low to high rate tenghishidh). (Bottom middle) The proadest part above middle (2); obtain the property of the proadest part above middle (3); obtain the proad

EXERCISE





		TGP.	/14: Color
		state of expression	example
	low	single color	yellow, orange, red
sion	,	color rango	(a) yellow, yellow orange, orange, orange red, red
f precis		color range	(b) white, yellowish white, yellow, yellowish orange
level of precision	↓	intensity	light yellow, medium yellow, dark yellow
	high	RHS Colour Chart No.	RHS 41 B
	·		Species?
		Leve	el of variation?

TGP/14: Color Single color

- A single color has the lowest precision to describe the state of expression.
- Example: Flower: color: white (1); yellow (2); orange (3); red (4)

TGP/14: Color

Color range

- (a) In color combinations the second color indicates the predominant color with blending of both colors, resulting in what can look like a single color. For example in "green red" the predominant color is red and in "red green" the predominant color is green.
- Example: Flower: color: white (1); yellow white (2); yellow (3); yellow orange (4); orange (5)
- (b) The use of "ish" in color combinations indicates that there is a predominant color (e.g. yellow) together with another minor color. For example,
- Example: Flower: color: whitish (1); yellowish (2); greenish (3)

TGP/14: Color Intensity

- Depending on the organ described, the intensity can be presented either in relation to a single color or in combination with different colors (example 2).
- Example 1: Leaf: green color of upper side: light (3); medium (5); dark (9)
- Example 2: Flower: color: white (1); light yellow (2); medium yellow (3); dark yellow (4); orange (5)

TGP/14: Color Color Chart

- The "RHS Colour Chart" because of its worldwide availability.
 UPOV names for colors in document TGP/14: ANNEX.
- "Because daylight varies, color determinations made against a color chart should be made either in a suitable cabinet providing artificial daylight or in the middle of the day in a room without direct sunlight. The spectral distribution of the illuminant for artificial daylight should conform with the CIE Standard of Preferred Daylight D 6500 and should fall within the tolerances set out in the British Standard 950, Part I. These determinations should be made with the plant part placed against a white background".
- Observations should not be made in direct sunlight. The observations should be made on a cloudy day with sufficient light intensity, or in a shaded area.

Allocation of UPOV	Color Groups for e	<u>ach RHS Color in</u>	<u>ı RHS Refer</u>	ence order
RHS COLORS (RHS	COLOUR CHART,	, EDITIONS 1986	, 1995, 200	1 AND 2007)
•	BY UPOV CO	OR GROUPS		,

UPOV Group No.	No. RHS	English	français	deutsch	español
11	001A	yellow	jaune	gelb	amarillo
5	001B	yellow green	vert-jaune	gelbgrün	verde amarillento
5	001C	yellow green	vert-jaune	gelbgrün	verde amarillento
5	001D	yellow green	vert-jaune	gelbgrün	verde amarillento
11	002A	yellow	jaune	gelb	amarillo
11	002B	yellow	jaune	gelb	amarillo
5	002C	yellow green	vert-jaune	gelbgrün	verde amarillento
5	002D	yellow green	vert-jaune	gelbgrün	verde amarillento
11	003A	yellow	jaune	gelb	amarillo
11	003B	yellow	jaune	gelb	amarillo
11	003C	yellow	jaune	gelb	amarillo
5	003D	yellow green	vert-jaune	gelbgrün	verde amarillento
11	004A	yellow	jaune	gelb	amarillo
11	004B	yellow	jaune	gelb	amarillo
5	004C	yellow green	vert-jaune	gelbgrün	verde amarillento
10	004D	light yellow	jaune clair	hellgelb	amarillo claro
11	005A	yellow	jaune	gelb	amarillo
11	005B	yellow	jaune	gelb	amarillo
11	005C	yellow	jaune	gelb	amarillo
10	005D	light yellow	jaune clair	hellgelb	amarillo claro
11	006A	yellow	jaune	gelb	amarillo
11	006B	yellow	jaune	gelb	amarillo
11	006C	yellow	jaune	gelb	amarillo
10	006D	light yellow	jaune clair	hellgelb	amarillo claro
11	007A	yellow	jaune	gelb	amarillo
11	007B	yellow	jaune	gelb	amarillo
11	007C	yellow	jaune	gelb	amarillo
11	007D	vellow	iaune	aelb	amarillo

TGP/14: Color APPROACHES TO DESCRIBE COLORS AND COLOR PATTERNS

- depends on the number of colors...
- the types of color distribution...
- and the number of color patterns possible for the species concerned.

TGP/14: Color

Approach according to the size of the surface area

- (a) only a few colors, a few types of color distribution and a few patterns to be described,
- the colors are described according to the size of the surface area they cover

"The main color is the color with the largest surface area. In cases where the areas of the main and secondary color are too similar to reliably decide which color has the largest area, [the darkest color] / [the color...[location]Q] is considered to be the main color."

TGP/14: Color Approach according to tissue layers

- one layer is covering the other:
- (a) Ground color (not always the largest surface area):
 - (i) the first color to appear chronologically.
 - (ii) has a continuous dispersion across the surface.
- (b) Over color (not always occupying the smallest surface area):
 - a second color, such as a flush, spots or blotches developed over time.

APPLE - TG/14/9

35. (*)		Fruit: ground color	
PQ	(f)	not visible	1
		whitish yellow	2
		yellow	3
		whitish green	4
		yellow green	5
		green	6

37. (*)		Fruit: hue of over color – with bloom removed	
PQ	(f)	orange red	1
		pink red	2
		red	3
		purple red	4
		brown red	5

Phalaenopsis (TG/213/2(proj.7))




Petal: ground color – RHS Colour Chart 155A - white Petal: over color – RHS Colour Chart 83A – dark violet

TGP/14: Color

Approach according to defined parts of an organ

- (a) If the different parts of a plant organ can have different colors, the color of these different parts can be described separately.
- Example:
 - Petal: color of margin
 - Petal: color of middle zone
 - Petal: color of base
- (b) When an organ has one color with different intensities, the parts
 of the organ which are lighter or darker could be described as follows:
- · Example:
 - Ray floret: color distribution on upper side:
 - lighter towards base (1); even (2); lighter towards apex (3)




Approach according to the RHS Colour Chart number ("Lisbon" approach)

- All colors of the plant part concerned are assessed using the RHS Colour Charts first.
- The color should first be described, followed by:
 - distribution,
 - pattern
 - area,
 - conspicuousness of the color (if necessary).
- The same sequence should be followed for color two, color three and so on.

Heuchera and Heucherella (TG/280/1)

- 36. Leaf blade: color one RHS Colour Chart Yellow-Green 144C
- 37. Leaf blade: color one: distribution marginal zone (7)
- · 38. Leaf blade: color one: pattern solid or nearly solid (5)
- 39. Leaf blade: color one: total area very small to small (2)
- 40. Leaf blade: color two RHS Colour Chart Greyed-Orange 176B
- 41: Leaf blade: color two: distribution along veins (2)
- 42: Leaf blade: color two: pattern solid or nearly solid (5)
- 43: Leaf blade: color two: total area small (3)
- · 44: Leaf blade: color three RHS Colour Chart Greyed-Orange 177D but more grey
- 45: Leaf blade: color three: distribution between veins in intermediate zone (6)
- · 46: Leaf blade: color three: pattern solid or nearly solid (5)
- 47: Leaf blade: color three: total area large (7)
- 48: Leaf blade: color four RHS Colour Chart not applicable
- 49: Leaf blade: color four: distribution none (1)
- 50: Leaf blade: color four: pattern not applicable
- 51: Leaf blade: color four: total area not applicable



TGP/14: Color Order of states of expression

- normally presented in the following order: white, green, yellow, orange, pink, red, purple, violet, blue, brown, black
- chronological appearance of the color (e.g. as the fruit ripens)

Questions

3. GUIDANCE ON DRAFTING TEST GUIDELINES

e) Example Varieties


TG/13/9 Lettuce/Laitue/Salat/Lechuga, 2004-03-31 - 7 -


7. <u>Table of Characteristics/Tableau des caractères/Merkmalstabelle/Tabla de caracteres</u>

	English	français	Deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
1. (*)	Seed: color	Semence: couleur	Samen: Farbe	Semilla: color		
	white	blanche	weiß	blanco	Verpia	1
	yellow	jaune	gelb	amarillo	Durango	2
	black	noire	schwarz	negro	Kagraner Sommer	3
2. (*) (+)	Seedling: anthocyanin coloration	Plantule: pigmentation anthocyanique	Keimpflanze: Anthocyanfärbung	Plántula: pigmentación antociánica		
	absent	absente	fehlend	ausente	Verpia	1
	present	présente	vorhanden	presente	Pirat	9
3.	Seedling: size of cotyledon (fully developed)	Plantule: taille du cotylédon (à complet développement)	Keimpflanze: Größe des Keimblatts (voll entwickelt)	Plántula: tamaño del cotiledón (plenamente desarrollado)		
	small	petit	klein	pequeño	Romance	3
	medium	moyen	mittel	medio	Expresse	5
	large	grand	groß	grande	Verpia	7

		English	français	deutsch	español	Example Varieties/ Exemples/ Beispielssorten/ Variedades ejemplo	Note Note
14.	VG	Leaf blade: intensity of purplish color of <u>lower</u> side		Blattspreite: Intensität der Purpurfarbe der Unterseite	Limbo: intensidad del color purpúreo del envés		
QN	(a)	very light	très claire	sehr hell	muy claro		1
		light	claire	hell	claro	Perlime	3
		medium	moyenne	mittel	medio		5
		dark	foncée	dunkel	oscuro	Perro	7
		very dark	très foncée	sehr dunkel	muy oscuro	Bora, Purple	9
15.	VG	Leaf blade: profile	Limbe: profil	Blattspreite: Profil	Limbo: perfil		
QN	(a)	concave	concave	konkav	cóncavo	Perro	3
		plane	plan	flach	plano	Pergro, Saeyeupsil	5
		convex	convexe	konvex	convexo		7

			Brachyscome/Bl	TG/223/1 aues Gänseblümchen, 2 - 7 -	2005-04-06		
7.	Table	e of Characteristics	Tableau des caracté	eres/Merkmalstabel	le/Tabla de caracte	eres	
		English	français	deutsch	español	Example Varieties/ Exemples/ Beispielssorten/ Variedades ejemplo	Note/ Nota
1. (*) (+)		Plant: growth type	Plante: type de croissance	Pflanze: Wuchstyp	Planta: tipo de crecimiento		
QL	(a)	basal clusters	en amas à la base	basale Büschel	en racimos basales	l	1
		bushy	buissonnant	buschig	arbustivo		2
2. (+)		Only varieties with bushy growth type: Plant: predominant attitude of stems	Variétés à type de croissance huissonnant uniquement: Plante: port le plus fréquent des tiges	Nur Sorten mit buschigem Wuchstyn: Pflanze: vorwiegende Haltung der Triebe	Sólo variedades con tipo de crecimiento arbustivo: Planta: porte predominante de los tallos		
QN	(a)	upright	dressées	aufrecht	erecto	l	1
		semi upright	demi-dressées	halbaufrecht	semierecto	l	3
		horizontal	horizontales	waagerecht	horizontal		5
3.		Only varieties with bushy growth type: Plant: number of stems	Variétés à type de croissance buissonnant uniquement: Plante: nombre de tiges	Nur Sorten mit buschigem Wuchstyn: Pflanze: Anzahl Triebe	Sólo variedades con tipo de crecimiento arbustivo: Planta: número de tallos		
QN	(a)	few	peu nombreuses	klein	bajo		3
		medium	moyennement nombreuses	mittel	medio		5
		many	nombreuses	groß	alto		7
4. (*) (+)		Plant: height including flowers	Plante: hauteur, fleurs comprises	Pflanze: Höhe einschließlich Blüten	Planta: altura, i incluidas las flores		
QN	(a)	short	basse	niedrig	corta	Mardi Gras	3
		medium	moyenne	mittel	media	Breakoday	5
		tall	élevée	hoch	larga	Happy Face Pink	7










in characteristics used to **harmonize descriptions**

NEED < and

which are influenced by the environment







Questions

3. GUIDANCE ON DRAFTING TEST GUIDELINES

f) The process for developing UPOV Test Guidelines, including: TG Template; Additional Standard Wording; and Guidance Notes;

Genera and Species

- >3,450 genera and species with varieties examined for PBR
- >3,305 genera and species for which UPOV members have practical DUS experience
- 321Test Guidelines adopted

Note: 321 Test Guidelines estimated to cover 92% of PBR-related varieties in UPOV Plant Variety Database

PRIORITY for UPOV Test Guidelines

PRIORITY for species or crops with high:

- number of authorities receiving PBR applications;
- number of PBR applications;
- number of foreign applications received by UPOV members;
- economic importance;
- level of breeding activity

EXAMPLE (New Test Guidelines)

Test Guidelines: *Plantus magnifica* L.

(Common name: Alpha)

Technical Working Party: TWX

TWX (2014):
TWX (2015):
Alpha (proj.1)
Alpha (proj.2)
Alpha (proj.3)
Alpha (proj.3)
Alpha (proj.4)
Alpha (proj.4)
Alpha (proj.5)
Final adopted document (2017):
TG/500/1

TGP/7: "Development of Test Guidelines"

Procedure for the Introduction and Revision of UPOV Test Guidelines

Proposals (New TG, Revisions, Corrections)

Criteria to be observed

Approval

Preparation of draft TG for the TWP

Leading Expert

Subgroup

Submission to the TC

Requirements for "final" draft

Consideration by the TC-EDC

Adoption of the TG by the TC

4. AGENDA for the TWP Session

Sunday	Mon	ıday	Tues	sday	Wedn	nesday	Thur	sday	Friday
[TECHNICAL WORKSHOP] (optional)	Reports on developmen	Reports on developments in PVP		TGP document development		development		s with new pecies ominations	Databases, Electronic application systems Exchangeable software
COFFEE	COF	FEE	COF	FEE	COFFEE		COF	FEE	COFFEE
[TECHNICAL WORKSHOP] (optional)	Reports (Co Molecular te	· ·		TGP document development		Room 1 Room 2 Uniformity method development			Recommendations on Test Guidelines
	LUN	СH	LUN	СН	LUI	NCH	LUN	СH	LUNCH
PREPARATORY WORKSHOP	Room 1 Test Guidelines subgroup	Room 2 Test Guidelines subgroup	Room 1 Test Guidelines subgroup	Room 2 Test Guidelines subgroup			Room 1 Test Guidelines subgroup	Room 2 Test Guidelines subgroup	Future program Adoption of report
COFFEE	COF	FEE	COF	FEE	TECHNIC	CAL VISIT	COF	FEE	
PREPARATORY WORKSHOP	Room 1 Test Guidelines subgroup	Room 2 Test Guidelines subgroup	Room 1 Test Guidelines subgroup	Room 2 Test Guidelines subgroup			Test Guidelines subgroup Subgroup		END OF SESSION
	Contin	nuation	RECEPTION				Contin	nuation	

	Monday, June 19 Start 9.00	Tuesday, June 20 Start 8.30	Wednesday, June 21 Early Start 8.00	Thursday, Start		Friday, June 23 Start 8.30
08.30	Start 9.00 0.00: 1_Opening 2_Allocation of the agenda (TWA46/1 Rex.) 26_Date and place of next session 3_(a) Reports from members and observers (TWA46/3) - Increasing participation of new members of the from (TWE/1/18)	8. TGP documents (contd) -Data processing for distinctness and producing descriptions (TWPPILIS) -Assessing uniformity by off-types on more than one growing cycle (TWPILIT Rev.) -B. Uniformity by off-types (TWA4694 and TWA4694 Add.) 14. Number of arrowing cycles in DUS examination (TWPILIS). TWA4692 and TWA4694 and TWA694	Room 1 Quinca (DK)	8. Information and dai (b) Variety description (TWPII2) 17. Development of It excluding varieties fro cycle when COVD is 1 18. Statistical method observed characterist 19. Image analysis 20. Management of v. (TWPII/14)	databases resholds for m second growing used (TWP/1/22) s for visually ics (TWP/1/23) WP/1/10) ariety collections	5. Molecular Techniques (TWP/II/) 7. Variety denominations (TWP/II/6) 8. Information and databases (Contd) (a) UPOV information databases (TWP/II/6) (c) Exchange and use of software and equipment (TWP/II/6) (d) Electronic application systems (TWP/II/6) 24. Recommendations on draft Test
	3 (b) Report on developments within UPOV (TWA/46/2)	Minimum distance between varieties (TWA/46/6 and TWA/46/6 Add.) TGP documents (cont'd)	9.45 – 10.15 COFFEE	21. Software for statistical analysis (TWP/1/18) 22. Matters to be resolved concerning TGs adopted by the TC (TC/53/31)		-New proposals for Test Guidelines
		-Illustrations for shape and ratio characteristics (TWP/1/18)	3.43 - 10.13 COFFEE	10.15 - 10.45		27. Future program
10.30	COFFEE 4. Organization of the UPOV sessions (TWP1/12) 5. TGP documents (TWP1/11 Rev.) -Confidentially of molecular information (TWP1/12) -Drustion of DDS (TWP1/11) -The COYU orterion (TWP1/11) -The COYU orterion (TWP1/12) 10. Esperiences with new types and sessions	COFFEE 1. TGP documents (cont.) Characteristics which only apply to certain varieties (CWFILIS) 2. Guidance for drafters of Test Suidelines (TWFILIS) 12. Impact of endophytes on DUS characteristics in prasses (TWAHIDE and TWAHIDE AND America (TWAHIDE AND America (TWAHIDE) 15. Beginnal set of example varieties in Wheat for South America (TWAHIDE) 15. Use of disease and insect resistance characteristics in DUS examination (TWAHIDE) LUNCH	Room 1 Soya Bean (AR) 12.00 – 13.00 EARLY LUNCH	Room 1 Outs (ES)		COFFEE 28. Adoption of resont 29. Closing of the session
14.00	Room 1 *Barley (DE)	Room 1 *Cotton (ES)	Departure from hotel 13.00 Field Trip:	Room 1 Rice (JP)	Room 2 *Elytrigia (AR)	Closing 1pm
15.45	COFFEE	COFFEE		COFF	EE	1
16.00	Room 1 Red Clover (ZA)	Room 1 "Field Bean (GB)		Room 1 Castor Bean (ZA)	Room 2 Ginseng (KR)	
7.45						1
18.00	Reserve	Official dinner (informal) 18.00 to 22.30	Return to hotel: 20.00	Rese	rve	

EXCHANGING INFORMATION

AN OPPORTUNITY for TRAINING