



TG/36/7(proj.1)

ORIGINAL: English

DATE: 2020-05-08

INTERNATIONAL UNION FOR THE PROTECTION OF NEW VARIETIES OF PLANTS

Geneva

DRAFT

RAPE SEED

UPOV Code(s): BRASS_NAP_NUS

Brassica napus L. oleifera

GUIDELINES

FOR THE CONDUCT OF TESTS

FOR DISTINCTNESS, UNIFORMITY AND STABILITY

*prepared by experts from the United Kingdom
to be considered by the
Technical Working Party for Agricultural Crops
at its forty-ninth session, to be held in Saskatoon, Canada,
from 2020-06-22 to 2020-06-26*

Disclaimer: this document does not represent UPOV policies or guidance

Alternative names:*

<i>Botanical name</i>	<i>English</i>	<i>French</i>	<i>German</i>	<i>Spanish</i>
<i>Brassica napus L. oleifera</i>	Oilseed Rape, Rapeseed, Swede Rape	Colza	Raps	Colza

The purpose of these guidelines ("Test Guidelines") is to elaborate the principles contained in the General Introduction (document TG/1/3), and its associated TGP documents, into detailed practical guidance for the harmonized examination of distinctness, uniformity and stability (DUS) and, in particular, to identify appropriate characteristics for the examination of DUS and production of harmonized variety descriptions.

ASSOCIATED DOCUMENTS

These Test Guidelines should be read in conjunction with the General Introduction and its associated TGP documents.

* These names were correct at the time of the introduction of these Test Guidelines but may be revised or updated. [Readers are advised to consult the UPOV Code, which can be found on the UPOV Website (www.upov.int), for the latest information.]

TABLE OF CONTENTS	PAGE
1. SUBJECT OF THESE TEST GUIDELINES.....	4
2. MATERIAL REQUIRED.....	4
3. METHOD OF EXAMINATION.....	4
3.1 Number of Growing Cycles.....	4
3.2 Testing Place.....	4
3.3 Conditions for Conducting the Examination.....	4
3.4 Test Design.....	5
3.5 Additional Tests.....	5
4. ASSESSMENT OF DISTINCTNESS, UNIFORMITY AND STABILITY.....	5
4.1 Distinctness.....	5
4.2 Uniformity.....	6
4.3 Stability.....	7
5. GROUPING OF VARIETIES AND ORGANIZATION OF THE GROWING TRIAL.....	8
6. INTRODUCTION TO THE TABLE OF CHARACTERISTICS.....	8
6.1 Categories of Characteristics.....	8
6.2 States of Expression and Corresponding Notes.....	8
6.3 Types of Expression.....	9
6.4 Example Varieties.....	9
6.5 Legend.....	10
7. TABLE OF CHARACTERISTICS/TABLEAU DES CARACTÈRES/MERKMALSTABELLE/TABLA DE CARACTERES.....	11
8. EXPLANATIONS ON THE TABLE OF CHARACTERISTICS.....	19
8.1 Explanations covering several characteristics.....	19
8.2 Explanations for individual characteristics.....	20
9. LITERATURE.....	25
10. TECHNICAL QUESTIONNAIRE.....	26

1. Subject of these Test Guidelines

These Test Guidelines apply to all varieties of *Brassica napus* L. *oleifera*.

2. Material Required

2.1 The competent authorities decide on the quantity and quality of the plant material required for testing the variety and when and where it is to be delivered. Applicants submitting material from a State other than that in which the testing takes place must ensure that all customs formalities and phytosanitary requirements are complied with.

2.2 The material is to be supplied in the form of seed.

2.3 The minimum quantity of plant material, to be supplied by the applicant, should be:

Candidate: 300g
Component of controlled-cross pollination hybrids: 100g

The seed should meet the minimum requirements for germination, species and analytical purity, health and moisture content, specified by the competent authority. In cases where the seed is to be stored, the germination capacity should be as high as possible and should be stated by the applicant.

2.4 The plant material supplied should be visibly healthy, not lacking in vigor, nor affected by any important pest or disease.

2.5 The plant material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If it has been treated, full details of the treatment must be given.

3. Method of Examination

3.1 *Number of Growing Cycles*

3.1.1 The minimum duration of tests should normally be two independent growing cycles.

3.1.2 The two independent growing cycles should be in the form of two separate plantings.

3.1.3 The testing of a variety may be conducted when the competent authority can determine with certainty the outcome of the test.

3.2 *Testing Place*

Tests are normally conducted at one place. In the case of tests conducted at more than one place, guidance is provided in TGP/9 "Examining Distinctness".

3.3 *Conditions for Conducting the Examination*

3.3.1 The tests should be carried out under conditions ensuring satisfactory growth for the expression of the relevant characteristics of the variety and for the conduct of the examination.

3.3.2 The optimum stage of development for the assessment of each characteristic is indicated by a number in the Table of Characteristics. The stages of development denoted by each number are described in Chapter 8.

3.4 *Test Design*

3.4.1 Each test should be designed to result in a total of at least 200 plants, which should be divided between at least 2 replicates.

3.4.2 The design of the tests should be such that plants or parts of plants may be removed for measurement or counting without prejudice to the observations which must be made up to the end of the growing cycle.

3.5 *Additional Tests*

Additional tests, for examining relevant characteristics, may be established.

4. Assessment of Distinctness, Uniformity and Stability

4.1 *Distinctness*

4.1.1 General Recommendations

It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding distinctness. However, the following points are provided for elaboration or emphasis in these Test Guidelines.

To assess distinctness of hybrids, the parent lines and the formula may be used according to the following recommendations:

(i) description of parent lines according to the Test Guidelines;

(ii) check of the originality of the parent lines in comparison with the variety collection, based on the characteristics in Chapter 7, in order to identify similar parent lines;

(iii) check of the originality of the hybrid formula in relation to the hybrids in the variety collection, taking into account the most similar lines; and

(iv) assessment of the distinctness at the hybrid level for varieties with a similar formula.

Further guidance is provided in documents TGP/9 "Examining Distinctness" and TGP/8 "Trial Design and Techniques Used in the Examination of Distinctness, Uniformity and Stability".

4.1.2 Consistent Differences

The differences observed between varieties may be so clear that more than one growing cycle is not necessary. In addition, in some circumstances, the influence of the environment is not such that more than a single growing cycle is required to provide assurance that the differences observed between varieties are sufficiently consistent. One means of ensuring that a difference in a characteristic, observed in a growing trial, is sufficiently consistent is to examine the characteristic in at least two independent growing cycles.

4.1.3 Clear Differences

Determining whether a difference between two varieties is clear depends on many factors, and should consider, in particular, the type of expression of the characteristic being examined, i.e. whether it is expressed in a qualitative, quantitative, or pseudo-qualitative manner. Therefore, it is important that users of these Test Guidelines are familiar with the recommendations contained in the General Introduction prior to making decisions regarding distinctness.

4.1.4 Number of Plants or Parts of Plants to be Examined

Unless otherwise indicated, for the purposes of distinctness, all observations on single plants should be made on 45 plants or parts of plants taken from each of 45 plants and any other observations made on all plants in the test, disregarding any off-type plants.

In the case of observations of parts taken from single plants, the number of parts to be taken from each of the plants should be 1.

4.1.5 Method of Observation

The recommended method of observing the characteristic for the purposes of distinctness is indicated by the following key in the Table of Characteristics (see document TGP/9 "Examining Distinctness", Section 4 "Observation of characteristics"):

MG: single measurement of a group of plants or parts of plants

MS: measurement of a number of individual plants or parts of plants

VG: visual assessment by a single observation of a group of plants or parts of plants

VS: visual assessment by observation of individual plants or parts of plants

Type of observation: visual (V) or measurement (M)

"Visual" observation (V) is an observation made on the basis of the expert's judgment. For the purposes of this document, "visual" observation refers to the sensory observations of the experts and, therefore, also includes smell, taste and touch. Visual observation includes observations where the expert uses reference points (e.g. diagrams, example varieties, side-by-side comparison) or non-linear charts (e.g. color charts). Measurement (M) is an objective observation against a calibrated, linear scale e.g. using a ruler, weighing scales, colorimeter, dates, counts, etc.

Type of record: for a group of plants (G) or for single, individual plants (S)

For the purposes of distinctness, observations may be recorded as a single record for a group of plants or parts of plants (G), or may be recorded as records for a number of single, individual plants or parts of plants (S). In most cases, "G" provides a single record per variety and it is not possible or necessary to apply statistical methods in a plant-by-plant analysis for the assessment of distinctness.

In cases where more than one method of observing the characteristic is indicated in the Table of Characteristics (e.g. VG/MG), guidance on selecting an appropriate method is provided in document TGP/9, Section 4.2.

4.2 *Uniformity*

4.2.1 It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding uniformity. However, the following points are provided for elaboration or emphasis in these Test Guidelines:

4.2.2 These Test Guidelines have been developed for the examination of seed-propagated. For varieties with other types of propagation, the recommendations in the General Introduction and document TGP/13 "Guidance for new types and species" Section 4.5 "Testing Uniformity" should be followed.

4.2.3 The assessment of uniformity for seed-propagated varieties should be according to the recommendations for cross-pollinated varieties in the General Introduction.

4.2.4 The assessment of uniformity for hybrid varieties depends on the type of hybrid and should be according to the recommendations for hybrid varieties in the General Introduction.

4.2.5 Where the assessment of a hybrid variety involves the parent lines, the uniformity of the hybrid variety should, in addition to an examination of the hybrid variety itself, also be assessed by examination of the uniformity of its parent lines.

4.2.6 For the assessment of uniformity of inbred varieties and component lines of hybrid varieties, a population standard of 2% and an acceptance probability of at least 95% should be applied. In the case of a sample size of 200 plants, 7 off-types are allowed. For the assessment of uniformity of single and multiple-cross hybrid varieties, a population standard of 10% and an acceptance probability of at least 95% should be applied. In the case of a sample size of 200 plants, 27 off-types are allowed.

4.3 *Stability*

4.3.1 In practice, it is not usual to perform tests of stability that produce results as certain as those of the testing of distinctness and uniformity. However, experience has demonstrated that, for many types of variety, when a variety has been shown to be uniform, it can also be considered to be stable.

- 4.3.2 Where appropriate, or in cases of doubt, stability may be further examined by testing a new seed stock to ensure that it exhibits the same characteristics as those shown by the initial material supplied.
- 4.3.3 Where appropriate, or in cases of doubt, the stability of a hybrid variety may, in addition to an examination of the hybrid variety itself, also be assessed by examination of the uniformity and stability of its parent lines.

5. Grouping of Varieties and Organization of the Growing Trial

- 5.1 The selection of varieties of common knowledge to be grown in the trial with the candidate varieties and the way in which these varieties are divided into groups to facilitate the assessment of distinctness are aided by the use of grouping characteristics.
- 5.2 Grouping characteristics are those in which the documented states of expression, even where produced at different locations, can be used, either individually or in combination with other such characteristics: (a) to select varieties of common knowledge that can be excluded from the growing trial used for examination of distinctness; and (b) to organize the growing trial so that similar varieties are grouped together.
- 5.3 The following have been agreed as useful grouping characteristics:
- (a) Seed: erucic acid (characteristic 1)
 - (b) Leaf: lobes (characteristic 13)
 - (c) Time of flowering (characteristic 19)
 - (d) Flower: color of petals (characteristic 20)
 - (e) Production of pollen (characteristic 24)
 - (f) Plant: total length (characteristic 25)
- 5.4 Guidance for the use of grouping characteristics, in the process of examining distinctness, is provided through the General Introduction and document TGP/9 "Examining Distinctness".

6. Introduction to the Table of Characteristics

6.1 *Categories of Characteristics*

6.1.1 Standard Test Guidelines Characteristics

Standard Test Guidelines characteristics are those which are approved by UPOV for examination of DUS and from which members of the Union can select those suitable for their particular circumstances.

6.1.2 Asterisked Characteristics

Asterisked characteristics (denoted by *) are those included in the Test Guidelines which are important for the international harmonization of variety descriptions and should always be examined for DUS and included in the variety description by all members of the Union, except when the state of expression of a preceding characteristic or regional environmental conditions render this inappropriate.

6.2 *States of Expression and Corresponding Notes*

- 6.2.1 States of expression are given for each characteristic to define the characteristic and to harmonize descriptions. Each state of expression is allocated a corresponding numerical note for ease of recording of data and for the production and exchange of the description.

6.2.2 In the case of qualitative and pseudo-qualitative characteristics (see Chapter 6.3), all relevant states of expression are presented in the characteristic. However, in the case of quantitative characteristics with 5 or more states, an abbreviated scale may be used to minimize the size of the Table of Characteristics. For example, in the case of a quantitative characteristic with 9 states, the presentation of states of expression in the Test Guidelines may be abbreviated as follows:

State	Note
small	3
medium	5
large	7

However, it should be noted that all of the following 9 states of expression exist to describe varieties and should be used as appropriate:

State	Note
very small	1
very small to small	2
small	3
small to medium	4
medium	5
medium to large	6
large	7
large to very large	8
very large	9

6.2.3 Further explanation of the presentation of states of expression and notes is provided in document TGP/7 "Development of Test Guidelines".

6.3 *Types of Expression*

An explanation of the types of expression of characteristics (qualitative, quantitative and pseudo-qualitative) is provided in the General Introduction.

6.4 *Example Varieties*

Where appropriate, example varieties are provided to clarify the states of expression of each characteristic.

6.5 Legend

		English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
1	2	3	4	5	6	7	
	Name of characteristics in English		Nom du caractère en français	Name des Merkmals auf Deutsch	Nombre del carácter en español		
	states of expression		types d'expression	Ausprägungsstufen	tipos de expresión		

- 1 Characteristic number
- 2 (*) Asterisked characteristic – see Chapter 6.1.2
- 3 Type of expression
 - QL Qualitative characteristic – see Chapter 6.3
 - QN Quantitative characteristic – see Chapter 6.3
 - PQ Pseudo-qualitative characteristic – see Chapter 6.3
- 4 Method of observation (and type of plot, if applicable)
 - MG, MS, VG, VS – see Chapter 4.1.5
- 5 (+) See Explanations on the Table of Characteristics in Chapter 8.2
- 6 (a)-(d) See Explanations on the Table of Characteristics in Chapter 8.1
- 7 Growth stage key See Explanations on the Table of Characteristics in Chapter 8

7. Table of Characteristics/Tableau des caractères/Merkmalstabelle/Tabla de caracteres

	English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
1. (*)	QL MG	(+)			00	
	Seed: erucic acid					
	low					1
	high					9
2.	QN MG/MS	(+)	(a)		15-17	
	Cotyledon: ratio length/width					
	small					3
	medium					5
	large					7
3.	QN MS	(+)	(a)		15-17	
	Cotyledon: saddle depth					
	small					3
	medium					5
	large					7
4.	QN MS	(+)	(a)		15-17	
	Cotyledon: ratio lobe separation/width					
	small					3
	medium					5
	large					7
5.	QN MG/MS	(+)	(a)		15-17	
	Cotyledon: lamina base to wide point/width ratio					
	small					3
	medium					5
	large					7
6.	QN MS	(+)	(a)		15-17	
	Cotyledon: ratio lobe separation/saddle depth					
	small					3
	medium					5
	large					7

	English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
7.	QN MG/MS	(+) (a)	15-17			
	Cotyledon: length/lamina length ratio					
	small					3
	medium					5
	large					7
8.	QN MG/MS	(+) (a)	15-17			
	Cotyledon: lamina base to wide point					
	small					3
	medium					5
	large					7
9.	QN VG	(+)	23-27			
	Plant: growth habit					
	erect					1
	intermediate					2
	prostrate					3
10. (*)	QN VG		23-27			
	Leaf: green color					
	light					3
	medium					5
	dark					7
11.	QL VG		23-27			
	Leaf: glaucosity					
	absent					1
	present					9
12.	QN VG		23-27			
	Leaf: intensity of glaucosity					
	weak					1
	medium					2
	strong					3

	English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
13. (*)	QL VG	(b)	23-27			
	Leaf: lobes					
	absent					1
	present					9
14. (*)	QN MS/VG	(+) (b)	23-27			
	Leaf: number of lobes					
	few					3
	medium					5
	many					7
15. (*)	QN VG	(+)	23-27			
	Leaf: dentation of margin					
	weak					3
	medium					5
	strong					7
16.	QN MG		23-27			
	Propose to DELETE: Leaf: length (blade and petiole)					
	short					3
	medium					5
	long					7
17.	QN MG		23-27			
	Proposed to DELETE Leaf: width (widest point)					
	narrow					3
	medium					5
	broad					7
18.	QN MG		23-27			
	<u>Proposed to DELETE</u> <u>Varieties with lobed leaves only:</u> Leaf: length of petiole					
	short					3
	medium					5
	long					7

	English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
19. (*)	QN MG/MS	(+)				
	Time of flowering					
	very early					1
	early					3
	medium					5
	late					7
	very late					9
20. (*)	PQ VG					
	Flower: color of petals					
	white					1
	cream					2
	yellow					3
	orange-yellow					4
21.	QN MS/VG	(c)	62-63			
	Flower: length of petals					
	short					3
	medium					5
	long					7
22.	QN MS/VG	(c)	62-63			
	Flower: width of petals					
	narrow					3
	medium					5
	broad					7
23.	QN MS/VG	(c)	62-63			
	Flower: ratio length/width of petals					
	small					3
	medium					5
	large					7
24. (*)	QL VG		62-63			
	Production of pollen					
	absent					1
	present					9

	English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
25. (*)	QN MG/MS	(+)	75-80			
	Plant: total length					
	short					3
	medium					5
	long					7
26.	QN MS	(+)	(d)	75-89		
	Siliqua: length					
	short					3
	medium					5
	long					7
27.	QN MS		(d)	75-89		
	Siliqua: length of beak					
	short					3
	medium					5
	long					7
28.	QN MS		(d)	75-89		
	Siliqua: length of peduncle					
	short					3
	medium					5
	long					7
29.	QN MS		(d)	75-89		
	Siliqua: width					
	narrow					3
	medium					5
	broad					7
30.	QN MS		(d)	75-89		
	Siliqua: ratio length/width					
	small					3
	medium					5
	large					7

	English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
31.	QN VG	(+)				
	Tendency to form inflorescences when sown in alternate season					
	absent or very weak					1
	weak					3
	medium					5
	strong					7
	very strong					9

8. Explanations on the Table of Characteristics

8.1 *Explanations covering several characteristics*

Characteristics containing the following key in the Table of Characteristics should be examined as indicated below:

- (a) The measurements should be taken on cotyledons of 40 seedlings. If the two cotyledons differ in size, the biggest one should be measured.

The length (characteristic no. 3) is defined as distance between the inclination at top of the cotyledon and the point where the width of the petiole is about 4 mm.

INSERT PICTURE

- (b) INSERT PICTURE

- (c) Picture required

- (d) All observations on the siliqua should be recorded in the midpart of the inflorescence of the main stem.


Picture required

8.2 *Explanations for individual characteristics*


Ad. 1: Seed: erucic acid

The erucic acid content should be observed on seed sent in by the applicant. It should be expressed as a percentage by mass of methyl esters in accordance with the ISO standard in document 5508, paragraph 6.2.2.1. Seed containing 2% or less would be classified as "low".


Ad. 2: Cotyledon: ratio length/width




Ad. 3: Cotyledon: saddle depth




Ad. 4: Cotyledon: ratio lobe separation/width




Ad. 5: Cotyledon: lamina base to wide point/width ratio




Ad. 6: Cotyledon: ratio lobe separation/saddle depth



Ad. 7: Cotyledon: length/lamina length ratio



Ad. 8: Cotyledon: lamina base to wide point



Ad. 9: Plant: growth habit

Picture required

Ad. 14: Leaf: number of lobes

To be observed on fully developed leaves.

Ad. 15: Leaf: dentation of margin

Picture required

Ad. 19: Time of flowering

The observation should be done at least three times per week and more frequently if there is any need to do so. A plant is at flowering when one flower is fully open. When assessing on the plot as a whole, the time of flowering should be recorded when 10% of plants have reached this stage. When assessed on individual plants, time of flowering is reached when 50% of plants have reached this stage.

Ad. 25: Plant: total length

The measurement should be taken from the base of the plant to the longest point (including side branches).

Ad. 26: Siliqua: length

To be measured between peduncle and beak.

Ad. 31: Tendency to form inflorescences when sown in alternate season

The tendency to form inflorescences in the year of sowing of winter rape varieties should be recorded in spring sown trials; that of spring rape varieties in late summer sown trials. The observation of the growth stage reached of winter rape varieties should be made in summer when the late spring rape varieties are flowering; that of spring rape varieties in autumn, when their development stagnates.

The design of the test should result in a total of at least 100 plants.

8.3 INSERT KEY FOR GROWTH STAGES

9. Literature

INSERT REFERENCE FOR GROWTH STAGE KEY

10. Technical Questionnaire

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
		Application date: (not to be filled in by the applicant)
TECHNICAL QUESTIONNAIRE to be completed in connection with an application for plant breeders' rights		
In the case of hybrid varieties which are the subject of an application for plant breeders' rights, and where the parent lines are to be submitted as a part of the examination of the hybrid variety, this Technical Questionnaire should be completed for each of the parent lines, in addition to being completed for the hybrid variety.		
1. Subject of the Technical Questionnaire		
1.1	Botanical name	<input type="text" value="Brassica napus L. oleifera"/>
1.2	Common name	<input type="text" value="Rape Seed"/>
2. Applicant		
	Name	<input type="text"/>
	Address	<input type="text"/>
	Telephone No.	<input type="text"/>
	Fax No.	<input type="text"/>
	E-mail address	<input type="text"/>
	Breeder (if different from applicant)	<input type="text"/>
3. Proposed denomination and breeder's reference		
	Proposed denomination (if available)	<input type="text"/>
	Breeder's reference	<input type="text"/>

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

#4. Information on the breeding scheme and propagation of the variety

4.1 Breeding scheme

Variety resulting from:

4.1.1 Crossing

(a) controlled cross

(please state parent variety)

(.....) x (.....)

female parent

male parent

(b) partially known cross

(c) unknown cross

4.1.2 Mutation
(please state parent variety)

4.1.3 Discovery and development
(please state where and when discovered and how developed)

4.1.4 Hybrid

MSL

Ogura

Other - please specify


4.1.5 Other
(Please provide details)

4.2 Method of propagating the variety


4.2.1 Other []
 (Please provide details)

In the case of hybrid varieties the production scheme for the hybrid should be provided on a separate sheet. This should provide details of all the parent lines required for propagating the hybrid e.g.

Single Hybrid



Three-Way Hybrid



and should identify in particular:

- (a) any male sterile lines
- (b) maintenance system of male sterile lines.

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

5. Characteristics of the variety to be indicated (the number in brackets refers to the corresponding characteristic in Test Guidelines; please mark the note which best corresponds).		
Characteristics	Example Varieties	Note
5.1 Seed: erucic acid (1)		
low		1 []
high		9 []
5.2 Leaf: lobes (13)		
absent		1 []
present		9 []
5.3 Time of flowering (19)		
very early		1 []
early		3 []
medium		5 []
late		7 []
very late		9 []
5.4 Flower: color of petals (20)		
white		1 []
cream		2 []
yellow		3 []
orange-yellow		4 []
5.5 Production of pollen (24)		
absent		1 []
present		9 []
5.6 Plant: total length (25)		
short		3 []
medium		5 []
long		7 []

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

6. Similar varieties and differences from these varieties

Please use the following table and box for comments to provide information on how your candidate variety differs from the variety (or varieties) which, to the best of your knowledge, is (or are) most similar. This information may help the examination authority to conduct its examination of distinctness in a more efficient way.

Denomination(s) of variety(ies) similar to your	Characteristic(s) in which your candidate variety differs	Describe the expression of the characteristic(s) for the	Describe the expression of the characteristic(s) for your
<i>Example</i>			
<p>Comments:</p>			

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

#7. Additional information which may help in the examination of the variety

7.1 In addition to the information provided in sections 5 and 6, are there any additional characteristics which may help to distinguish the variety?

Yes No

(If yes, please provide details)

7.2 Are there any special conditions for growing the variety or conducting the examination?

Yes No

(If yes, please provide details)

7.3 Other information

Seasonal type:
Winter
Spring

Use:
Forage
Oil

Miscellaneous information:
Dwarf-type
Clearfield tolerant
Clubroot resistance
Other (please specify)

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

8. Authorization for release

(a) Does the variety require prior authorization for release under legislation concerning the protection of the environment, human and animal health?

Yes [] No []

(b) Has such authorization been obtained?

Yes [] No []

If the answer to (b) is yes, please attach a copy of the authorization.

9. Information on plant material to be examined or submitted for examination

9.1 The expression of a characteristic or several characteristics of a variety may be affected by factors, such as pests and disease, chemical treatment (e.g. growth retardants or pesticides), effects of tissue culture, different rootstocks, scions taken from different growth phases of a tree, etc.

9.2 The plant material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If the plant material has undergone such treatment, full details of the treatment must be given. In this respect, please indicate below, to the best of your knowledge, if the plant material to be examined has been subjected to:

(a) Microorganisms (e.g. virus, bacteria, phytoplasma)	Yes []	No []
(b) Chemical treatment (e.g. growth retardant, pesticide)	Yes []	No []
(c) Tissue culture	Yes []	No []
(d) Other factors	Yes []	No []

Please provide details for where you have indicated "yes".

.....

10. I hereby declare that, to the best of my knowledge, the information provided in this form is correct:

Applicant's name

Signature Date

[End of document]