



TG/186/2(proj.2)

ORIGINAL: English

DATE: 2021-05-06

INTERNATIONAL UNION FOR THE PROTECTION OF NEW VARIETIES OF PLANTS

Geneva

DRAFT

SUGARCANE

UPOV Code(s): SACCH

Saccharum L.

GUIDELINES

FOR THE CONDUCT OF TESTS

FOR DISTINCTNESS, UNIFORMITY AND STABILITY

*prepared by experts from Australia
to be considered by the
Technical Working Party for Agricultural Crops
at its fiftieth session, to be held in Arusha, United Republic of Tanzania,
from 2021-06-21 to 2021-06-25*

Disclaimer: this document does not represent UPOV policies or guidance

Alternative names:*

<i>Botanical name</i>	<i>English</i>	<i>French</i>	<i>German</i>	<i>Spanish</i>
<i>Saccharum</i> L.	Sugarcane	Canne à sucre	Zuckerrohr	Caña de azúcar

The purpose of these guidelines ("Test Guidelines") is to elaborate the principles contained in the General Introduction (document TG/1/3), and its associated TGP documents, into detailed practical guidance for the harmonized examination of distinctness, uniformity and stability (DUS) and, in particular, to identify appropriate characteristics for the examination of DUS and production of harmonized variety descriptions.

ASSOCIATED DOCUMENTS

These Test Guidelines should be read in conjunction with the General Introduction and its associated TGP documents.

* These names were correct at the time of the introduction of these Test Guidelines but may be revised or updated. [Readers are advised to consult the UPOV Code, which can be found on the UPOV Website (www.upov.int), for the latest information.]

TABLE OF CONTENTS	PAGE
1. SUBJECT OF THESE TEST GUIDELINES.....	3
2. MATERIAL REQUIRED.....	3
3. METHOD OF EXAMINATION.....	3
3.1 Number of Growing Cycles.....	3
3.2 Testing Place.....	3
3.3 Conditions for Conducting the Examination.....	3
3.4 Test Design.....	4
3.5 Additional Tests.....	4
4. ASSESSMENT OF DISTINCTNESS, UNIFORMITY AND STABILITY.....	4
4.1 Distinctness.....	4
4.2 Uniformity.....	5
4.3 Stability.....	5
5. GROUPING OF VARIETIES AND ORGANIZATION OF THE GROWING TRIAL.....	6
6. INTRODUCTION TO THE TABLE OF CHARACTERISTICS.....	6
6.1 Categories of Characteristics.....	6
6.2 States of Expression and Corresponding Notes.....	6
6.3 Types of Expression.....	6
6.4 Example Varieties.....	6
6.5 Legend.....	8
7. TABLE OF CHARACTERISTICS/TABLEAU DES CARACTÈRES/MERKMALSTABELLE/TABLA DE CARACTERES.....	9
8. EXPLANATIONS ON THE TABLE OF CHARACTERISTICS.....	23
8.1 Explanations covering several characteristics.....	23
8.2 Explanations for individual characteristics.....	24
9. LITERATURE.....	32
10 TECHNICAL QUESTIONNAIRE.....	33

1. Subject of these Test Guidelines

These Test Guidelines apply to all varieties of *Saccharum L.*

2. Material Required

2.1 The competent authorities decide on the quantity and quality of the plant material required for testing the variety and when and where it is to be delivered. Applicants submitting material from a State other than that in which the testing takes place must ensure that all customs formalities and phytosanitary requirements are complied with.

2.2 The material is to be supplied in the form of vegetative cuttings which are about 8 to 12 months old.

2.3 The minimum quantity of plant material, to be supplied by the applicant, should be:

12 segments of culm with 3 buds each, properly packaged to minimize damage to the buds

2.4 The plant material supplied should be visibly healthy, not lacking in vigor, nor affected by any important pest or disease.

2.5 The plant material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If it has been treated, full details of the treatment must be given.

3. Method of Examination

3.1 *Number of Growing Cycles*

3.1.1 The minimum duration of tests should normally be a single growing cycle.

3.1.2 The testing of a variety may be concluded when the competent authority can determine with certainty the outcome of the test.

3.2 *Testing Place*

Tests are normally conducted at one place. In the case of tests conducted at more than one place, guidance is provided in TGP/9 "Examining Distinctness".

3.3 *Conditions for Conducting the Examination*

The tests should be carried out under conditions ensuring satisfactory growth for the expression of the relevant characteristics of the variety and for the conduct of the examination.

3.4 *Test Design*

Each test should be designed to result in a total of at least 24 culms, which should be divided between at least 2 replicates.

3.5 *Additional Tests*

Additional tests, for examining relevant characteristics, may be established.

4. Assessment of Distinctness, Uniformity and Stability

4.1 *Distinctness*

4.1.1 General Recommendations

It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding distinctness. However, the following points are provided for elaboration or emphasis in these Test Guidelines.

4.1.2 Consistent Differences

The differences observed between varieties may be so clear that more than one growing cycle is not necessary. In addition, in some circumstances, the influence of the environment is not such that more than a single growing cycle is required to provide assurance that the differences observed between varieties are sufficiently consistent. One means of ensuring that a difference in a characteristic, observed in a growing trial, is sufficiently consistent is to examine the characteristic in at least two independent growing cycles.

4.1.3 Clear Differences

Determining whether a difference between two varieties is clear depends on many factors, and should consider, in particular, the type of expression of the characteristic being examined, i.e. whether it is expressed in a qualitative, quantitative, or pseudo-qualitative manner. Therefore, it is important that users of these Test Guidelines are familiar with the recommendations contained in the General Introduction prior to making decisions regarding distinctness.

4.1.4 Number of Plants or Parts of Plants to be Examined

Unless otherwise indicated, for the purposes of distinctness, all observations on single plants should be made on 12 plants or parts of plants taken from each of 12 plants and any other observations made on all plants in the test, disregarding any off-type plants.

4.1.5 Method of Observation

The recommended method of observing the characteristic for the purposes of distinctness is indicated by the following key in the Table of Characteristics (see document TGP/9 "Examining Distinctness", Section 4 "Observation of characteristics"):

MG: single measurement of a group of plants or parts of plants

MS: measurement of a number of individual plants or parts of plants

VG: visual assessment by a single observation of a group of plants or parts of plants

VS: visual assessment by observation of individual plants or parts of plants

Type of observation: visual (V) or measurement (M)

"Visual" observation (V) is an observation made on the basis of the expert's judgment. For the purposes of this document, "visual" observation refers to the sensory observations of the experts and, therefore, also includes smell, taste and touch. Visual observation includes observations where the expert uses reference points (e.g. diagrams, example varieties, side-by-side comparison) or non-linear charts (e.g. color charts). Measurement (M) is an objective observation against a calibrated, linear scale e.g. using a ruler, weighing scales, colorimeter, dates, counts, etc.

Type of record: for a group of plants (G) or for single, individual plants (S)

For the purposes of distinctness, observations may be recorded as a single record for a group of plants or parts of plants (G), or may be recorded as records for a number of single, individual plants or parts of plants (S). In most cases, "G" provides a single record per variety and it is not possible or necessary to apply statistical methods in a plant-by-plant analysis for the assessment of distinctness.

In cases where more than one method of observing the characteristic is indicated in the Table of Characteristics (e.g. VG/MG), guidance on selecting an appropriate method is provided in document TGP/9, Section 4.2.

4.2 *Uniformity*

- 4.2.1 It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding uniformity. However, the following points are provided for elaboration or emphasis in these Test Guidelines:
- 4.2.2 These Test Guidelines have been developed for the examination of vegetatively propagated varieties. For varieties with other types of propagation, the recommendations in the General Introduction and document TGP/13 "Guidance for new types and species" Section 4.5 "Testing Uniformity" should be followed.
- 4.2.3 For the assessment of uniformity of vegetatively propagated varieties, a population standard of 1% and an acceptance probability of at least 95% should be applied. In the case of a sample size of 24 plants, 1 off-type is allowed.

4.3 *Stability*

- 4.3.1 In practice, it is not usual to perform tests of stability that produce results as certain as those of the testing of distinctness and uniformity. However, experience has demonstrated that, for many types of variety, when a variety has been shown to be uniform, it can also be considered to be stable.
- 4.3.2 Where appropriate, or in cases of doubt, stability may be further examined by testing a new seed or plant stock to ensure that it exhibits the same characteristics as those shown by the initial material supplied.

5. Grouping of Varieties and Organization of the Growing Trial

- 5.1 The selection of varieties of common knowledge to be grown in the trial with the candidate varieties and the way in which these varieties are divided into groups to facilitate the assessment of distinctness are aided by the use of grouping characteristics.
- 5.2 Grouping characteristics are those in which the documented states of expression, even where produced at different locations, can be used, either individually or in combination with other such characteristics: (a) to select varieties of common knowledge that can be excluded from the growing trial used for examination of distinctness; and (b) to organize the growing trial so that similar varieties are grouped together.
- 5.3 The following have been agreed as useful grouping characteristics:
- (a) Internode: cross section (characteristic 8)
 - (b) Internode: color where not exposed to sun (characteristic 10)
 - (c) Node: presence of wing on bud (characteristic 20)
 - (d) Node: shape of bud (characteristic 21)
- 5.4 Guidance for the use of grouping characteristics, in the process of examining distinctness, is provided through the General Introduction and document TGP/9 "Examining Distinctness".

6. Introduction to the Table of Characteristics

6.1 *Categories of Characteristics*

6.1.1 Standard Test Guidelines Characteristics

Standard Test Guidelines characteristics are those which are approved by UPOV for examination of DUS and from which members of the Union can select those suitable for their particular circumstances.

6.1.2 Asterisked Characteristics

Asterisked characteristics (denoted by *) are those included in the Test Guidelines which are important for the international harmonization of variety descriptions and should always be examined for DUS and included in the variety description by all members of the Union, except when the state of expression of a preceding characteristic or regional environmental conditions render this inappropriate.

6.2 *States of Expression and Corresponding Notes*

6.2.1 States of expression are given for each characteristic to define the characteristic and to harmonize descriptions. Each state of expression is allocated a corresponding numerical note for ease of recording of data and for the production and exchange of the description.

6.2.2 All relevant states of expression are presented in the characteristic.

6.2.3 Further explanation of the presentation of states of expression and notes is provided in document TGP/7 "Development of Test Guidelines".

6.3 *Types of Expression*

An explanation of the types of expression of characteristics (qualitative, quantitative and pseudo-qualitative) is provided in the General Introduction.

6.4 *Example Varieties*

Where appropriate, example varieties are provided to clarify the states of expression of each characteristic.

6.5 Legend

		English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
1	2	3	4	5	6	7	
		Name of characteristics in English	Nom du caractère en français	Name des Merkmals auf Deutsch	Nombre del carácter en español		
		states of expression	types d'expression	Ausprägungsstufen	tipos de expresión		

- 1 Characteristic number
- 2 (*) Asterisked characteristic – see Chapter 6.1.2
- 3 Type of expression
 QL Qualitative characteristic – see Chapter 6.3
 QN Quantitative characteristic – see Chapter 6.3
 PQ Pseudo-qualitative characteristic – see Chapter 6.3
- 4 Method of observation (and type of plot, if applicable)
 MG, MS, VG, VS – see Chapter 4.1.5
- 5 (+) See Explanations on the Table of Characteristics in Chapter 8.2
- 6 (a)-(d) See Explanations on the Table of Characteristics in Chapter 8.1
- 7 Not applicable

7. Table of Characteristics/Tableau des caractères/Merkmalstabelle/Tabla de caracteres

	English		français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
1.	QN	VG	(+)				
	Plant: stool growth habit						
	erect					Q 121, Q186	1
	semi-erect					Q96, RB72-454	3
	intermediate					Q168	5
	semi-prostrate					H56-752	7
	prostrate						9
2. (*)	QN	VS	(+)				
	Plant: adherence of leaf sheath						
	weak					H56-752, Q96	3
	medium					Q124, Q186	5
	strong					NC0 310, Q120, Q201	7
3.	QN	VG					
	Plant: number of tillers						
	few					Q124	3
	medium					RB72-454	5
	many					Q138	7
4. (*)	QN	MS	(+)				
	Culm: height						
	short					Q117	3
	medium					Q124, Q138, Q170	5
	long					Q136, RB72-454	7
5.	QN	MS	(+)	(a)			
	Internode: length on the bud side						
	short					Q117	3
	medium					Q138, Q170	5
	long					Q124	7
6. (*)	QN	MS	(+)	(a)			
	Internode: diameter						
	thin					Q136	3
	medium					H56-752, Q124, Q170	5
	thick					Q117	7

	English		français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
7. (*)	PQ	VG	(+)				
	Internode: shape						
		cylindrical				Q169, RB72-454	1
		tumescens				Q205	2
		bobbin-shaped				H56-752	3
		conoidal				Q177, Q178	4
		obconoidal				H60-3802	5
		concave-convex				Q115	6
8. (*)	QN	VG					
	Internode: cross section						
		circular				Q 121, RB72-454	1
		circular to ovate					2
		ovate				Q152, Q186, Q96	3
9. (*)	PQ	VG	(+)				
	Internode: color where <u>exposed</u> to sun						
		yellow				Q230	1
		yellow green				SRA24, SRA25	2
		grey yellow				SRA10	3
		grey orange				Q165	4
		grey red					5
		grey purple				RB72-454	6
		purple					7
10 (*)	PQ	VG	(+)				
	Internode: color where <u>not exposed</u> to sun						
		yellow					1
		yellow green				SRA24, SRA25	2
		grey yellow				QS01-1078	3
		grey orange				Q220	4
		grey red					5
		grey purple				SRA9	6
		purple					7

	English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
11	QN	MS/VS				
	Internode: number of growth crack					
	absent or very few				H56-752, RB72-454	1
	few				Q124	3
	medium				Q121	5
	many				Q179	7
	very many					9
12 (*)	QN	VG	(+)			
	Internode: expression of zigzag alignment					
	absent or very weak				Q124	1
	weak				Q135, Q152	3
	moderate				Q117	5
	strong				H56-752	7
13	QN	VS	(+)			
	Internode: waxiness					
	absent or very weak				Q179	1
	weak				Q138	3
	medium				Q121, RB72-454	5
	strong				H56-752, Q117	7
14	QN	VS	(+)	(a)		
	Internode: length of bud groove					
	short				Q121	3
	medium				Q135, Q138	5
	long				H56-752, Q179, Q96	7
15	QN	VS	(+)	(a)		
	Internode: depth of bud groove					
	absent or very shallow				Q117, Q121, Q186	1
	shallow				Q138, Q170, RB72-454	3
	medium				Q179	5
	deep				Q174	7

	English		français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
16	QN	VS	(+)				
	Internode: depth of growth crack						
	absent or very shallow					RB72-454	1
	shallow					Q124	2
	medium					Q121	3
	deep					Q179	4
	very deep						5
17	QN	MS/VG	(+)	(a)			
	Node: width of root band						
	narrow					Q215	3
	medium					SRA6	5
	wide					Q202	7
18	PQ	VG	(+)				
	Node: shape of root band						
	tall						1
	constricted						2
	conoidal						3
	obconoidal						4
19	QN	VS		(a)			
	Node: width of wax ring						
	absent or very narrow					Q179	1
	narrow					Q180	3
	medium					Q113, Q96, RB72-454	5
	wide					Q115, Q138	7
	very wide					Q195	9
20 (*)	QL	VG	(+)				
	Node: presence of wing on bud						
	absent						1
	present						9

	English		français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
21	(*)	PQ	VG	(+)			
		Node: shape of bud					
		triangular-pointed				RB72-454	1
		oval				Q138	2
		obovate				Q202	3
		pentagonal				Q182	4
		rhomboid				Q217	5
		round				Q124, Q179	6
		ovate				Q115, Q170, Q186	7
		rectangular				Q215	8
22		QN	MS/VG	(+)	(a)		
		Node: length of bud					
		very short					1
		short					3
		medium					5
		long					7
		very long					9
23		QN	MS	(+)	(a)		
		Node: width of bud					
		very narrow				Q186	1
		narrow				Q138	3
		medium				Q178	5
		wide				Q121, Q124	7
		very wide				H56-752, Q136	9
24	(*)	QN	VS	(+)			
		Node: bud prominence					
		very weak				Q152	1
		weak				RB72-454	3
		medium				H56-752, Q121	5
		strong				Q136	7

	English		français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
25	QN	VS	(a)				
	Node: position of bud tip in relation to growth ring						
	clearly below					Q171, WSRA18	1
	intermediate					Q179, RB72-454	2
	clearly above					Q172, SRA9	3
26	QN	VS	(+)	(a)			
	Node: bud cushion						
	absent or very narrow					Q121, Q186	1
	narrow					Q96	3
	medium					Q181, RB72-454	5
	wide					Q170	7
27	QN	VS	(+)				
	Node: width of bud wing						
	narrow					RB72-454	1
	medium					Q121	3
	wide					BN81-1394	5
28	PQ	VG	(+)				
	Node: color of root band where <u>not exposed</u> to sun						
	white and green						1
	yellow and green						2
	yellow and purple						3
	green						4
	purple						6
29	PQ	VG	(+)				
	Node: color of growth ring where <u>not exposed</u> to sun						
	white and green						1
	yellow and green						2
	yellow and purple						3
	green						4
	green and purple						5
	purple						6

	English		français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
30	QN	MS	(b)				
	Leaf sheath: length						
	short					Q117	3
	medium					Q136, Q170	5
	long					Q121, Q124	7
31	QN	VG	(b), (c)				
	Leaf sheath: number of hairs						
	absent or very few					Q186, RB72-454	1
	few					Q170	3
	medium					Q117, Q179	5
	many					Q124	7
	very many					Q169	9
32	QN	VG	(b), (c)				
	Leaf sheath: length of hairs						
	short					Q186	3
	medium					Q117, Q138, Q179	5
	long					Q121	7
33	PQ	VG	(b), (c)				
	Leaf sheath: distribution of hairs						
	only lateral					Q138, Q170	1
	lateral and dorsal					SRA5	2
	only dorsal					SRA19	3
34	PQ	VS	(+)	(b)			
	Leaf sheath: shape of ligule						
	strap-shaped					Argos	1
	deltoid					H56-752, Q170	2
	crescent-shaped					Q121, Q179, Q96	3
	bow-shaped						4
	asymmetrical, steeply sloping					Vertex 1 Vertex 7	5
	asymmetrical, horizontal					IACSP942094, RB72-454	6

	English		français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
35	QN	VS	(b)				
	Leaf sheath: width of ligule						
	narrow					WSRA17	1
	medium					Q115, Q179, Q186	2
	wide					H56-752, Q170	3
36	QN	VS	(b)				
	Leaf sheath: length of ligule hairs (c)						
	short					Q152, Q170, Q96	3
	medium					Q179, RB72-454	5
	long					BN81-1394, Q124	7
37	QN	VS	(b)				
	Leaf sheath: density of ligule hairs (c)						
	absent or very sparse					SRA6	1
	sparse					SRA25	3
	medium					Q152	5
	dense					Q121, RB72-454	7
	very dense					Q179	9
38	QL	VS	(b)				
	Leaf sheath: underlapping auricle						
	absent						1
	present					Q186	9
39 (*)	PQ	VS	(+)	(b)			
	Leaf sheath: shape of underlapping auricle						
	deltoid					Q186	1
	dentoid					SRA1, SRA2	2
	unciform						3
	calcarifom					Q196	4
	lanceolate					H56-752, RB72-454	5
	falcate					SRA16	6

	English		français	deutsch	español	Example Varieties Exemples Bei ejemplo	Note/
40	QN	VS	(b)				
	Leaf sheath: size of underlapping auricle						
		small				Q96	3
		medium				Q201	5
		large				Q135	7
41	QL	VS	(b)				
	Leaf sheath: overlapping auricle						
		absent					1
		present				SRA24	9
42 (*)	PQ	VS	(+)	(b)			
	Leaf sheath: shape of overlapping auricle						
		deltoid				Q117, RB72-454	1
		dentoid					2
		unciform					3
		calcariform					4
		lanceolate				Q138	5
		falcate					6
43	QN	MS/VS	(b)				
	Leaf sheath: size of overlapping auricle						
		small				SRA20, SRA25	3
		medium				Q251, SRA11	5
		large				Q198, Q215	7
44	QN	MS	(b)				
	Leaf blade: length						
		short				Q124	3
		medium				Q136	5
		long				Q170	7
45 (*)	QN	MS	(b)				
	Leaf blade: width at the longitudinal mid-point						
		narrow				Q113, Q186	3
		medium				Q121, Q124	5
		broad				Q138, Q179	7


	English		français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
46	QN	MS	(b)				
	Leaf: midrib width at the longitudinal mid-point						
	very narrow					Q203	1
	narrow					Q121	3
	medium					Q124, Q170	5
	wide					Q202, SRA5	7
	very wide					Q138	9
47	QN	MS	(b)				
	Leaf: ratio leaf blade width/midrib width						
	low					SRA5, SRA6	3
	medium					H56-752, Q124	5
	high					Q215, SRA11	7
48	QN	MS/VG	(d)				
	Cane top: length						
	short						3
	medium						5
	long						7
49	QL	VG	(d)				
	Cane top: shape of cross-section						
	circular						1
	ovate						2
50	QN	VG	(d)				
	Cane top: waxiness						
	absent or very weak						1
	weak						3
	medium						5
	strong						7

8. Explanations on the Table of Characteristics

8.1 *Explanations covering several characteristics*


Characteristics containing the following key in the Table of Characteristics should be examined as indicated below:

- (a) Observations on the internode and node to be made on the longest internode of a representative culm. Different parts of internode and node are illustrated:



- (b) Observations on the leaf sheath and leaf blade to be made on the top visible dewlap (TVD) leaf.
- (c) Leaf sheath hairs to be observed on hair groups 57 and 60. Distribution of hairs is dorsal when only hair group 57 is present. Distribution of hairs is lateral and dorsal when both hair groups 57 and 60 are present.


Leaf sheath ligule hairs to be observed on hair group 61.



- (d) The cane top is the region between the youngest exposed visible dewlap and the insertion of the fourth youngest fully extended leaf (leaf + 4) in the culm.

8.2 Explanations for individual characteristics

Ad. 1: Plant: stool growth habit




To be observed 2-3 months after transplantation.

Ad. 2: Plant: adherence of leaf sheath

To be observed on the lower half of the stool on the senescing leaves.

Ad. 4: Culm: height

To be observed from the base of the culm at soil level to the base of the Top Visible Dewlap (TVD) leaf.



Ad. 5: Internode: length on the bud side

To be observed on the bud side of the longest internode of the culm.


Ad. 6: Internode: diameter

Observations should be made at central part of the internode on the axis going through the bud


Ad. 7: Internode: shape



cylindrical




tumescent




bobbin-shaped



conoidal



obconoidal



concave-convex


Ad. 9: Internode: color where exposed to sun

After three days of exposure to the sun on a culm on which the wax has been removed

Ad. 10: Internode: color where not exposed to sun

On a culm protected from the sun, on which the wax has been removed.

Ad. 12: Internode: expression of zigzag alignment



Ad. 13: Internode: waxiness

Observation should be made on the longest internode

Ad. 14: Internode: length of bud groove

Observations should be made on the longest internode.

Ad. 15: Internode: depth of bud groove

Observations should be made on the longest internode.


Ad. 16: Internode: depth of growth crack

Observation should be made on the longest internode

Ad. 17: Node: width of root band

Observation should be made on the longest internode

Ad. 18: Node: shape of root band




Ad. 20: Node: presence of wing on bud

Observation should be made on the longest internode

Ad. 21: Node: shape of bud

To be observed excluding the bud wings



Ad. 22: Node: length of bud

To be observed excluding the bud wings.

Ad. 23: Node: width of bud

To be observed excluding the bud wings.

Ad. 24: Node: bud prominence

To be observed on second senescent leaf from the top.

Ad. 26: Node: bud cushion

To be observed in the space between base of bud and leaf scar.

Ad. 27: Node: width of bud wing

Observation should be made on the longest internode

Ad. 28: Node: color of root band where not exposed to sun

Observation should be made on the longest internode

Ad. 29: Node: color of growth ring where not exposed to sun

Observation should be made on the longest internode

Ad. 34: Leaf sheath: shape of ligule



1. strap-shaped



2. deltoid




3. crescent-shaped



4. bow-shaped

Ad. 39: Leaf sheath: shape of underlapping auricle



1
deltoid



2
dentoid



3
unciform



4
calcariform



5
lanceolate



6
falcate

Ad. 42: Leaf sheath: shape of overlapping auricle

Same as Leaf sheath: shape of underlapping auricle (Characteristics 39)

9. Literature

- Artschwager, E., 1940: Journal of Agricultural Research, v. 60, n. 8, pp. 503-508.
- Gallacher, D.J., 1994: Development of a minimum descriptor set for individuals of *Saccharum* spp. Hybrid germplasm. Thesis submitted for Ph.D., Department of Botany and Tropical Agriculture, James Cook University of North Queensland, AU.
- Gallacher, D.J. and Berding, N. 1997: Purpose selection and application of descriptors for sugarcane germplasm. *Aust. J. Agric. Res* 48: 759-67.
- Gallacher, D.J., 1997: Evaluation of sugarcane morphological descriptors using variance components analysis. *Aust. J. Agric. Res* 48: 769-73.
- Gallacher, D.J., 1997: Optimised descriptors recommended for Australian sugarcane germplasm (*Saccharum* spp. hybrid) *Aust. J. Agric. Res* 48: 775-79.

10. Technical Questionnaire

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
		Application date: (not to be filled in by the applicant)
TECHNICAL QUESTIONNAIRE to be completed in connection with an application for plant breeders' rights		
1. Subject of the Technical Questionnaire		
1.1	Botanical name	<input type="text" value="Saccharum L."/>
1.2	Common name	<input type="text" value="Sugarcane"/>
2. Applicant		
	Name	<input type="text"/>
	Address	<input type="text"/>
	Telephone No.	<input type="text"/>
	Fax No.	<input type="text"/>
	E-mail address	<input type="text"/>
	Breeder (if different from applicant)	<input type="text"/>
3. Proposed denomination and breeder's reference		
	Proposed denomination (if available)	<input type="text"/>
	Breeder's reference	<input type="text"/>

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

#4. Information on the breeding scheme and propagation of the variety

4.1 Breeding scheme

Variety resulting from:

4.1.1 Crossing

(a) controlled cross

(please state parent variety)

(.....) x (.....)

female parent male parent

(b) partially known cross

(please state known parent variety(ies))

(.....) x (.....)

female parent male parent

(c) unknown cross

4.1.2 Mutation

(please state parent variety)

4.1.3 Discovery and development

(please state where and when discovered and how developed)

4.1.4 Other

(Please provide details)

Authorities may allow certain of this information to be provided in a confidential section of the Technical Questionnaire.

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

4.2 Method of propagating the variety

4.2.1 Vegetative propagation

4.2.2 Other []
(Please provide details)

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

5. Characteristics of the variety to be indicated (the number in brackets refers to the corresponding characteristic in Test Guidelines; please mark the note which best corresponds).

Characteristics	Example Varieties	Note
5.1 Plant: adherence of leaf sheath (2)		
weak	H56-752, Q96	3 []
medium	Q124, Q186	5 []
strong	NC0 310, Q120, Q201	7 []
5.2 Internode: shape (7)		
cylindrical	Q169, RB72-454	1 []
tumescient	Q205	2 []
bobbin-shaped	H56-752	3 []
conoidal	Q177, Q178	4 []
obconoidal	H60-3802	5 []
concave-convex	Q115	6 []
5.3 Internode: color where <u>exposed</u> to sun (9)		
yellow	Q230	1 []
yellow green	SRA24, SRA25	2 []
grey yellow	SRA10	3 []
grey orange	Q165	4 []
grey red		5 []
grey purple	RB72-454	6 []
purple		7 []
5.4 Internode: color where <u>not exposed</u> to sun (10)		
yellow		1 []
yellow green	SRA24, SRA25	2 []
grey yellow	QS01-1078	3 []
grey orange	Q220	4 []
grey red		5 []
grey purple	SRA9	6 []
purple		7 []

Characteristics	Example Varieties	Note
5.5 Internode: expression of zigzag alignment (12)		
absent or very weak	Q124	1 []
weak	Q135, Q152	3 []
moderate	Q117	5 []
strong	H56-752	7 []
5.6 Node: shape of bud (21)		
triangular-pointed	RB72-454	1 []
oval	Q138	2 []
obovate	Q202	3 []
pentagonal	Q182	4 []
rhomboid	Q217	5 []
round	Q124, Q179	6 []
ovate	Q115, Q170, Q186	7 []
rectangular	Q215	8 []
5.7 Leaf blade: width at the longitudinal mid-point (45)		
narrow	Q113, Q186	3 []
medium	Q121, Q124	5 []
broad	Q138, Q179	7 []

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

6. Similar varieties and differences from these varieties

Please use the following table and box for comments to provide information on how your candidate variety differs from the variety (or varieties) which, to the best of your knowledge, is (or are) most similar. This information may help the examination authority to conduct its examination of distinctness in a more efficient way.

Denomination(s) of variety(ies) similar to your candidate variety	Characteristic(s) in which your candidate variety differs from the similar variety(ies)	Describe the expression of the characteristic(s) for the similar variety(ies)	Describe the expression of the characteristic(s) for your candidate variety
<i>Example</i>			
<p>Comments:</p>			

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

#7.	Additional information which may help in the examination of the variety		
7.1	In addition to the information provided in sections 5 and 6, are there any additional characteristics which may help to distinguish the variety?		
	Yes	<input type="checkbox"/>	No <input type="checkbox"/>
	(If yes, please provide details)		
7.2	Are there any special conditions for growing the variety or conducting the examination?		
	Yes	<input type="checkbox"/>	No <input type="checkbox"/>
	(If yes, please provide details)		
7.3	Other information		

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

8. Authorization for release

(a) Does the variety require prior authorization for release under legislation concerning the protection of the environment, human and animal health?

Yes [] No []

(b) Has such authorization been obtained?

Yes [] No []

If the answer to (b) is yes, please attach a copy of the authorization.

9. Information on plant material to be examined or submitted for examination

9.1 The expression of a characteristic or several characteristics of a variety may be affected by factors, such as pests and disease, chemical treatment (e.g. growth retardants or pesticides), effects of tissue culture, different rootstocks, scions taken from different growth phases of a tree, etc.

9.2 The plant material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If the plant material has undergone such treatment, full details of the treatment must be given. In this respect, please indicate below, to the best of your knowledge, if the plant material to be examined has been subjected to:

(a) Microorganisms (e.g. virus, bacteria, phytoplasma)	Yes []	No []
(b) Chemical treatment (e.g. growth retardant, pesticide)	Yes []	No []
(c) Tissue culture	Yes []	No []
(d) Other factors	Yes []	No []

Please provide details for where you have indicated "yes".

.....

10. I hereby declare that, to the best of my knowledge, the information provided in this form is correct:

Applicant's name

Signature Date

TG/186/2(proj.2) - Annex

Sugarcane, 2021-05-06

36

[End of document]