Technical Working Party on Automation and Computer Programs TWC/35/8

Thirty-Fifth Session Buenos Aires, Argentina, November 14 to 17, 2017 Original: English Date: October 23, 2017

ASSESSING UNIFORMITY BY OFFTYPES ON THE BASIS OF MORE THAN ONE GROWING CYCLE: EXAMPLES FROM NL

Document prepared by an expert from the Netherlands

Disclaimer: this document does not represent UPOV policies or guidance

The Annex to this document contains a copy of a presentation "Assessing Uniformity by Offtypes on the Basis of More than One Growing Cycle: examples from NL" to be made at its thirty-fifth session of the Technical Working Party on Automation and Computer Programs (TWC).

Lysbeth Hof, Researcher / Specialist Agricultural Crops, Variety Research Department, Naktuinbouw, Netherlands

[Annex follows]

TWC/35/8

ANNEX





TWC/35/8 Annex, page 2

Overview approaches
Summary of the three approaches:Approach 1: Third growing cycle in case of inconsistent results
 Approach 2: Combining the results of two growing cycles in the case of inconsistent results
Approach 3: Combining the results of two growing cycles
 A variety may be rejected after a single growing cycle if the number of offtypes exceeds the number of allowed offtypes for the combined sample (over two cycles)

7	Case 1: Tomato Variety A					
	Crop: Tomato Variety A Population standard 1%, Acceptance Prob. ≥ 95% Sample size per growing cycle = 20 Maximum number of offtypes per growing cycle = 1 Maximum number of offtypes growing cycle 1 and 2 combined (n=40) = 2					
	Number of offtypes per growing cycle			Decision		
	First	Second	Third	Approach 1	Approach 2	Approach 3
	3	0	0	third cycle: => uniform	non-uniform	non-uniform*
4			_	of document TWI ave been rejected		

TWC/35/8 Annex, page 3

Case 2: Tomato Variety B

Crop: Tomato Variety B Population standard 1%, Acceptance Prob. \geq 95% Sample size per growing cycle = 20 Maximum number of offtypes per growing cycle = 1 Maximum number of offtypes growing cycle 1 and 2 combined (n=40) = 2

Number of offtypes per growing cycle					Decision	
	First	Second	Third	Approach 1	Approach 2	Approach 3
	3	1	4	third cycle: => non-uniform	non-uniform	non-uniform

7	Comparing Variety A and Variety B
	 Both in Variety A and in Variety B the number of offtypes in the first growing cycle was 3 (non-uniform) In Variety A, approach 1 eventually lead to decision 'uniform' after 3 growing cycles, and in Variety B the final decision after 3 growing cycles was non-uniform In Variety A, the current wording of the document (TWP/1/17) could/would have resulted in a rejection after the first growing cycle. In retrospect a 'wrong' decision? Early decisions, based on small deviations from the allowable number of offtypes, can be premature and are risky
6	

5

TWC/35/8 Annex, page 4



Population standard 1%, Acceptance Prob. ≥ 95% Sample size per growing cycle = 50 Maximum number of offtypes per growing cycle = 2 Maximum number of offtypes growing cycle 1 and 2 combined (n=100) = 3

Number of offtypes per growing cycle				Decision	
First	Second	Third	Approach 1	Approach 2	Approach 3
2	2	-	uniform	uniform	non-uniform

This example illustrates the pitfall of approach 3. It considers the variety non-uniform, while it is considered uniform in both separate cycles!

7	Conclusions
	 The number of allowed off-types in a sample is sometimes small. This increases the risk of a 'wrong' decision: one offtype more or less could lead to a different decision In approach 3, the allowed number of offtypes of the combined cycles can be smaller than the sum of allowed offtypes of both cycles (see example previous slide). This may lead to a 'questionable' decision Decisions on non-uniformity should only be made when cases are <u>clear</u>. In case of doubt: the benefit should be for the applicant. Naktuinbouw has decided to use approach 1, as with this approach the chance of a 'wrong' decision seems smallest
8	

[End of Annex and of document]