

UPOV

TG/PAPAYA(proj.4)

ORIGINAL: English

DATE: 2008-04-22

INTERNATIONAL UNION FOR THE PROTECTION OF NEW VARIETIES OF PLANTS

GENEVA

DRAFT

PAPAYA

UPOV Code: CARIC_PAP

Carica papaya L.

GUIDELINES

FOR THE CONDUCT OF TESTS

FOR DISTINCTNESS, UNIFORMITY AND STABILITY

prepared by an expert from Mexico

to be considered by the

Technical Working Party for Fruit Crops

at its thirty-ninth session, to be held in Lisbon, Portugal, from June 2 to 6, 2008

Alternative Names:*

| <i>Botanical name</i> | <i>English</i> | <i>French</i> | <i>German</i> | <i>Spanish</i> |
|-------------------------|----------------|------------------------|---------------------|-----------------|
| <i>Carica papaya</i> L. | Papaya, Papaw | Arbre à melon, Papayer | Melonenbaum, Papaya | Papayo, Lechosa |

The purpose of these guidelines (“Test Guidelines”) is to elaborate the principles contained in the General Introduction (document TG/1/3), and its associated TGP documents, into detailed practical guidance for the harmonized examination of distinctness, uniformity and stability (DUS) and, in particular, to identify appropriate characteristics for the examination of DUS and production of harmonized variety descriptions.

ASSOCIATED DOCUMENTS

These Test Guidelines should be read in conjunction with the General Introduction and its associated TGP documents.

* These names were correct at the time of the introduction of these Test Guidelines but may be revised or updated. [Readers are advised to consult the UPOV Code, which can be found on the UPOV Website (www.upov.int), for the latest information.]

| <u>TABLE OF CONTENTS</u> | <u>PAGE</u> |
|--|-------------|
| 1. SUBJECT OF THESE TEST GUIDELINES..... | 4 |
| 2. MATERIAL REQUIRED | 4 |
| 3. METHOD OF EXAMINATION..... | 4 |
| 3.1 Number of Growing Cycles | 4 |
| 3.2 Testing Place | 4 |
| 3.3 Conditions for Conducting the Examination..... | 4 |
| 3.4 Test Design | 5 |
| 3.5 Number of Plants / Parts of Plants to be Examined..... | 5 |
| 3.6 Additional Tests | 5 |
| 4. ASSESSMENT OF DISTINCTNESS, UNIFORMITY AND STABILITY | 5 |
| 4.1 Distinctness | 5 |
| 4.2 Uniformity..... | 6 |
| 4.3 Stability | 6 |
| 5. GROUPING OF VARIETIES AND ORGANIZATION OF THE GROWING TRIAL..... | 6 |
| 6. INTRODUCTION TO THE TABLE OF CHARACTERISTICS | 7 |
| 6.1 Categories of Characteristics..... | 7 |
| 6.2 States of Expression and Corresponding Notes..... | 7 |
| 6.3 Types of Expression..... | 7 |
| 6.4 Example Varieties | 7 |
| 6.5 Legend..... | 7 |
| 7. TABLE OF CHARACTERISTICS/TABLEAU DES CARACTÈRES/MERKMALSTABELLE/TABLA DE CARACTERES..... | 8 |
| 8. EXPLANATIONS ON THE TABLE OF CHARACTERISTICS | 17 |
| 8.1 Explanations covering several characteristics | 17 |
| 8.2 Explanations for individual characteristics | 17 |
| 9. LITERATURE | 20 |
| 10. TECHNICAL QUESTIONNAIRE | 21 |

1. Subject of these Test Guidelines

These Test Guidelines apply to all varieties of *Carica papaya* L. of the family *Caricaceae*.

2. Material Required

2.1 The competent authorities decide on the quantity and quality of the plant material required for testing the variety and when and where it is to be delivered. Applicants submitting material from a State other than that in which the testing takes place must ensure that all customs formalities and phytosanitary requirements are complied with.

2.2 The material is to be supplied in the form of seed or plants.

2.3 The minimum quantity of plant material, to be supplied by the applicant, should be:

200 seeds in the case of seed-propagated varieties,
or 10 plants in the case of vegetatively propagated varieties.

In the case of seed, the seed should meet the minimum requirements for germination, species and analytical purity, health and moisture content, specified by the competent authority.

2.4 The plant material supplied should be visibly healthy, not lacking in vigor, nor affected by any important pest or disease.

2.5 The plant material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If it has been treated, full details of the treatment must be given.

3. Method of Examination

3.1 *Number of Growing Cycles*

3.1.1 The minimum duration of tests should normally be two growing cycles.

3.1.2 The growing cycle is considered to be the duration of a single growing season, beginning with vegetative growth, followed by flowering and fruit harvest.

3.2 *Testing Place*

Tests are normally conducted at one place. In the case of tests conducted at more than one place, guidance is provided in TGP/9 "Examining Distinctness".

3.3 *Conditions for Conducting the Examination*

The tests should be carried out under conditions ensuring satisfactory growth for the expression of the relevant characteristics of the variety and for the conduct of the examination. In particular, it is essential that the trees produce a satisfactory crop of fruit in each of the two growing cycles.

3.4 *Test Design*

3.4.1 Each test should be designed to result in a total of at least 25 hermaphrodite plants in the case of seed-propagated plants or, in the case of vegetatively propagated varieties, in a total of at least 10 hermaphrodite plants.

3.4.2 The design of the tests should be such that plants or parts of plants may be removed for measurement or counting without prejudice to the observations which must be made up to the end of the growing cycle.

3.5 *Number of Plants / Parts of Plants to be Examined*

Unless otherwise indicated, all observations should be made on 25 hermaphrodite plants in the case of seed-propagated varieties or, in the case of vegetatively propagated varieties, on 10 hermaphrodite plants.

3.6 *Additional Tests*

Additional tests, for examining relevant characteristics, may be established.

4. Assessment of Distinctness, Uniformity and Stability

4.1 *Distinctness*

4.1.1 *General Recommendations*

It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding distinctness. However, the following points are provided for elaboration or emphasis in these Test Guidelines.

4.1.2 *Consistent Differences*

The differences observed between varieties may be so clear that more than one growing cycle is not necessary. In addition, in some circumstances, the influence of the environment is not such that more than a single growing cycle is required to provide assurance that the differences observed between varieties are sufficiently consistent. One means of ensuring that a difference in a characteristic, observed in a growing trial, is sufficiently consistent is to examine the characteristic in at least two independent growing cycles.

4.1.3 *Clear Differences*

Determining whether a difference between two varieties is clear depends on many factors, and should consider, in particular, the type of expression of the characteristic being examined, i.e. whether it is expressed in a qualitative, quantitative, or pseudo-qualitative manner. Therefore, it is important that users of these Test Guidelines are familiar with the

recommendations contained in the General Introduction prior to making decisions regarding distinctness.

4.2 *Uniformity*

4.2.1 It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding uniformity. However, the following points are provided for elaboration or emphasis in these Test Guidelines.

4.2.2 For the assessment of uniformity for seed-propagated varieties, a population standard of 1% and an acceptance probability of at least 95% should be applied. In the case of a sample size of 25 hermaphrodite plants, one off-type is allowed.

4.2.3. For the assessment of uniformity for vegetatively propagated varieties, a population standard of 1% and an acceptance probability of 95% should be applied. In the case of a sample size of 10 hermaphrodite plants, one off-type is allowed.

4.3 *Stability*

4.3.1 In practice, it is not usual to perform tests of stability that produce results as certain as those of the testing of distinctness and uniformity. However, experience has demonstrated that, for many types of variety, when a variety has been shown to be uniform, it can also be considered to be stable.

4.3.2 Where appropriate, or in cases of doubt, stability may be tested, either by growing a further generation, or by testing a new plant stock to ensure that it exhibits the same characteristics as those shown by the previous material supplied.

5. Grouping of Varieties and Organization of the Growing Trial

5.1 The selection of varieties of common knowledge to be grown in the trial with the candidate varieties and the way in which these varieties are divided into groups to facilitate the assessment of distinctness are aided by the use of grouping characteristics.

5.2 Grouping characteristics are those in which the documented states of expression, even where produced at different locations, can be used, either individually or in combination with other such characteristics: (a) to select varieties of common knowledge that can be excluded from the growing trial used for examination of distinctness; and (b) to organize the growing trial so that similar varieties are grouped together.

5.3 The following have been agreed as useful grouping characteristics:

- (a) Plant: height to first flower (characteristic 2)
- (b) Leaf blade: ratio length/width (characteristic 9)
- (c) Fruit: ratio length/maximum diameter (characteristic 24).
- (d) Fruit: shape (characteristic 25).

5.4 Guidance for the use of grouping characteristics, in the process of examining distinctness, is provided through the General Introduction.

6. Introduction to the Table of Characteristics

6.1 *Categories of Characteristics*

6.1.1 Standard Test Guidelines Characteristics

Standard Test Guidelines characteristics are those which are approved by UPOV for examination of DUS and from which members of the Union can select those suitable for their particular circumstances.

6.1.2 Asterisked Characteristics

Asterisked characteristics (denoted by *) are those included in the Test Guidelines which are important for the international harmonization of variety descriptions and should always be examined for DUS and included in the variety description by all members of the Union, except when the state of expression of a preceding characteristic or regional environmental conditions render this inappropriate.

6.2 *States of Expression and Corresponding Notes*

States of expression are given for each characteristic to define the characteristic and to harmonize descriptions. Each state of expression is allocated a corresponding numerical note for ease of recording of data and for the production and exchange of the description.

6.3 *Types of Expression*

An explanation of the types of expression of characteristics (qualitative, quantitative and pseudo-qualitative) is provided in the General Introduction.

6.4 *Example Varieties*

Where appropriate, example varieties are provided to clarify the states of expression of each characteristic.

6.5 *Legend*

(*) Asterisked characteristic – see Chapter 6.1.2

QL Qualitative characteristic – see Chapter 6.3

QN Quantitative characteristic – see Chapter 6.3

PQ Pseudo-qualitative characteristic – see Chapter 6.3

(a)-(g) See Explanations on the Table of Characteristics in Chapter 8.1

(+) See Explanations on the Table of Characteristics in Chapter 8.2

7. Table of Characteristics/Tableau des caractères/Merkmalstabelle/Tabla de caracteres

| | English | français | deutsch | español | Example Varieties Exemples Beispielssorten Variedades ejemplo | Note/ Nota |
|-----------|--|----------|---------|---------|--|---------------|
| 1. | Young plant: color of stem | | | | | |
| (+) | | | | | | |
| PQ | green | | | | Ishigaki sango | 1 |
| | brown | | | | | 2 |
| | green and purple | | | | Sunrise | 3 |
| | purple | | | | | 4 |
| 2. | Plant: height to first flower | | | | | |
| (*) | | | | | | |
| QN | (a) short | | | | Ishigaki sango | 3 |
| | medium | | | | Sunrise | 5 |
| | tall | | | | | 7 |
| 3. | Plant: branching | | | | | |
| (+) | | | | | | |
| QL | absent | | | | Ishigaki sango, Maradol, Sunrise | 1 |
| | present | | | | | 9 |
| 4. | Stem: maximum diameter | | | | | |
| QN | (a) small | | | | | 3 |
| | medium | | | | Ishigaki sango, Sunrise | 5 |
| | large | | | | | 7 |
| 5. | Stem: number of nodes from ground to first flower | | | | | |
| QN | (a) few | | | | Ishigaki sango | 3 |
| | medium | | | | Sunrise | 5 |
| | many | | | | | 7 |

| | English | français | deutsch | español | Example Varieties Exemples Beispielssorten Variedades ejemplo | Note/ Nota |
|--------------------|--|----------|---------|---------|--|---------------|
| 6. | Stem: length of internode at the middle between ground and first flower | | | | | |
| QN | (a) short | | | | Ishigaki sango | 3 |
| | medium | | | | Sunrise | 5 |
| | long | | | | | 7 |
| 7. | Leaf blade: length | | | | | |
| QN | (b) short | | | | | 3 |
| | medium | | | | Ishigaki sango, Sunrise | 5 |
| | long | | | | | 7 |
| 8. | Leaf blade: width | | | | | |
| QN | (b) narrow | | | | | 3 |
| | medium | | | | | 5 |
| | broad | | | | | 7 |
| 9. (*) | Leaf blade: ratio length/ width | | | | | |
| QN | (b) small | | | | | 3 |
| | medium | | | | Ishigaki sango, Sunrise | 5 |
| | large | | | | | 7 |
| 10. (+) | Leaf blade: presence of tertiary lobes | | | | | |
| QL | (b) absent | | | | | 1 |
| | present | | | | Ishigaki sango, Sunrise | 9 |

| | English | français | deutsch | español | Example Varieties Exemples Beispielssorten Variedades ejemplo | Note/ Nota |
|------------|--|----------|---------|---------|--|---------------|
| 11. | Leaf blade: waxiness | | | | | |
| QL | (b) absent | | | | Ishigaki sango, Sunrise | 1 |
| | present | | | | | 9 |
| 12. | Leaf blade: pubescence on lower side | | | | | |
| QL | (b) absent (c) | | | | Ishigaki sango, Sunrise | 1 |
| | present | | | | | 9 |
| 13. | Petiole: length | | | | | |
| QN | (b) short | | | | | 3 |
| | medium | | | | Ishigaki sango, Sunrise | 5 |
| | long | | | | | 7 |
| 14. | Petiole: anthocyanin coloration | | | | | |
| QL | (b) absent | | | | Ishigaki sango | 1 |
| | present | | | | Sunrise | 9 |
| 15. | Petiole: intensity of anthocyanin coloration | | | | | |
| QN | (b) weak | | | | | 1 |
| | medium | | | | Sunrise | 2 |
| | strong | | | | | 3 |
| 16. | Inflorescence (excluding solitary flowers): number of flowers | | | | | |
| QN | (d) few | | | | Ishigaki sango | 3 |
| | medium | | | | | 5 |
| | many | | | | | 7 |

| | English | français | deutsch | español | Example Varieties Exemples Beispielssorten Variedades ejemplo | Note/ Nota |
|------------|--|----------|---------|---------|--|---------------|
| 17. | Inflorescence: length of main axis | | | | | |
| QN | (d) short | | | | Ishigaki sango, Sunrise | 3 |
| | medium | | | | | 5 |
| | long | | | | | 7 |
| 18. | Inflorescence: color of axis | | | | | |
| PQ | (d) green | | | | Ishigaki sango, Sunrise | 1 |
| | light purple | | | | | 2 |
| | dark purple | | | | | 3 |
| 19. | Hermaphrodite flower: length of corolla | | | | | |
| QN | (e) short | | | | | 3 |
| | medium | | | | Sunrise | 5 |
| | long | | | | | 7 |
| 20. | Hermaphrodite flower: color of corolla | | | | | |
| PQ | (e) white | | | | | 1 |
| | cream | | | | | 2 |
| | yellow | | | | | 3 |
| | dark yellow to orange | | | | | 4 |
| | green | | | | | 5 |
| | dark green | | | | | 6 |
| | yellow green and purple | | | | | 7 |
| | purple | | | | | 8 |
| | dark purple | | | | | 9 |

| | English | français | deutsch | español | Example Varieties Exemples Beispielssorten Variedades ejemplo | Note/ Nota |
|------------------------------|---|----------|---------|---------|--|---------------|
| 21. | Peduncle: length | | | | | |
| QN | (f) short | | | | Ishigaki sango, Sunrise | 3 |
| | medium | | | | | 5 |
| | long | | | | | 7 |
| 22. | Fruit: length | | | | | |
| QN | (f) short | | | | Sunrise | 3 |
| | medium | | | | Ishigaki sango | 5 |
| | long | | | | | 7 |
| 23. | Fruit: maximum diameter | | | | | |
| QN | (f) small | | | | Sunrise | 3 |
| | medium | | | | Ishigaki sango | 5 |
| | large | | | | | 7 |
| 24. (*) | Fruit: ratio length/maximum diameter | | | | | |
| QN | (f) small | | | | | 3 |
| | medium | | | | Ishigaki sango | 5 |
| | large | | | | Sunrise | 7 |
| 25. (*) (+) | Fruit: shape | | | | | |
| PQ | (f) ovoid | | | | | 1 |
| | ellipsoid | | | | Ishigaki sango | 2 |
| | obovoid | | | | Red Lady | 3 |
| | piriform | | | | Kapoho, Rainbow | 4 |
| | oblong | | | | Amarela | 5 |

| | English | français | deutsch | español | Example Varieties Exemples Beispielssorten Variedades ejemplo | Note/ Nota |
|------------|---|----------|---------|---------|--|---------------|
| 26. | Fruit: shape of stalk end | | | | | |
| (+) | | | | | | |
| PQ | (f) depressed | | | | Ishigaki sango | 1 |
| | truncate | | | | | 2 |
| | rounded | | | | | 3 |
| | pointed | | | | | 4 |
| 27. | Fruit: shape at distal end | | | | | |
| PQ | (f) rounded | | | | | 1 |
| | weakly pointed | | | | Ishigaki sango, Sunrise | 2 |
| | strongly pointed | | | | | 3 |
| 28. | Ripe fruit: main color | | | | | |
| PQ | (g) green | | | | | 1 |
| | yellow green | | | | | 2 |
| | yellow | | | | Amarela, Kapoho | 3 |
| | medium orange | | | | Ishigaki sango, Maradol, Mulata | 4 |
| | dark orange | | | | Mamey | 5 |
| 29. | Ripe fruit: ridges | | | | | |
| QL | (g) absent | | | | Sunrise | 1 |
| | present | | | | Ishigaki sango | 9 |
| 30. | Ripe fruit: prominence of ridges | | | | | |
| QN | (g) weakly expressed | | | | Ishigaki sango | 1 |
| | medium | | | | | 2 |
| | strongly expressed | | | | | 3 |

| | English | français | deutsch | español | Example Varieties Exemples Beispielssorten Variedades ejemplo | Note/ Nota |
|------------|--|----------|---------|---------|--|---------------|
| 31. | Ripe fruit: thickness of skin | | | | | |
| QN | (g) thin | | | | | 1 |
| | medium | | | | Sunrise | 2 |
| | thick | | | | | 3 |
| 32. | Ripe fruit: color of flesh | | | | | |
| PQ | (g) yellow | | | | Amarela, Cera, Kapoho | 1 |
| | orange | | | | | 2 |
| | red orange | | | | Ishigaki sango, Maradol | 3 |
| 33. | Ripe fruit: firmness of flesh | | | | | |
| QN | (g) soft | | | | Cera, Mamey | 3 |
| | medium | | | | Maradol | 5 |
| | firm | | | | Sunrise | 7 |
| 34. | Ripe fruit: sweetness | | | | | |
| QN | (g) low | | | | Cera | 3 |
| | medium | | | | Maradol | 5 |
| | high | | | | Ishigaki sango, Sunrise | 7 |
| 35. | Ripe fruit: aroma of flesh | | | | | |
| QN | (g) weak | | | | Maradol | 1 |
| | moderate | | | | Ishigaki sango, Sunrise | 2 |
| | strong | | | | Caera | 3 |
| 36. | Ripe fruit: placental tissue | | | | | |
| QN | (g) scarce | | | | Mamey | 3 |

| | English | français | deutsch | español | Example Varieties Exemples Beispielssorten Variedades ejemplo | Note/ Nota |
|----------------|---|----------|---------|---------|--|---------------|
| | medium | | | | Sunrise | 5 |
| | abundant | | | | Cera | 7 |
| 37. | Ripe fruit: central cavity: maximum width | | | | | |
| QN | (g) narrow | | | | Sunrise | 3 |
| | medium | | | | Ishigaki sango | 5 |
| | broad | | | | | 7 |
| 38. | Ripe fruit: central cavity predominant shape | | | | | |
| (+) | | | | | | |
| PQ | (g) circular | | | | | 1 |
| | angular | | | | | 2 |
| | star-shaped | | | | Ishigaki sango | 3 |
| | irregular | | | | | 4 |
| 39. (*) | Ripe fruit: seeds | | | | | |
| QL | (g) absent or very few | | | | Ishigaki sango | 1 |
| | few | | | | | 3 |
| | many | | | | Sunrise | 7 |
| | very many | | | | Cera | 9 |
| 40. | Seed: color (only normal seeds) | | | | | |
| PQ | (f) grey yellow | | | | | 1 |
| | grey | | | | | 2 |
| | brown | | | | | 3 |
| | dark brown | | | | Sunrise | 4 |
| | black | | | | Maradol | 5 |

| | English | français | deutsch | español | Example Varieties Exemples Beispielssorten Variedades ejemplo | Note/ Nota | |
|------------|-------------------------------------|----------|---------|---------|--|---------------|--|
| 41. | Seed: length | | | | | | |
| QN | (f) short | | | | | 3 | |
| | medium | | | | Sunrise | 5 | |
| | long | | | | Cera | 7 | |
| 42. | Seed: width | | | | | | |
| QN | (f) narrow | | | | | 3 | |
| | medium | | | | Sunrise | 5 | |
| | broad | | | | | 7 | |
| 43. | Seed: length/width ratio | | | | | | |
| QN | (f) small | | | | | 3 | |
| | medium | | | | Sunrise | 5 | |
| | large | | | | | 7 | |
| 44. | Seed: shape | | | | | | |
| PQ | (f) round | | | | Sunrise | 1 | |
| | ellipsoid | | | | | 2 | |
| | ovoid | | | | | 3 | |
| 45. | Seed: amount of mucilage | | | | | | |
| QN | (f) small | | | | | 1 | |
| | intermediate | | | | Sunrise | 2 | |
| | large | | | | Cera | 3 | |

8. Explanations on the Table of Characteristics

8.1 *Explanations covering several characteristics*

Characteristics containing the following key in the second column of the Table of Characteristics should be examined as indicated below:

- (a) Plant and stem: All observations on the tree and stem should be made at the beginning of fruit maturity.
- (b) Leaf blade and petiole: All observations on the leaf blade and petiole should be made on mature leaves. Leaves should be taken from the middle third of the current season's growth at the beginning of fruit maturity.
- (c) Pubescence: All observations on pubescence should be made with the aid of a magnifying glass.
- (d) Inflorescence: All observations on inflorescence should be made after the fourth one has appeared.
- (e) Flower: All observations on the flower should be made during the first flower opening, at the start of anther dehiscence.
- (f) Peduncle, fruit and seed: All observations on the peduncle, fruit and seed should be made on 5 typical fruits from hermaphrodite flowers, taken from the middle part of the fruiting region with a minimum sample of 10 fruits, at the time of maturity for harvest.
- (g) Ripe fruit: Observations on the ripe fruit should be made when the fruit is ready for eating.

8.2 *Explanations for individual characteristics*


Ad. 1. Young plant: color of stem

To be observed when the first bud appears.

Ad. 3. Plant branching


To be observed at the beginning of flowering.

Ad. 10: Leaf blade: presence of tertiary lobes




Ad. 25: Fruit: shape


To be observed from hermaphrodite flower.



Ad. 26: Fruit: shape of stalk end



Ad. 38: Fruit: central cavity predominant shape



1
circular



2
angular



3
star-shaped



4
irregular

9. Literature

IBPGR, 1988: Descriptors for Papaya. International Board for Plant Genetic Resources. Rome, Italy, 34 p.

Loyola, J. L. D., Pinto, R. M. de S., Lima, J. F. de, Ferreira, F. R. 2000: Catálogo de germoplasma de mamão (*Carica papaya* L.). Embrapa Mandioca e Fruticultura, Cruz das Almas, Bahia, Brasil, 40 p.

10. Technical Questionnaire

| | | |
|--|-----------------|---|
| TECHNICAL QUESTIONNAIRE | Page {x} of {y} | Reference Number: |
| | | Application date: (not to be filled in by the applicant) |
| TECHNICAL QUESTIONNAIRE to be completed in connection with an application for plant breeders' rights | | |
| 1. Subject of the Technical Questionnaire 1.1 Botanical name <input data-bbox="576 730 1374 786" type="text" value="Carica papaya L."/> 1.2 Common name <input data-bbox="576 819 1374 875" type="text" value="Papaya"/> | | |
| 2. Applicant Name <input data-bbox="576 1021 1374 1066" type="text"/> Address <input data-bbox="576 1099 1374 1245" type="text"/> Telephone No. <input data-bbox="576 1290 1374 1335" type="text"/> Fax No. <input data-bbox="576 1368 1374 1413" type="text"/> E-mail address <input data-bbox="576 1447 1374 1491" type="text"/> Breeder (if different from applicant) <input data-bbox="576 1559 1374 1603" type="text"/> | | |
| 3. Proposed denomination and breeder's reference Proposed denomination <input data-bbox="576 1749 1374 1794" type="text"/> (if available) Breeder's reference <input data-bbox="576 1861 1374 1906" type="text"/> | | |

| | | |
|-------------------------|-----------------|-------------------|
| TECHNICAL QUESTIONNAIRE | Page {x} of {y} | Reference Number: |
|-------------------------|-----------------|-------------------|

#4. Information on the breeding scheme and propagation of the variety

4.1 Breeding Scheme

Variety resulting from:

4.1.1 Crossing

- (a) controlled cross []
(please state parent varieties)
- (b) partially known cross []
(please state known parent variety(ies))
- (c) unknown cross []

4.1.2 Mutation []
(please state parent variety)

4.1.3 Discovery and development []
(please state where and when discovered and how developed)

4.1.4 Other []
(please provide details)

4.2 Method of propagating the variety

Authorities may allow certain of this information to be provided in a confidential section of the Technical Questionnaire.

| | | |
|-------------------------|-----------------|-------------------|
| TECHNICAL QUESTIONNAIRE | Page {x} of {y} | Reference Number: |
|-------------------------|-----------------|-------------------|

5. Characteristics of the variety to be indicated (the number in brackets refers to the corresponding characteristic in Test Guidelines; please mark the note which best corresponds).

| Characteristics | Example Varieties | Note |
|--|-------------------|------|
| 5.1 Plant: height to first flower (2) | | |
| short | Ishigaki sango | 3[] |
| medium | Sunrise | 5[] |
| tall | | 7[] |
| 5.2 Leaf blade: ratio length/width (9) | | |
| small | | 3[] |
| medium | | 5[] |
| large | | 7[] |
| 5.3 Fruit: ratio length/maximum diameter (24) | | |
| small | | 3[] |
| medium | | 5[] |
| large | | 7[] |
| 5.4 Fruit: shape (25) | | |
| ovoid | | 1[] |
| ellipsoid | Ishigaki sango | 2[] |
| obovoid | Red Lady | 3[] |
| piriform | Kapoho, Rainbow | 4[] |
| oblong | Amarela | 5[] |

| | | |
|-------------------------|-----------------|-------------------|
| TECHNICAL QUESTIONNAIRE | Page {x} of {y} | Reference Number: |
|-------------------------|-----------------|-------------------|

6. Similar varieties and differences from these varieties

Please use the following table and box for comments to provide information on how your candidate variety differs from the variety (or varieties) which, to the best of your knowledge, is (or are) most similar. This information may help the examination authority to conduct its examination of distinctness in a more efficient way.

| Denomination(s) of variety(ies) similar to your candidate variety | Characteristic(s) in which your candidate variety differs from the similar variety(ies) | Describe the expression of the characteristic(s) for the similar variety(ies) | Describe the expression of the characteristic(s) for your candidate variety |
|---|---|--|--|
| Example | <i>Petiole: anthocyanin coloration</i> | <i>e.g. note 1</i> | <i>note 9</i> |
| | | <i>e.g. absent</i> | <i>present</i> |

Comments:

| | | |
|-------------------------|-----------------|-------------------|
| TECHNICAL QUESTIONNAIRE | Page {x} of {y} | Reference Number: |
|-------------------------|-----------------|-------------------|

#7. Additional information which may help in the examination of the variety

7.1 In addition to the information provided in sections 5 and 6, are there any additional characteristics, which may help to distinguish the variety?

Yes [] No []

(If yes, please provide details)

7.2 Are there any special conditions for growing the variety or conducting the examination?

Yes [] No []

(If yes, please provide details)

7.3 Other information

A representative color photograph of the variety should accompany the Technical Questionnaire

8. Authorization for release

(a) Does the variety require prior authorization for release under legislation concerning the protection of the environment, human and animal health?

Yes [] No []

(b) Has such authorization been obtained?

Yes [] No []

If the answer to (b) is yes, please attach a copy of the authorization.

#

Authorities may allow certain of this information to be provided in a confidential section of the Technical Questionnaire.

| | | |
|-------------------------|-----------------|-------------------|
| TECHNICAL QUESTIONNAIRE | Page {x} of {y} | Reference Number: |
|-------------------------|-----------------|-------------------|

9. Information on plant material to be examined or submitted for examination.

9.1 The expression of a characteristic or several characteristics of a variety may be affected by factors, such as pests and disease, chemical treatment (e.g. growth retardants or pesticides), effects of tissue culture, different rootstocks, scions taken from different growth phases of a tree, etc.

9.2 The plant material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If the plant material has undergone such treatment, full details of the treatment must be given. In this respect, please indicate below, to the best of your knowledge, if the plant material to be examined has been subjected to:

- | | | |
|---|---------|--------|
| (a) Microorganisms (e.g. virus, bacteria, phytoplasma) | Yes [] | No [] |
| (b) Chemical treatment (e.g. growth retardant, pesticide) | Yes [] | No [] |
| (c) Tissue culture | Yes [] | No [] |
| (d) Other factors | Yes [] | No [] |

Please provide details for where you have indicated “yes”.

.....

10. I hereby declare that, to the best of my knowledge, the information provided in this form is correct:

Applicant's name

Signature

Date

[End of document]