

TWF/44/11

ORIGINAL: English **DATE:** April 18, 2013

INTERNATIONAL UNION FOR THE PROTECTION OF NEW VARIETIES OF PLANTS

Geneva

TECHNICAL WORKING PARTY FOR FRUIT CROPS

Forty-Fourth Session Napier, New Zealand, April 29 to May 3, 2013

REVISION OF DOCUMENT TGP/7: INDICATION OF GROWTH STAGE IN TEST GUIDELINES.

Document prepared by the Office of the Union

- 1. The purpose of this document is to consider guidance on indicating the growth stage at which to observe characteristics in the Test Guidelines for a future revision of document TGP/7: "Development of Test Guidelines".
- 2. The following abbreviations are used in this document:

TC: Technical Committee

TC-EDC: Enlarged Editorial Committee

TWA: Technical Working Party for Agricultural Crops

TWC: Technical Working Party on Automation and Computer Programs

TWF: Technical Working Party for Fruit Crops

TWO: Technical Working Party for Ornamental Plants and Forest Trees

TWPs: Technical Working Parties

TWV: Technical Working Party for Vegetables

3. The structure of this document is as follows:

BACKGROUNDEXISTING GUIDANCE IN DOCUMENT TGP/7	
TG TemplateGN 23	2
Chapter 3.3 and GN 9	3 3
PROPOSALS FOR INDICATING GROWTH STAGE AT WHICH TO OBSERVE CHARACTERISTICS IN	0
THE TEST GUIDELINES	4

ANNEX: EXAMPLES OF INDICATIONS OF GROWTH STAGE AT WHICH TO OBSERVE CHARACTERISTICS IN EXISTING TEST GUIDELINES


BACKGROUND

4. The Technical Committee (TC), at its forty-ninth session held in Geneva from March 18 to 20, 2013, agreed that clarification should be provided in a future revision of document TGP/7 with regard to the inclusion of growth stage keys in Chapter 8 of the Test Guidelines and requested the Office of the Union to prepare draft guidance for consideration by the TWPs at their sessions in 2013 (see document TC/49/41 "Report on the Conclusions", paragraph 83).

EXISTING GUIDANCE IN DOCUMENT TGP/7

TG Template

5. The TG Template and associated Guidance Note (GN24) provide the following guidance for indicating the growth stage for the observation of characteristics in the Test Guidelines:



GN 24 (TG Template: Chapter 7: column 2, header row 1) – Growth stage

In some Test Guidelines, the growth stage at which the examination of the characteristic should be done is provided here. In such cases, the stages of development denoted by each number are described in a section within Chapter 8, according to ASW 4(a).

ASW 4 (TG Template: Chapter 3.3) - Conditions for conducting the examination

Information for conducting the examination of particular characteristics

(a) Stage of development for the assessment

"The optimum stage of development for the assessment of each characteristic is indicated by a number in the second column of the Table of Characteristics. The stages of development denoted by each number are described in Chapter 8 [...]."

GN 23

6. Guidance Note GN 23 for drafters of Test Guidelines: "Explanations covering several characteristics", also states as follows:

GN 23 (TG Template: Chapter 7: column 2, state of expression row 1) – Explanations covering several characteristics

In cases where an explanation applies to several characteristics (e.g. part of the plant on which to observe particular characteristics, illustration of plant parts, etc.), particularly for characteristics that are not immediately consecutive in the Table of Characteristics, a note is placed in column 2 and the explanation provided in Chapter 8.1, according to ASW 11. In the case of indications of the stage of observation, those indications should be made according to GN 24 "Growth stage".

Chapter 3.3 and GN 9

7. The TG Template and Guidance Note (GN 9) provide further guidance as follows:

3.2 Testing Place

Tests are normally conducted at one place. In the case of tests conducted at more than one place, guidance is provided in TGP/9 "Examining Distinctness".

- 3.3 Conditions for Conducting the Examination
- [3.3.1] The tests should be carried out under conditions ensuring satisfactory growth for the expression of the relevant characteristics of the variety and for the conduct of the examination.

{ ASW 4 (Chapter 3.3) – conditions for conducting the examination }

{ GN 9 (Chapter 3.3) – growth stage key}

GN 9 (TG Template: Chapter 3.3) – Growth stage key

In cases where it is appropriate to provide a growth stage key for the observation of characteristics, the following is a useful guide:

"Growth stages of mono-and dicotyledonous plants - BBCH Monograph" (Federal Biological Research Centre for Agriculture and Forestry)
ISBN Number: 3-8263-3152-4

http://www.bba.de/veroeff/bbch/bbcheng.pdf

- 8. The Annex to this document provides examples of growth stage reference in existing Test Guidelines as follows:
 - TG/7/10: Pea
 - TG/4/8: Ryegrass
 - TG/276/1: Hemp

- TG/PINEAP(proj.12): Pineapple

PROPOSALS FOR INDICATING GROWTH STAGE AT WHICH TO OBSERVE CHARACTERISTICS IN THE TEST GUIDELINES

- 9. The guidance in document TGP/7 indicating the growth stage at which to observe characteristics in the Test Guidelines clarifies that such indication should be provided in the relevant cell of the Table of Characteristics (see paragraph 5) and accompanied by an explanation of the growth stage key in a separate subchapter of Chapter 8 of the Test Guidelines.
 - 10. The TWF is invited to consider whether there is a need to amend the existing guidance in document TGP/7 with regard to the indication of growth stage at which to observe characteristics in the Test Guidelines.

[Annex follows]

TWF/44/11

ANNEX

EXAMPLES OF INDICATIONS OF GROWTH STAGE AT WHICH TO OBSERVE CHARACTERISTICS IN EXISTING TEST GUIDELINES

EXTRACT FROM TG/7/10: PEA

TG/7/10 Pea, 2009-04-01 - 4 -

- 3.3 Conditions for Conducting the Examination
- 3.3.1 The tests should be carried out under conditions ensuring satisfactory growth for the expression of the relevant characteristics of the variety and for the conduct of the examination.
- 3.3.2 The optimum stage of development for the assessment of each characteristic is indicated by a number in the second column of the Table of Characteristics. The stages of development denoted by each number are described at the end of Chapter 8.

TG/7/10 Pea/Pois/Erbse/Guisante, 2009-04-01 Example Varieties/ Exemples/ Beispielssorten/ English français deutsch español Note/ Nota Variedades ejemplo 9. 200- Leaf: maximum Feuille : nombre Blatt: maximale Hoja: número maximum de folioles Anzahl Blattfiedern | máximo de folíolos 240 number of leaflets MS/ VG QN few petit gering bajo Jof 3 Dark Skin Perfection, 5 medium mittel medio moyen Finale 7 alto Ultimo many grand groß 10. 216- Leaflet: size Foliole: taille Blattfieder: Größe Folíolo: tamaño 226 MS/ QN sehr klein Payette 1 very small très petite muy pequeño (a) small klein Mini 3 petite pequeño medium moyenne mittel medio Finale 5 7 grande groß grande Alderman large Mammoth Melting 9 very large très grande sehr groß muy grande Sugar 11. 216- Leaflet: length Foliole: longueur Blattfieder: Länge Folíolo: longitud 226 MS/ VG 3 ON short courte kurz corto Eagle, Polar (a)

TG/7/10 Pea, 2009-04-01 - 38 -

KEY FOR THE GROWTH STAGES CLE POUR LES STADES DE CROISSANCE SCHLÜSSEL FÜR DIE ENTWICKLUNGSSTADIEN CLAVE PARA LOS ESTADOS DE DESARROLLO

Key Clé Schlüssel Clave	General Description	Description générale	Allgemeine Beschreibung	Descripción general
0	Germination	Germination	Keimung	Germinación
00	Dry seed	Graine sèche	Trockenkorn	Semilla seca
10	Seedling growth	Croissance de la plantule	Wachstum des Keimlings	Desarrollo de las plántulas
16	Young seedling with first scale leaf developed	Jeune plantule avec première feuille à écailles développée	Junger Keimling mit ersten entwickelten Schuppenblättern	Plántula joven con la primera hoja escamosa desarrollada
18	Young seedling with second scale leaf developed	Jeune plantule avec deuxième feuille à écailles développée	Junger Keimling mit zweiten entwickelten Schuppenblättern	Plántula joven con la segunda hoja escamosa desarrollada
20	First pair of stipules at the third node fully opened	Première paire de stipules au niveau du troisième noeud complètement ouverte	Erstes Paar Nebenblätter am dritten Knoten voll geöffnet	Primer par de estípulas en el nivel del tercer nudo completamente abiertas
22	Stipules at the fourth node fully opened	Stipules au niveau du quatrième noeud complètement ouverts	Nebenblätter am vierten Knoten voll geöffnet	Estípulas en el nivel del cuarto nudo completamente abiertas
25	Stipules at the fifth node fully opened	Stipules au niveau du cinquième noeud complètement ouverts	Nebenblätter am fünften Knoten voll geöffnet	Estípulas en el nivel del quinto nudo completamente abiertas

TG/7/10 Pea, 2009-04-01 - 39 -

Key Clé Schlüssel Clave	General Description	Description générale	Allgemeine Beschreibung	Descripción general
210	Emergence of first flower bud from stipules	Apparition du premier bourgeon à fleurs hors des stipules	Erscheinen der ersten Blütenknospe aus den Nebenblättern	Aparición de la primera yema floral fuera de las hojas escamosas
212	Emergence of standards from the calyx	Apparition des étendards hors du calice	Erscheinen der Fahne aus dem Kelch	Aparición de los estandartes fuera del cáliz
214	Opening of the standards and emergence of the wings	Ouverture des étendards et apparition des ailes	Oeffnen der Fahne und Erscheinen der Flügel	Apertura de los estandartes y aparición de las alas
216	Slight opening of the wings to show the keel	Légère ouverture des ailes découvrant la carène	Leichtes Oeffnen der Flügel und Erscheinen des Kieles	Ligera apertura de las alas para mostrar la quilla
218	Standards usually fully opened	Etendards généralement complètement ouverts	Fahnen normalerweise voll geöffnet	Estandartes normalmente abiertos por completo
220	Standards beginning to crumple at the margins	Etendards commençant à se friper sur les bords	Fahnen beginnen am Rand zu kräuseln	Los estandartes comienzan a arrugarse en los bordes
222	Standards and wings showing signs of withering	Etendards et ailes présentant des signes de flétrissure	Fahnen und Flügel weisen Zeichen des Welkens auf	Los estandartes y las alas presentan signos de marchitez
224	Emergence of the first flat pod	Apparition de la première gousse aplatie	Erscheinen der ersten flachen Hülse	Aparición de la primera vaina plana

EXTRACT FROM TG/4/8: RYEGRASS

TG/4/8 Ryegrass, 2006-04-05 - 4 -

3.3.2 The optimum stage of development for the assessment of each characteristic is indicated by a number in the second column of the Table of Characteristics. The stages of development denoted by each number are described at the end of Chapter 8.

	TG/4/8 Ryegrass/Ray-grass/Weidelgras/Raygrás, 2006-04-05 - 9 -							
		English	français	deutsch	español	Example Varieties/ Exemples/ Beispielssorten/ Variedades ejemplo	Note/ Nota	
4.	20-29 VG B	Leaf: width (at vegetative stage)	Feuille: largeur (au stade végétatif)	Blatt: Breite (im vegetativen Stadium)	Hoja: anchura (en estado vegetativo)			
QN		very narrow	très étroite	sehr schmal	muy estrecha		1	
		narrow	étroite	schmal	estrecha	Aragon (Lp)	3	
		medium	moyenne	mittel	media	Mondial (Lp)	5	
		broad	large	breit	ancha	Baroldi (Lmw), Veritas (Lp)	7	
		very broad	très large	sehr breit	muy ancha	Lipo (Lmi), Promenade (Lmw)	9	

TG/4/8 Ryegrass, 2006-04-05 - 19 -

8.3 Growth stages for grasses

All characteristics should be recorded at the appropriate time for the plant concerned. Growth stages of grasses are indicated by decimal codes which are derived from the decimal code for the growth stages of cereals (Zadoks, et al., 1974). This decimal code is in close conformity with the BBCH-code (Meier, 1997).

- Seedling growth (seedling: one shoot)
- DC 10 First leaf through coleoptile
- DC 15 Five leaves unfolded
- DC 19 Nine or more leaves unfolded

Tillering

- DC 20 Main shoot only (beginning of tillering)
- DC 23 Main shoot and 3 tillers
- DC 25 Main shoot and 5 tillers
- DC 29 Main shoot and 9 or more tillers

EXTRACT FROM TG/276/1: HEMP

TG/276/1 Hemp/Chanvre/Hanf/Cáñamo, 2012-03-28 - 9 -

7. Table of Characteristics/Tableau des caractères/Merkmalstabelle/Tabla de caracteres

		English	français	deutsch	español	Example Varieties/ Exemples/ Beispielssorten/ Variedades ejemplo	Note Nota
1. (+)	0003 VG	Cotyledon: shape	Cotylédon : forme	Keimblatt: Form	Cotiledón: forma		
QN	C	narrow obovate	obovale étroit	schmal verkehrt eiförmig	oboval estrecha	Fibrimon	1
		medium obovate	obovale moyen	mittel verkehrt eiförmig	oboval media	Epsilon 68	2
		broad obovate	obovale large	breit verkehrt eiförmig	oboval ancha	Futura 75	3
2.	0003 VG	Cotyledon: color	Cotylédon : couleur	Keimblatt: Farbe	Cotiledón: color		
PQ	C	yellow	jaune	gelb	amarillo	Chamaeleon	1
		light green	vert clair	hellgrün	verde claro	Fedora 17	2
		medium green	vert moyen	mittelgrün	verde medio	Ferimon	3
		dark green	vert foncé	dunkelgrün	verde oscuro	Dioica 88	4

TG/276/1 Hemp, 2012-03-28 - 19 -

Principal growth stages

Four principal stages describe the life cycle of a plant and are coded by their first digit of the four-digit code.

First-digit of code	Definition
0	Germination and emergence
1	Vegetative stage
2	Flowering and seed formation
3	Senescence

Secondary growth stages

The secondary growth stages are described by the second digit, which indicates the sex of the plant, the third and fourth digits indicating the developmental stage of the plant.

Code	Definition	Remarks			
Germina	tion and emergence				
0000	Dry seed				
0003	Cotyledons unfolded				
Vegetativ	Vegetative stage refers to main stem. Leaves are considered unfolded when leaflets are at least one cm long				
1002	1st leaf pair	1 leaflet			
1004	2 nd leaf pair	3 leaflets			

EXTRACT FROM TG/PINEAP(proj.12): PINEAPPLE

Pflanze: Anzahl der

Blätter

gering

mittel

groß

kurz

lang

Blatt: Länge

Planta: número de

Perola

BRS Imperial, Gold,

Smooth Cayenne

Smooth Cayenne

Aus-Carnival, Perola

Gomo de Mel

Queen

hojas

bajo

medio

alto

corta

media

larga

Hoja: longitud

7.	TG/PINEAP(proj. 12) Pineapple/Ananas/Ananas/Piña, 2013-02-12 - 8 - 7. Table of Characteristics/Tableau des caractères/Merkmalstabelle/Tabla de caracteres						
		English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	
1.	VG	Plant: growth habit	Plante : port	Pflanze: Wuchsform	Planta: porte		
(*) (+)	1-T						
QN	(a)	upright	dressé	aufrecht	erguido	Perola	
		semi upright	demi-dressé	halbaufrecht	semierguido	Smooth Cayenne	
		spreading	étalé	breitwüchsig	extendido	Perolera	

Plante : nombre de

feuilles

faible

moyen

élevé

petite

moyenne

grande

Feuille : longueur

TG/PINEAP(proj.12) Pineapple, 2013-02-12

8. Explanations on the Table of Characteristics

VGI

MS

1-T

(a) few

MS 1-T

(a)

QN

(*) (+)

QN

Plant: number of

leaves

medium

many

short medium

long

Leaf: length

8.1 Explanations covering several characteristics

Characteristics containing the following key in the second column of the Table of Characteristics should be examined as indicated below:

- (Characteristics 1 to 13) Should be observed at the time floral induction is provoked (a) artificially (stage 1-T), which is generally about 36 to 54 weeks after planting, depending on the location and the varieties.
- (Characteristics 14 to 18) Observations related to flowering, inflorescence and flowers (b) should be made on 10 inflorescences, at the time of anthesis (stage 2-A). Measurements of floral parts to be taken on 10 flowers removed at mid-anthesis.
- (c) (Characteristic 19) Observations of fruits before maturity should be made on 10 fruits, 4-6 months after floral induction is provoked (immature fruit-stage 3-I), at maximum size before the fruits starts to mature.
- (Characteristics 20 to 48) Observations related to plant and fruit at harvest should be made (d) in the plot on 10 plants and 10 fruits. It is considered that harvest time is the stage at which the fruit is ready for consumption (actual maturity-stage 4-M). Measures to be made on 10 fruits.

Growth stages

- 1-T: At fully vegetative growth stage, before flower emergence
- 2-A: Anthesis stage
- 3-I: Immature fruit stage, before physiologically ripe
- 4-M: Maturity stage, when physiologically ripe

Note/

Nota

3 5

3

5