

**TWO/33/2****ORIGINAL:** English**DATE:** May 15, 2000

INTERNATIONAL UNION FOR THE PROTECTION OF NEW VARIETIES OF PLANTS
GENEVA

**TECHNICAL WORKING PARTY
FOR
ORNAMENTAL PLANTS AND FOREST TREES**

**Thirty-Third Session
Budapest, June 26 to 30, 2000**

WORKING PAPER ON REVISED DRAFT TEST GUIDELINES FOR EUSTOMA
(*Eustoma grandiflorum* (Raf.) Shinnery)

Document prepared by experts from Japan

<u>TABLE OF CONTENTS</u>	<u>PAGE</u>
I. Subject of these Guidelines	3
II. Material Required	3
III. Conduct of Tests	3
IV. Methods and Observations.....	4
V. Grouping of Varieties	4
VI. Characteristics and Symbols	5
VII. Table of Characteristics	6
VIII. Explanations on the Table of Characteristics	12
IX. Literature	13
X. Technical Questionnaire	14

I. Subject of these Guidelines

These Test Guidelines apply to all varieties of *Eustoma grandiflorum* (Raf.) Shinn. of the family Gentianaceae.

II. Material Required

1. The competent authorities decide when, where and in what quantity and quality the plant material /seed required for testing the variety is to be delivered. Applicants submitting material from a State other than that in which the testing takes place must make sure that all customs formalities are complied with. As a minimum, the following quantity of plant material is recommended:

- (a) seed propagated varieties: sufficient seeds to propagate 60 plants.
- (b) vegetatively propagated varieties: 40 plantlets

2. The plant material/seed supplied should be visibly healthy, not lacking in vigor or affected by any important pests or diseases. The quality of the seed to be delivered should not be below the standards of seeds for certification or marketing in the country concerned. The germination capacity should be stated.

3. The plant material/seed must not have undergone any treatment unless the competent authorities allow or request such treatment. If it has been treated, full details of the treatment must be given.

III. Conduct of Tests

1. To assess distinctness, it is essential that the plants under test should produce at least two satisfactory crops of fruit.

2. The tests should normally be conducted at one place. If any important characteristics of the variety cannot be seen at that place, the variety may be tested at an additional place.

Sowing: time: February to March (in Northern hemisphere)
temperature: 15°C minimum
soil: a well drained fertile soil, pH 6.0 to 6.5
use cell tray (3 x 3 x 4)
no soil covering
subirrigation

Planting: time: April to May
plant stage: 4 true leaves
soil: well-drained and moisture holding fertile soil, rich in organic material,
pH 6.0 to 6.5

Planting density: 12 x 12 cm

Temperature: day: +20 to 25°C
night: +10 to 15°C

3. The tests should be carried out in the green house under the conditions ensuring normal growth. The size of the plots should be such that plants or parts of plants may be removed for measurement and counting without prejudice to the observations which must be made up to the end of the growing period. As a minimum, each test should include a total of 60 plants for seed propagated varieties and 40 plants for vegetatively propagated varieties which should be divided between two or more replicates. Separate plots for observation and measuring can only be used if they have been subject to similar environmental conditions.
4. Additional tests for special purposes may be established.

IV. Methods and Observations

1. All observations should be made on 40 plants or 40 parts of plants.
2. For the assessment of uniformity a population standard of 2% and an acceptance probability of at least 95% should be applied in the case of vegetatively propagated varieties. In the case of a sample size of 40 plants, the maximum number of off-types allowed would be 2.
3. All observations should be made on flowering plants.
4. All observations on the leaf should be made on the third leaf from the top. For determining the color of the leaf and the stem the bloom should be removed.
5. Length and color of internode should be observed on the fourth internode from the top.
6. All observations on the flower and the pedicel should be made on the second flower to open. Color observation on the petal should be made on its inner side.
7. Because daylight varies, color determinations made against a color chart should be made either in a suitable cabinet providing artificial daylight or in the middle of the day in a room without direct sunlight. The spectral distribution of the illuminant for artificial daylight should conform with the CIE Standard of Preferred Daylight D 6500 and should fall within the tolerances set out in British Standard 950, Part I. These determinations should be made with the plant part placed against a white background.

V. Grouping of Varieties

1. The collection of varieties to be grown should be divided into groups to facilitate the assessment of distinctness. Characteristics which are suitable for grouping purposes are those which are known from experience not to vary, or to vary only slightly, within a variety. Their various states of expression should be fairly evenly distributed throughout the collection.

2. It is recommended that the competent authorities use the following characteristics for grouping varieties:

- (a) Plant: height (characteristic 1)
- (b) Flower: diameter (characteristic 17)
- (c) Flower: shape (characteristic 18)
- (d) Self-colored varieties only: Petal: color (characteristic 23)
- (e) Bi-colored varieties only: Petal: color (characteristic 24)
- (f) Flower: color of base (characteristic 28)

VI. Characteristics and Symbols

1. To assess distinctness, uniformity and stability, the characteristics and their states as given in the Table of Characteristics should be used.

2. Notes (numbers), for the purposes of electronic data processing, are given opposite the states of expression for each characteristic.

3. Legend

(*) Characteristics that should be used on all varieties in every growing period over which examinations are made and always be included in the variety descriptions, except when the state of expression of a preceding characteristic or regional environmental conditions render this impossible.

(+) See Explanations on the Table of Characteristics in Chapter VIII.

VII. Table of Characteristics/Tableau des caractères/Merkmalstabelle/Tabla de caracteres

English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
1. Plant: height					
(*)					
short				White Coronet	3
medium				Momo Sen, Deep Purple	5
tall				Yuki no Mine	7
2. Stem: thickness					
thin				White Coronet	3
medium				Momo Sen	5
thick				Yuki no Mine	7
3. Stem: number of nodes					
few				White Coronet	3
medium				Momo Sen	5
many				Purple Robin	7
4. Stem: length of fourth internode below the top flower					
short				White Coronet	3
medium				Momo Sen	5
long					7
5. Stem: number of branches					
few				White Coronet	3
medium				Momo Sen	5
many				Purple Robin	7

English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
6. Stem: green color					
light					3
medium				Yuki no Mine	5
dark				Fuku Shihai	7
7. Stem: position of branching					
upper part only				Purple moon	1
upper middle part				Momo Sen	2
whole stem					3
8. Leaf: attitude to the stem (+)					
semi-erect				White Coronet	1
horizontal				Momo Sen	2
semi-drooping					3
9. Leaf: length (*) (+)					
short				White Colonet	3
medium				Momo Sen	5
long					7
10. Leaf: width (*) (+)					
narrow				White Colonet	3
medium				Momo Sen	5
broad					7

English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
11. Leaf: shape					
(*)					
(+)					
lanceolate					1
ovate				Momo Sen	2
broad ovate					3
12. Leaf: bloom					
(*)					
absent					1
present					9
13. Leaf: green color of upper side (without bloom)					
(*)					
light					3
medium				Momo Sen	5
dark					7
14. Leaf: green color of lower side (without bloom)					
(*)					
light					3
medium					5
dark					7
15. Flower: type					
(*)					
single					1
double					2


English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
16. Variety with double flowers only: (*) Flower: number of petals					
few					3
medium				King of Blue Flash	5
many				Deep Purple	7
17. Flower: diameter (*)					
small					3
medium				Momo Sen	5
large				Deep Purple	7
18. Flower: shape					
campanulate				Momo Sen	1
funnel-shaped				Fuku Shihai	2
19. Petal: length (*)					
short					3
medium				Momo Sen	5
long				Yuki no Mine	7
20. Petal: width (*)					
narrow				Deep Purple	3
medium				Yuki no Mine	5
broad					7
21. Petal: undulation of margin					
weak				Momo Sen	3
medium				Yuki no Mine	5
strong					7

English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
22. Petal: color pattern (*)					
self-colored					1
bi-colored					2
23. Self-colored varieties only: Petal: color (*)					
RHS Colour Chart (indicate reference number)					
24. Bi-colored varieties only: Petal: main color (*)					
RHS Colour Chart (indicate reference number)					
25. Bi-colored varieties only: Petal: secondary color (*)					
RHS Colour Chart (indicate reference number)					
26. Petal: relative area of secondary color (*)					
small					3
medium					5
large					7
27. Petal: position of secondary color					
upper third					1
middle					2
lower third					3

English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
28. Petal: color of base					
(*)					
green				Haku Sen	1
brown				Fuku Shihai	2
29. Calyx: length					
(+)					
short					3
medium				Yuki no Mine	5
long					7
30. Calyx: anthocyanin coloration					
absent				Haku Sen	1
present				Shi Sen	9
31. Sepal: relative attitude to petal					
(+)					
adpressed					1
spreading					2
32. Pedicel: length					
short				White Coronet	3
medium				Haku Sen	5
long					7
33. Time of beginning of flowering					
(*)					
early				Azuma no Yosooi	3
medium				Haku Sen	5
late				Fuku Shihai	7

VIII. Explanations on the Table of Characteristics

Ad. 18: Flower: shape




1
campanurate




2
funnel-shaped


Ad. 29: Calyx: length



Ad. 31: Sepal: relative attitude to petal



1
adpressed



2
spreading

IX. Literature

Kiyoshi Okawa, 1992: Eustoma (Torukogikyo) Seibunndo-Shinkosya Co., Tokyo, JP

X. Technical Questionnaire

	Reference Number (not to be filled in by the applicant)
<p>TECHNICAL QUESTIONNAIRE to be completed in connection with an application for plant breeders' rights</p>	
1. Species	<p><i>Eustoma grandiflorum</i> (Raf.) Shinnery EUSTOMA</p>
2. Applicant (Name and address)	
3. Proposed denomination or breeder's reference	

4. Information on origin, maintenance and reproduction of the variety

4.1 Origin

(a) Seedling of unknown parentage

..... []

(b) Produced by controlled pollination (indicate parent varieties)

- Seed bearing parent (indicate parent)

..... []

- Pollen parent (indicate parent)

..... []

(c) Produced by open pollination of (indicate seed bearing parent plant)

..... []

(d) Mutation or sport from (indicate original parent variety)

..... []

(e) Discovery (indicate where and when)

..... []

4.2 *In vitro* propagation:

The plant material of the candidate variety has been obtained by *in vitro* propagation

yes []

no []

4.3 Other information

5. Characteristics of the variety to be indicated (the number in brackets refers to the corresponding characteristic in Test Guidelines; please mark the state of expression which best corresponds).

Characteristics	Example Varieties	Note
5.1 Plant: height (1)		
low	White Coronet	3[]
medium	Momo Sen	5[]
high	Yuki no Mine	7[]
5.2 Flower: diameter (17)		
small	Minetta	3[]
medium	Momo Sen	5[]
large	Deep Purple	7[]
5.3 Flower: shape (18)		
campanulate	Momo Sen	1[]
funnel-shaped	Fuku Shihai	2[]
5.4 Petal: color of base (28)		
green	Haku Sen	1[]
brown	Fuku Shihai	2[]
5.5 Self-colored variety only: Petal: color (23)		
RHS Colour Chart (indicate reference number)		
5.6 Bi-color variety only: Petal: main color (24)		
RHS Colour Chart (indicate reference number)		

6. Similar varieties and differences from these varieties

Denomination of similar variety	Characteristic in which the similar variety is different ^{o)}	State of expression of similar variety	State of expression of candidate variety
---------------------------------	--	--	--

^{o)} In the case of identical states of expressions of both varieties, please indicate the size of the difference.

7. Additional information which may help to distinguish the variety

7.1 Resistance to pests and diseases

7.2 Special conditions for the examination of the variety

7.3 Other information

A representative color photo of the variety should be added to the Technical Questionnaire.

8. Authorization for release

- (a) Does the variety require prior authorization for release under legislation concerning the protection of the environment, human and animal health?

Yes [] No []

- (b) Has such authorization been obtained?

Yes [] No []

If the answer to that question is yes, please attach a copy of such an authorization.

[End of document]