

TG/TOREN(proj.2)
ORIGINAL: English
DATE: 2010-08-02

INTERNATIONAL UNION FOR THE PROTECTION OF NEW VARIETIES OF PLANTS GENEVA



TORENIA

UPOV Code: TOREN

Torenia L.

GUIDELINES

FOR THE CONDUCT OF TESTS

FOR DISTINCTNESS, UNIFORMITY AND STABILITY

prepared by an expert from Japan

to be considered by the

Technical Working Party for Ornamental Plants and Forest Trees at its forty-third session, to be held in Cuernavaca, Morelos State, Mexico, from September 20 to 24, 2010

Alternative Names:*

| Botanical name | English | French | German | Spanish |
|----------------|--|---------|---------|-----------------------------|
| Torenia L. | Torenia, Bluewings, Wishbone-flower | Torenia | Torenie | Torenia, Legazpia blanco |

The purpose of these guidelines ("Test Guidelines") is to elaborate the principles contained in the General Introduction (document TG/1/3), and its associated TGP documents, into detailed practical guidance for the harmonized examination of distinctness, uniformity and stability (DUS) and, in particular, to identify appropriate characteristics for the examination of DUS and production of harmonized variety descriptions.

ASSOCIATED DOCUMENTS

These Test Guidelines should be read in conjunction with the General Introduction and its associated TGP documents.

*

These names were correct at the time of the introduction of these Test Guidelines but may be revised or updated. [Readers are advised to consult the UPOV Code, which can be found on the UPOV Website (www.upov.int), for the latest information.]

TG/TOREN(proj.2) Torenia, 2010-08-02 - 2 -

| <u>TA</u> | BLE OF CONTENTS | <u>PAGE</u> |
|-----------|---|-------------|
| 1. | SUBJECT OF THESE TEST GUIDELINES | 3 |
| 2. | MATERIAL REQUIRED | 3 |
| 3. | METHOD OF EXAMINATION | 3 |
| | 3.1 Number of Growing Cycles | 3 |
| | 3.2 Testing Place | |
| | 3.3 Conditions for Conducting the Examination | 3 |
| | 3.4 Test Design | 4 |
| | 3.5 Additional Tests | |
| 4. | ASSESSMENT OF DISTINCTNESS, UNIFORMITY AND STABILITY | 4 |
| | 4.1 Distinctness | 4 |
| | 4.2 Uniformity | |
| | 4.3 Stability | 6 |
| 5. | GROUPING OF VARIETIES AND ORGANIZATION OF THE GROWING TRIAL | 6 |
| 6. | INTRODUCTION TO THE TABLE OF CHARACTERISTICS | 7 |
| | 6.1 Categories of Characteristics | 7 |
| | 6.2 States of Expression and Corresponding Notes | 7 |
| | 6.3 Types of Expression | 8 |
| | 6.4 Example Varieties | 8 |
| | 6.5 Legend | 8 |
| 7. | TABLE OF CHARACTERISTICS/TABLEAU DES CARACTERES/MERKMALSTABELLE/TABLA DE CARACTERES | 10 |
| 8. | EXPLANATIONS ON THE TABLE OF CHARACTERISTICS | |
| 0. | 8.1 Explanations covering several characteristics | |
| | 8.2 Explanations for individual characteristics | |
| 9. | LITERATURE | |
| - | TECHNICAL OLIESTIONNAIDE | |

- 3 -

1. <u>Subject of these Test Guidelines</u>

These Test Guidelines apply to all varieties of *Torenia* L. of the family Scrophulariaceae.

2. Material Required

- 2.1 The competent authorities decide on the quantity and quality of the plant material required for testing the variety and when and where it is to be delivered. Applicants submitting material from a State other than that in which the testing takes place must ensure that all customs formalities and phytosanitary requirements are complied with.
- 2.2 The material is to be supplied in the form of seed or rooted cuttings.
- 2.3 The minimum quantity of plant material, to be supplied by the applicant, should be:

Vegetatively propagated varieties: 10 rooted cuttings Seed-propagated varieties: sufficient seed to produce 20 plants

In the case of seed, the seed should meet the minimum requirements for germination, species and analytical purity, health and moisture content, specified by the competent authority. In cases where the seed is to be stored, the germination capacity should be as high as possible and should be stated by the applicant.

- 2.4 The plant material supplied should be visibly healthy, not lacking in vigor, nor affected by any important pest or disease.
- 2.5 The plant material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If it has been treated, full details of the treatment must be given.

3. Method of Examination

3.1 Number of Growing Cycles

The minimum duration of tests should normally be a single growing cycle.

3.2 Testing Place

Tests are normally conducted at one place. In the case of tests conducted at more than one place, guidance is provided in TGP/9 "Examining Distinctness".

3.3 Conditions for Conducting the Examination

3.3.1 The tests should be carried out under conditions ensuring satisfactory growth for the expression of the relevant characteristics of the variety and for the conduct of the examination.

3.3.2 Observation of color by eye

Because daylight varies, color determinations made against a color chart should be made either in a suitable cabinet providing artificial daylight or in the middle of the day in a room without direct sunlight. The spectral distribution of the illuminant for artificial daylight should conform with the CIE Standard of Preferred Daylight D 6500 and should fall within the tolerances set out in the British Standard 950, Part I. These determinations should be made with the plant part placed against a white background. The color chart and version used should be specified in the variety description.

3.4 Test Design

- 3.4.1 Each test should be designed to result in a total of at least 10 plants for vegetatively propagated varieties or 20 plants for seed propagated varieties.
- 3.4.2 The design of the tests should be such that plants or parts of plants may be removed for measurement or counting without prejudice to the observations which must be made up to the end of the growing cycle.

3.5 Additional Tests

Additional tests, for examining relevant characteristics, may be established.

4. Assessment of Distinctness, Uniformity and Stability

4.1 Distinctness

4.1.1 General Recommendations

It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding distinctness. However, the following points are provided for elaboration or emphasis in these Test Guidelines.

4.1.2 Consistent Differences

The differences observed between varieties may be so clear that more than one growing cycle is not necessary. In addition, in some circumstances, the influence of the environment is not such that more than a single growing cycle is required to provide assurance that the differences observed between varieties are sufficiently consistent. One means of ensuring that a difference in a characteristic, observed in a growing trial, is sufficiently consistent is to examine the characteristic in at least two independent growing cycles.

4.1.3 Clear Differences

Determining whether a difference between two varieties is clear depends on many factors, and should consider, in particular, the type of expression of the characteristic being examined, i.e. whether it is expressed in a qualitative, quantitative, or pseudo-qualitative manner. Therefore, it is important that users of these Test Guidelines are familiar with the

recommendations contained in the General Introduction prior to making decisions regarding distinctness.

4.1.4 Number of Plants / Parts of Plants to be Examined

Unless otherwise indicated, all observations for the purposes of distinctness should be made on 10 plants or parts taken from each of 10 plants for vegetatively propagated varieties or 20 plants or parts of plants taken from 20 plants for seed propagated varieties, disregarding any off-type plants.

4.1.5 Method of Observation

The recommended method of observing the characteristic for the purposes of distinctness is indicated by the following key in the second column of the Table of Characteristics (see document TGP/9 "Examining Distinctness", Section 4 "Observation of characteristics"):

MG: single measurement of a group of plants or parts of plants

MS: measurement of a number of individual plants or parts of plants

VG: visual assessment by a single observation of a group of plants or parts of plants

VS: visual assessment by observation of individual plants or parts of plants

Type of observation: visual (V) or measurement (M)

"Visual" observation (V) is an observation made on the basis of the expert's judgment. For the purposes of this document, "visual" observation refers to the sensory observations of the experts and, therefore, also includes smell, taste and touch. Visual observation includes observations where the expert uses reference points (e.g. diagrams, example varieties, side-by-side comparison) or non-linear charts (e.g. color charts). Measurement (M) is an objective observation against a calibrated, linear scale e.g. using a ruler, weighing scales, colorimeter, dates, counts, etc.

Type of record: for a group of plants (G) or for single, individual plants (S)

For the purposes of distinctness, observations may be recorded as a single record for a group of plants or parts of plants (G), or may be recorded as records for a number of single, individual plants or parts of plants (S). In most cases, "G" provides a single record per variety and it is not possible or necessary to apply statistical methods in a plant-by-plant analysis for the assessment of distinctness."

In cases where more than one method of observing the characteristic is indicated in the Table of Characteristics (e.g. VG/MG), guidance on selecting an appropriate method is provided in document TGP/9, Section 4.2.

4.2 Uniformity

4.2.1 It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding uniformity. However, the following points are provided for elaboration or emphasis in these Test Guidelines:

- 4.2.2 For the assessment of uniformity of vegetatively propagated varieties, a population standard of 1% and an acceptance probability of at least 95 % should be applied. In the case of a sample size of 10 plants, one off-type is allowed.
- 4.2.3 For the assessment of uniformity of seed-propagated varieties which are self-pollinated, a population standard of 1% and an acceptance probability of at least 95% should be applied. In the case of a sample size of 20 plants, one off-type is allowed.
- 4.2.4 The assessment of uniformity for hybrid varieties depends on the type of hybrid and should be according to the recommendations for hybrid varieties in the General Introduction.

4.3 Stability

- 4.3.1 In practice, it is not usual to perform tests of stability that produce results as certain as those of the testing of distinctness and uniformity. However, experience has demonstrated that, for many types of variety, when a variety has been shown to be uniform, it can also be considered to be stable.
- 4.3.2 Where appropriate, or in cases of doubt, stability may be further examined by testing a new seed or plant stock to ensure that it exhibits the same characteristics as those shown by the initial material supplied.

5. Grouping of Varieties and Organization of the Growing Trial

- 5.1 The selection of varieties of common knowledge to be grown in the trial with the candidate varieties and the way in which these varieties are divided into groups to facilitate the assessment of distinctness are aided by the use of grouping characteristics.
- 5.2 Grouping characteristics are those in which the documented states of expression, even where produced at different locations, can be used, either individually or in combination with other such characteristics: (a) to select varieties of common knowledge that can be excluded from the growing trial used for examination of distinctness; and (b) to organize the growing trial so that similar varieties are grouped together.
- 5.3 The following have been agreed as useful grouping characteristics:
 - (a) Plant: growth habit (characteristic 1)
 - (b) Upper corolla lobe: color of basal part (characteristic 21) with the following groups:

Gr.1: white

Gr.2: yellow

Gr.3: orange

Gr.4: red

Gr.5: pink

Gr.6: purple

Gr.7: violet

- 7 -

(c) Upper corolla lobe: color of distal part (characteristic 22) with the following groups:

Gr.1: white

Gr.2: yellow

Gr.3: orange

Gr.4: red

Gr.5: pink

Gr.6: purple

Gr.7: violet

(d) Lateral corolla lobe: color of central part (characteristic 23) with the following groups:

Gr.1: white

Gr.3: yellow

Gr.4: orange

Gr.5: red

Gr.2: pink

Gr.6: purple

Gr.7: violet

(e) Lower corolla lobe: color of blotch (characteristic 27) with the following groups:

Gr.1: white

Gr.2: yellow

Guidance for the use of grouping characteristics, in the process of examining distinctness, is provided through the General Introduction and document TGP/9 "Examining Distinctness".

6. Introduction to the Table of Characteristics

6.1 Categories of Characteristics

6.1.1 Standard Test Guidelines Characteristics

Standard Test Guidelines characteristics are those which are approved by UPOV for examination of DUS and from which members of the Union can select those suitable for their particular circumstances.

6.1.2 Asterisked Characteristics

Asterisked characteristics (denoted by *) are those included in the Test Guidelines which are important for the international harmonization of variety descriptions and should always be examined for DUS and included in the variety description by all members of the Union, except when the state of expression of a preceding characteristic or regional environmental conditions render this inappropriate.

6.2 States of Expression and Corresponding Notes

6.2.1 States of expression are given for each characteristic to define the characteristic and to harmonize descriptions. Each state of expression is allocated a corresponding numerical note for ease of recording of data and for the production and exchange of the description.

6.2.2 In the case of qualitative and pseudo-qualitative characteristics (see Chapter 6.3), all relevant states of expression are presented in the characteristic. However, in the case of quantitative characteristics with 5 or more states, an abbreviated scale may be used to minimize the size of the Table of Characteristics. For example, in the case of a quantitative characteristic with 9 states, the presentation of states of expression in the Test Guidelines may be abbreviated as follows:

| State | Note |
|--------|------|
| small | 3 |
| medium | 5 |
| large | 7 |

However, it should be noted that all of the following 9 states of expression exist to describe varieties and should be used as appropriate:

| State | Note |
|---------------------|------|
| very small | 1 |
| very small to small | 2 |
| small | 3 |
| small to medium | 4 |
| medium | 5 |
| medium to large | 6 |
| large | 7 |
| large to very large | 8 |
| very large | 9 |

6.2.3 Further explanation of the presentation of states of expression and notes is provided in document TGP/7 "Development of Test Guidelines".

6.3 Types of Expression

An explanation of the types of expression of characteristics (qualitative, quantitative and pseudo-qualitative) is provided in the General Introduction.

6.4 Example Varieties

Where appropriate, example varieties are provided to clarify the states of expression of each characteristic.

6.5 Legend

- (*) Asterisked characteristic see Chapter 6.1.2
- QL Qualitative characteristic see Chapter 6.3
- QN Quantitative characteristic see Chapter 6.3
- PQ Pseudo-qualitative characteristic see Chapter 6.3

MG, MS, VG, VS – see Chapter 4.1.5

(a) See Explanations on the Table of Characteristics in Chapter 8.1

(+) See Explanations on the Table of Characteristics in Chapter 8.

7. <u>Table of Characteristics/Tableau des caractères/Merkmalstabelle/Tabla de caracteres</u>

| | | English | français | deutsch | español | Example Varieties Exemples Beispielssorten Variedades ejemplo | Note/ Nota |
|------------------|-----------|--|----------------|-------------------|---|--|---------------|
| 1. (*) (+) | VG | Plant: growth habit | to 9)? I can't | remember how y | have a wider range (1 ou grow the plants, if hould be possible to | | |
| PQ | | upright | observe a tra | ailing habit as w | ell. The characteristic | Crown Blue | 1 |
| | | semi upright | should be ind | icated as "QN". | | | 2 |
| | | horizontal | | | | Danmoon20 | 3 |
| 2. | | Plant: height | | | | | |
| (+) | MS | | | | | | |
| QN | | short | | | | Danmoon20 | 3 |
| | | medium | | | | Crown Blue | 5 |
| | | tall | | | | | 7 |
| 3. | VG/ MS | Plant: width | | | | | |
| (+) | WIS | | | | | | |
| QN | | narrow | | | | Crown Blue | 3 |
| | | medium | | | | | 5 |
| | | broad | | | | | 7 |
| 4. | VG | Shoot: anthocyanin coloration of nodes | our variet | ies. We have te | rved anthocyanin on sted 'Dantomidmon' reak expression of | | |
| QN | | absent or very weak | | | Is this characteristic | Crown Rose | 1 |
| | | weak | | | | Dantomidmon | 2 |
| | | medium | | | | Crown Blue | 3 |
| | | strong | | | | | 4 |
| 5. | VG/ MS | Petiole: length | | | | | |
| (+) | IVIS | | | | | | |
| QN | (a) | short | | | | | 1 |
| | | medium | | | | Danmoon20 | 2 |
| | | long | | | | Crown Blue | 3 |

| | | English | français | deutsch | español | Example Varieties Exemples Beispielssorten Variedades ejemplo | Note/ Nota |
|------------------|------------------------|---|----------|---------|---------|--|---------------|
| 6. (*) (+) | VG/ MS | Leaf blade: length | | | | | |
| QN | (a) | short | | | | | 3 |
| | | medium | | | | HATO-04-05 | 5 |
| | | long | | | | Crown Blue | 7 |
| 7. (*) (+) | <mark>VG/</mark> MS | Leaf blade: width | | | | | |
| QN | (a) | narrow | | | | | 3 |
| | | medium | | | | HATO-04-05 | 5 |
| | | broad | | | | Crown Blue | 7 |
| <mark>8</mark> . | | Leaf blade: depth of incision of margin | | | | | |
| | (a) | absent or very shallow | | | | | 1 |
| | | shallow | | | | | 3 |
| | | medium | | | | | 5 |
| | | deep | | | | | 7 |
| 9. | | Leaf blade: incisions of margin | 1 | | | | |
| (+) | IVID | meisions of margn | 1 | | | | |
| PQ | (a) | dentate | | | | Danmoon20 | 1 |
| | | crenate | | | | HATO-04-05 | 2 |
| 10. | VG | Leaf blade: intensity of green color on upper sid | e | | | | |
| QN | (a) | weak light | | | | | 3 |
| | | medium | | | | | 5 |
| | | strong dark | | | | | 7 |

TG/TOREN(proj.2) Torenia, 2010-08-02 - 12 -

| | | English | français | deutsch | español | Example Varieties Exemples Beispielssorten Variedades ejemplo | Note/ Nota |
|-------------------|----|-----------------------------------|----------|---------|---------|--|---------------|
| 11. (+) | VG | Inflorescence: axillary flower | | | | | |
| QL | | absent | | | | Crown Blue | 1 |
| | | present | | | | Danmoon20 | 9 |
| 12. (*) (+) | VG | Calyx: number of lobes | | | | | |
| QL | | two | | | | Crown Blue | 1 |
| | | five | | | | Danmoon20 | 2 |
| 13. (*) (+) | MS | Flower: length in front view | | | | | |
| QN | | short | | | | | 3 |
| | | medium | | | | Crown Blue | 5 |
| | | long | | | | | 7 |
| 14. (*) (+) | MS | Flower: width in front view | | | | | |
| QN | | narrow | | | | | 3 |
| | | medium | | | | Crown Blue | 5 |
| | | broad | | | | | 7 |
| 15. (+) | MS | Corolla tube: length | | | | | |
| QN | | short | | | | | 3 |
| | | medium | | | | | 5 |
| | | long | | | | | 7 |

| | | English | français | deutsch | español | Example Varieties Exemples Beispielssorten Variedades ejemplo | Note/ Nota |
|-------------------|----|--|-------------|---------------------|---------|--|---------------|
| 16. (+) | VG | Corolla tube: color of outer side | | | | | |
| PQ | | RHS Colour Chart (indicate reference number) | | | | | |
| 17. (+) | VG | Corolla tube: vertical lines of inner side on corolla | Whether the | stripes are truly Q | L? | | |
| QL | | absent | | | | Crown Blue | 1 |
| QL. | | present | | | | Danmoon20 | 9 |
| 18. | VC | Corolla tube: color | | | | Damiloon20 | |
| (+) | VG | of inner side at basal part | | | | | |
| PQ | | RHS Colour Chart (indicate reference number) | | | | | |
| 19. (*) (+) | VG | Corolla lobe: incision of margin | | | | | |
| QL | | absent | | | | Danmoon20 | 1 |
| | | present | | | | Danmoon15 | 9 |
| 20. (+) | VG | Upper corolla lobe: undulation | : | | | | |
| QN | | absent or weak | | | | Danmoon20 | 1 |
| | | medium | | | | | 2 |
| | | strong | | | | Crown Blue | 3 |
| 21. (*) (+) | VG | Upper corolla lobe: color of basal part | : | | | | |
| PQ | | RHS Colour Chart (indicate reference number) | | | | | |

TG/TOREN(proj.2) Torenia, 2010-08-02 - 14 -

| | | English | français | deutsch | español | Example Varieties Exemples Beispielssorten Variedades ejemplo | Note/ Nota |
|-------------------|----|--|----------|---------|---------|--|---------------|
| 22. (*) (+) | VG | Upper corolla lobe: color of distal part | | | | | |
| PQ | | RHS Colour Chart (indicate reference number) | | | | | |
| 23. (*) (+) | VG | Lateral corolla lobe: color of central part | | | | | |
| PQ | | RHS Colour Chart (indicate reference number) | | | | | |
| 24. (*) (+) | VG | Lateral corolla lobe: color of marginal part | | | | | |
| PQ | | RHS Colour Chart (indicate reference number) | | | | | |
| 25. (*) (+) | VG | Lower corolla lobe: color of <mark>distal</mark> part | | | | | |
| PQ | | RHS Colour Chart (indicate reference number) | | | | | |
| 26. | VG | Lower corolla lobe: conspicuousness of blotch | | | | | |
| QN | | absent or very weak | | | | | 1 |
| | | medium | | | | | 2 |
| | | strong | | | | | 3 |
| 27. (*) | VG | Lower corolla lobe: color of blotch | | | | | |
| PQ | | RHS Colour Chart (indicate reference number)white | | | | | |

8. Explanations on the Table of Characteristics

8.1 Explanations covering several characteristics


Unless otherwise indicated, all characteristics should be observed at the time of full flowering.

Characteristics containing the following key in the second column of the Table of Characteristics should be examined as indicated below:


(a) Observations on the leaf blade should be made on fully expanded leaves from the lower middle third of shoot.)

8.2 Explanations for individual characteristics


Ad. 1: Plant: growth habit



Ad. 2: Plant: height



Ad. 5: Petiole: length
Ad. 6: Leaf blade: length
Ad. 7: Leaf blade: width




Ad. 9: Leaf blade: incisions of margin







Ad. 11: Inflorescence: axillary flower




Ad. 12: Calyx: number of lobes




Ad. 13: Flower: length in front view Ad. 14: Flower: width in front view



Ad. 15: Corolla tube: length
Ad. 16: Corolla tube: color of outer side



Ad. 17: Corolla tube: vertical lines of inner side on corolla



Ad. 18: Corolla tube: color of inner side at basal part


Ad. 21: Upper corolla lobe: color of basal part

Ad. 22: Upper corolla lobe: color of distal part


Ad. 23: Lateral corolla lobe: color of central part

Ad. 24: Lateral corolla lobe: color of marginal part


Ad. 25: Lower corolla lobe: color of distal part



Ad. 19: Corolla lobe: incision of margin



Ad. 20: Upper corolla lobe: undulation



9. <u>Literature</u>

Inoue, Y, Kobayashi, A., etc., 1982: Encyclopedia of horticulture. Seibundoshinkosya. Tokyo, JP, vol.11, pp.161 to 162

Tsukamoto, Y., 1994: The grand dictionary of Horticulture. Shogakukan. Tokyo, JP, pp.1638 to 1639

10. <u>Technical Questionnaire</u>

| TECHNICAL QUESTION | NAIRE | Page {x} of {y} | Reference Number: |
|--------------------------|------------|--|--|
| | | | Application date: (not to be filled in by the applicant) |
| to be completed | | INICAL QUESTIONN tion with an applicatio | JAIRE n for plant breeders' rights |
| 1. Subject of the Techni | cal Questi | ionnaire (please compl | ete): |
| 1.1 Botanical name | To | renia L. | |
| 1.2 Common name | To | renia, Bluewings, Wisl | hbone-flower |
| | | | |
| 2. Applicant | | | |
| Name | | | |
| Address | | | |
| | | | |
| | | | |
| Telephone No. | | | |
| Fax No. | | | |
| E-mail address | | | |
| Breeder (if different f | rom appli | cant) | |
| · · | | , | |
| 3. Proposed denomination | on and bre | eeder's reference | |
| Proposed denomination | | | |
| (if available) | лі | | |
| Breeder's reference | | | |

| TECHNICAL QUESTIONNAIRE | | Reference Number: |
|-------------------------|------------------------|-------------------|
| | I Page Jy Lot Jy L | Reference Nilmber |
| TECHNICAL OULS HONNAINE | I I ago i A (OI i v (| Reference Number. |
| | | |

| 4.1 Breed | ling scheme | | |
|-----------|--|-----------------------------------|---|
| Varie | ty resulting from: | | |
| 4.1.1 | Crossing | | |
| | (a) controlled cross (please state parent varie | [] | |
| (|) | x (|) |
| | female parent | male parent | |
| | (b) partially known cross (please state known pare | ent variety(ies)) | |
| (|) | х (|) |
| | female parent | male parent | |
| | (c) unknown cross | [] | |
| 4.1.2 | Mutation (please state parent variety) | [] | |
| 4.1.3 | Discovery and development (please state where and when | [] discovered and how developed) | |
| 4.1.4 | Other (please provide details) | [] | |
| | | | |
| | (T) | | |

[#] Authorities may allow certain of this information to be provided in a confidential section of the Technical Questionnaire.

| TECHNICAL QUESTIONNAIRE | Page $\{x\}$ of $\{y\}$ | Reference Number: |
|-------------------------|-------------------------|-------------------|
| | | |

| (a | Vegetative propagation) cuttings | [] |
|-------|-----------------------------------|-----|
| (b | in vitro propagation | [] |
| (c | other (state method) | [] |
| 4.2.2 | Seed | [] |
| 4.2.2 | Seed | [] |
| | Other (please provide details) | [] |

| TECHNICAL QUESTIONNAIRE | Page $\{x\}$ of $\{y\}$ | Reference Number: |
|-------------------------|-------------------------|-------------------|

In the case of hybrid varieties the production scheme for the hybrid should be provided on a separate sheet. This should provide details of all the parent lines required for propagating the hybrid e.g. Single Hybrid female parent male parent Three-Way Hybrid (.....) (.....) female line male line (....<u>.....</u>) X single hybrid used as female parent male parent and should identify in particular: (a) any male sterile lines maintenance system of male sterile lines. (b)

| TECHNICAL QUESTIONNAIRE | Page $\{x\}$ of $\{y\}$ | Reference Number: |
|-------------------------|-------------------------|-------------------|

5. Characteristics of the variety to be indicated (the number in brackets refers to the corresponding characteristic in Test Guidelines; please mark the note which best corresponds).

| | Characteristics | Example Varieties | Note |
|----------------|--|-------------------|------|
| 5.1 (1) | Plant: growth habit | | |
| | upright | Crown Blue | 1[] |
| | semi upright | | 2[] |
| | horizontal | Danmoon20 | 3[] |
| 5.2(i) (21) | Upper corolla lobe: color of basal part | | |
| | RHS Colour Chart (indicate reference number) | | |
| 5.2(ii) (21) | Upper corolla lobe: color of basal part | | |
| | white | | 1[] |
| | yellow | | 2[] |
| | orange | | 3[] |
| | red | | 4[] |
| | pink | | 5[] |
| | purple | | 6[] |
| | violet | | 7[] |
| | other (indicate color) | | |
| 5.3(i) (22) | Upper corolla lobe: color of distal part | | |
| | RHS Colour Chart (indicate reference number) | | |

TECHNICAL QUESTIONNAIRE Page {x} of {y} Reference Number:

| | Characteristics | Example Varieties | Note |
|-----------------|--|-------------------|------|
| 5.3(ii) (22) | Upper corolla lobe: color of distal part | | |
| | white | | 1[] |
| | yellow | | 2[] |
| | orange | | 3[] |
| | red | | 4[] |
| | pink | | 5[] |
| | purple | | 6[] |
| | violet | | 7[] |
| | other (indicate color) | | |
| 5.4(i) (23) | Lateral corolla lobe: color of central part | | |
| | RHS Colour Chart (indicate reference number) | | |
| 5.4(ii) (23) | Lateral corolla lobe: color of central part | | |
| | white | | 1[] |
| | yellow | | 2[] |
| | orange | | 3[] |
| | red | | 4[] |
| | pink | | 5[] |
| | purple | | 6[] |
| | violet | | 7[] |
| | other (indicate color) | | |
| 5.5 (27) | Lower corolla lobe: color of blotch | | |
| | white | | 1[] |
| | yellow | | 2[] |

| TECHNICAL QUESTI | ONNAIRE I | Page {x} c | of {y} | Reference Nu | mber: | |
|---|-----------------------------|-------------------|--|--------------|--|--|
| 6. Similar varieties and differences from these varieties Please use the following table and box for comments to provide information on how your candidate variety differs from the variety (or varieties) which, to the best of your knowledge, is (or are) most similar. This information may help the examination authority to conduct its examination of distinctness in a more efficient way. | | | | | | |
| Denomination(s) of variety(ies) similar to your candidate variety Characteristics which your can variety differs fr similar variety | | andidate from the | Describe the expression of the characteristic(s) for the similar variety(ies) | | Describe the expression of the characteristic(s) for your candidate variety | |
| Example | Example Plant: height short | | hort | medium | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| Comments: | | | | | | |
| | | | | | | |
| | | | | | | |

| TEC | HNICAL QUESTIONNAIRE Page {x} of {y} Reference Number: | | | | |
|-----------------|---|--|--|--|--|
| [#] 7. | Additional information which may help in the examination of the variety | | | | |
| 7.1 | In addition to the information provided in sections 5 and 6, are there any additional characteristics which may help to distinguish the variety? | | | | |
| | Yes [] No [] | | | | |
| | (If yes, please provide details) | | | | |
| 7.2 | Are there any special conditions for growing the variety or conducting the examination? | | | | |
| | Yes [] No [] | | | | |
| | (If yes, please provide details) | | | | |
| 7.3 | Other information | | | | |
| A rep | A representative color image of the variety should accompany the Technical Questionnaire. | | | | |
| 8. | Authorization for release | | | | |
| | (a) Does the variety require prior authorization for release under legislation concerning the protection of the environment, human and animal health? | | | | |
| | Yes [] No [] | | | | |
| | (b) Has such authorization been obtained? | | | | |
| | | | | | |

No

If the answer to (b) is yes, please attach a copy of the authorization.

[]

Yes

[]

[#] Authorities may allow certain of this information to be provided in a confidential section of the Technical Questionnaire.

| IEC. | HNIC. | AL QUESTIONNAIRE Page {x} of {y} | Reference N | umber: | | | | |
|---|---|--|-------------|---------|--------|--|--|--|
| | | | | | | | | |
| 9. | Information on plant material to be examined or submitted for examination. | | | | | | | |
| effec | O.1 The expression of a characteristic or several characteristics of a variety may be affected by factors, such as pests and disease, chemical treatment (e.g. growth retardants or pesticides), effects of tissue culture, different rootstocks, scions taken from different growth phases of a cree, etc. | | | | | | | |
| 9.2 The plant material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If the plant material has undergone such treatment, full details of the treatment must be given. In this respect, please indicate below, to the best of your knowledge, if the plant material to be examined has been subjected to: | | | | | | | | |
| | (a) | Microorganisms (e.g. virus, bacteria, phytoplasi | ma) | Yes [] | No [] | | | |
| | (b) | Chemical treatment (e.g. growth retardant, pesti | icide) | Yes [] | No [] | | | |
| | (c) | Tissue culture | | Yes [] | No [] | | | |
| | (d) | Other factors | | Yes [] | No [] | | | |
| | Please provide details for where you have indicated "yes". | | | | | | | |
| | | | •••• | | | | | |
| 9.3 Has the plant material to be examined been tested for the presence of virus or other pathogens? | | | | | | | | |
| | Yes [] (please provide details as specified by the Authority) | | | | | | | |
| | No | [] | | | | | | |
| 10. I hereby declare that, to the best of my knowledge, the information provided in this form is correct: | | | | | | | | |
| | Appl | icant's name | | | | | | |
| | Signa | ature | Date | | | | | |
| | | | | | | | | |

[End of document]