

TG/29/8(proj.2)
ORIGINAL: English
DATE: 2017-07-28

INTERNATIONAL UNION FOR THE PROTECTION OF NEW VARIETIES OF PLANTS

Geneva

DRAFT

ALSTROEMERIA

UPOV Code(s): ALSTR

Alstroemeria L.

GUIDELINES

FOR THE CONDUCT OF TESTS

FOR DISTINCTNESS, UNIFORMITY AND STABILITY

prepared by experts from the Netherlands to be considered by the Technical Working Party for Ornamental Plants and Forest Trees at its fiftieth session, to be held in Victoria, British Columbia, Canada from 2017-09-11 to 2017-09-15

Disclaimer: this document does not represent UPOV policies or guidance

Alternative names:*

| Botanical name | English | French | German | Spanish | |
|-----------------|-------------------------|-------------------------------|-----------|-------------|--|
| Alstroemeria L. | Alstroemeria, Herb Lily | Alstroemère, Lis des Incas | Inkalilie | Alstromeria | |

The purpose of these guidelines ("Test Guidelines") is to elaborate the principles contained in the General Introduction (document TG/1/3), and its associated TGP documents, into detailed practical guidance for the harmonized examination of distinctness, uniformity and stability (DUS) and, in particular, to identify appropriate characteristics for the examination of DUS and production of harmonized variety descriptions.

ASSOCIATED DOCUMENTS

These Test Guidelines should be read in conjunction with the General Introduction and its associated TGP documents.

^{*} These names were correct at the time of the introduction of these Test Guidelines but may be revised or updated. [Readers are advised to consult the UPOV Code, which can be found on the UPOV Website (www.upov.int), for the latest information.]

| - | TABLE C | OF CONTENTS | PAGE |
|---|---------------------------------|---|--|
| | I. SUBJ | ECT OF THESE TEST GUIDELINES | <u>3</u> |
| 2 | 2. MATE | ERIAL REQUIRED | . <u>3</u> |
| 3 | B. METH | HOD OF EXAMINATION | . <u>4</u> |
| | 3.1 3.2 | Number of Growing CyclesTesting Place | |
| | 3.3 | Conditions for Conducting the Examination | . <u>4</u> |
| | 3.4 3.5 | Test DesignAdditional Tests | . <u>4</u> |
| 4 | 1. ASSE | SSMENT OF DISTINCTNESS, UNIFORMITY AND STABILITY | . <u>5</u> |
| | 4.1 4.2 4.3 | Distinctness | <u>6</u> |
| į | 5. GROI | JPING OF VARIETIES AND ORGANIZATION OF THE GROWING TRIAL | . <u>7</u> |
| (| 6. INTRO | ODUCTION TO THE TABLE OF CHARACTERISTICS | . <u>8</u> |
| | 6.1 6.2 6.3 6.4 6.5 | Categories of Characteristics States of Expression and Corresponding Notes Types of Expression Example Varieties Legend | . <u>8</u> . <u>8</u> . <u>8</u> |
| 7 | | E OF CHARACTERISTICS/TABLEAU DES CARACTÈRES/MERKMALSTABELLE/TABLA DE ACTERES | <u>10</u> |
| 8 | 3. EXPL | ANATIONS ON THE TABLE OF CHARACTERISTICS | <u>22</u> |
| | 8.1 8.2 | Explanations covering several characteristics Explanations for individual characteristics | . <u>22</u> . <u>24</u> |
| (| 9. LITER | RATURE | .35 |

10. TECHNICAL QUESTIONNAIRE.....36

1. <u>Subject of these Test Guidelines</u>

These Test Guidelines apply to all varieties of Alstroemeria L.

2. Material Required

- 2.1 The competent authorities decide on the quantity and quality of the plant material required for testing the variety and when and where it is to be delivered. Applicants submitting material from a State other than that in which the testing takes place must ensure that all customs formalities and phytosanitary requirements are complied with.
- 2.2 The material is to be supplied in the form of young plants.
- 2.3 The minimum quantity of plant material, to be supplied by the applicant, should be:

8 plants

- 2.4 The plant material supplied should be visibly healthy, not lacking in vigor, nor affected by any important pest or disease.
- 2.5 The plant material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If it has been treated, full details of the treatment must be given.

3. Method of Examination

3.1 Number of Growing Cycles

The minimum duration of tests should normally be a single growing cycle.

3.2 Testing Place

Tests are normally conducted at one place. In the case of tests conducted at more than one place, guidance is provided in TGP/9 "Examining Distinctness".

- 3.3 Conditions for Conducting the Examination
- 3.3.1 The tests should be carried out under conditions ensuring satisfactory growth for the expression of the relevant characteristics of the variety and for the conduct of the examination.
- 3.3.2 Because daylight varies, color determinations made against a color chart should be made either in a suitable cabinet providing artificial daylight or in the middle of the day in a room without direct sunlight. The spectral distribution of the illuminant for artificial daylight should conform with the CIE Standard of Preferred Daylight D 6500 and should fall within the tolerances set out in the British Standard 950, Part I. These determinations should be made with the plant part placed against a white background. The color chart and version used should be specified in the variety description.
- 3.4 Test Design

Each test should be designed to result in a total of at least 8 plants.

3.5 Additional Tests

Additional tests, for examining relevant characteristics, may be established.

4. Assessment of Distinctness, Uniformity and Stability

4.1 Distinctness

4.1.1 General Recommendations

It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding distinctness. However, the following points are provided for elaboration or emphasis in these Test Guidelines.

4.1.2 Consistent Differences

The differences observed between varieties may be so clear that more than one growing cycle is not necessary. In addition, in some circumstances, the influence of the environment is not such that more than a single growing cycle is required to provide assurance that the differences observed between varieties are sufficiently consistent. One means of ensuring that a difference in a characteristic, observed in a growing trial, is sufficiently consistent is to examine the characteristic in at least two independent growing cycles.

4.1.3 Clear Differences

Determining whether a difference between two varieties is clear depends on many factors, and should consider, in particular, the type of expression of the characteristic being examined, i.e. whether it is expressed in a qualitative, quantitative, or pseudo-qualitative manner. Therefore, it is important that users of these Test Guidelines are familiar with the recommendations contained in the General Introduction prior to making decisions regarding distinctness.

4.1.4 Number of plants or parts of plants to be Examined

Unless otherwise indicated, for the purposes of distinctness, all observations on single plants should be made on 7 plants or parts of plants taken from each of 7 plants and any other observations made on all plants in the test, disregarding any off-type plants.

4.1.5 Method of Observation

The recommended method of observing the characteristic for the purposes of distinctness is indicated by the following key in the second column of the Table of Characteristics (see document TGP/9 "Examining Distinctness", Section 4 "Observation of characteristics"):

MG: single measurement of a group of plants or parts of plants

MS: measurement of a number of individual plants or parts of plants

VG: visual assessment by a single observation of a group of plants or parts of plants

VS: visual assessment by observation of individual plants or parts of plants

Type of observation: visual (V) or measurement (M)

"Visual" observation (V) is an observation made on the basis of the expert's judgment. For the purposes of this document, "visual" observation refers to the sensory observations of the experts and, therefore, also includes smell, taste and touch. Visual observation includes observations where the expert uses reference points (e.g. diagrams, example varieties, side-by-side comparison) or nonlinear charts (e.g. color charts). Measurement (M) is an objective observation against a calibrated, linear scale e.g. using a ruler, weighing scales, colorimeter, dates, counts, etc.

Type of record: for a group of plants (G) or for single, individual plants (S)

For the purposes of distinctness, observations may be recorded as a single record for a group of plants or parts of plants (G), or may be recorded as records for a number of single, individual plants or parts of plants (S). In most cases, "G" provides a single record per variety and it is not possible or necessary to apply statistical methods in a plant-by-plant analysis for the assessment of distinctness.

In cases where more than one method of observing the characteristic is indicated in the Table of Characteristics (e.g. VG/MG), guidance on selecting an appropriate method is provided in document TGP/9, Section 4.2.

- 4.2 Uniformity
- 4.2.1 It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding uniformity. However, the following points are provided for elaboration or emphasis in these Test Guidelines:
- 4.2.2 For the assessment of uniformity of vegetatively propagated varieties, a population standard of 1% and an acceptance probability of at least 95% should be applied. In the case of a sample size of 8 plants, 1 off-type is allowed.
- 4.3 Stability
- 4.3.1 In practice, it is not usual to perform tests of stability that produce results as certain as those of the testing of distinctness and uniformity. However, experience has demonstrated that, for many types of variety, when a variety has been shown to be uniform, it can also be considered to be stable.
- 4.3.2 Where appropriate, or in cases of doubt, stability may be further examined by testing a new plant stock to ensure that it exhibits the same characteristics as those shown by the initial material supplied.
- 5. <u>Grouping of Varieties and Organization of the Growing Trial</u>
- 5.1 The selection of varieties of common knowledge to be grown in the trial with the candidate varieties and the way in which these varieties are divided into groups to facilitate the assessment of distinctness are aided by the use of grouping characteristics.
- 5.2 Grouping characteristics are those in which the documented states of expression, even where produced at different locations, can be used, either individually or in combination with other such characteristics: (a) to select varieties of common knowledge that can be excluded from the growing trial used for examination of distinctness; and (b) to organize the growing trial so that similar varieties are grouped together.
- 5.3 The following have been agreed as useful grouping characteristics:
 - (a) Plant: height (characteristic 1)
 - (b) Leaf blade: number of colors on inner side (silvery colored stripe excluded) (characteristic 8)
 - (c) Flower: main color (characteristic 13)
- 5.4 Guidance for the use of grouping characteristics, in the process of examining distinctness, is provided through the General Introduction and document TGP/9 "Examining Distinctness".

6. Introduction to the Table of Characteristics

6.1 Categories of Characteristics

6.1.1 Standard Test Guidelines Characteristics

Standard Test Guidelines characteristics are those which are approved by UPOV for examination of DUS and from which members of the Union can select those suitable for their particular circumstances.

6.1.2 Asterisked Characteristics

Asterisked characteristics (denoted by *) are those included in the Test Guidelines which are important for the international harmonization of variety descriptions and should always be examined for DUS and included in the variety description by all members of the Union, except when the state of expression of a preceding characteristic or regional environmental conditions render this inappropriate.

6.2 States of Expression and Corresponding Notes

- 6.2.1 States of expression are given for each characteristic to define the characteristic and to harmonize descriptions. Each state of expression is allocated a corresponding numerical note for ease of recording of data and for the production and exchange of the description.
- 6.2.2 In the case of qualitative and pseudo-qualitative characteristics (see Chapter 6.3), all relevant states of expression are presented in the characteristic. However, in the case of quantitative characteristics with 5 or more states, an abbreviated scale may be used to minimize the size of the Table of Characteristics. For example, in the case of a quantitative characteristic with 9 states, the presentation of states of expression in the Test Guidelines may be abbreviated as follows:

| State | Note |
|--------|------|
| small | 3 |
| medium | 5 |
| large | 7 |

However, it should be noted that all of the following 9 states of expression exist to describe varieties and should be used as appropriate:

| State | Note |
|---------------------|------|
| very small | 1 |
| very small to small | 2 |
| small | 3 |
| small to medium | 4 |
| medium | 5 |
| medium to large | 6 |
| large | 7 |
| large to very large | 8 |
| very large | 9 |

6.2.3 Further explanation of the presentation of states of expression and notes is provided in document TGP/7 "Development of Test Guidelines".

6.3 Types of Expression

An explanation of the types of expression of characteristics (qualitative, quantitative and pseudo-qualitative) is provided in the General Introduction.

6.4 Example Varieties

Where appropriate, example varieties are provided to clarify the states of expression of each characteristic.

6.5 Legend

| | English | | français | | deutsch | español | Example Varieties Exemples Beispielssorten Variedades ejemplo | Note/ Nota | | |
|-----|------------------------|----------------------|--------------------------|--------------|----------------------------------|--------------------------------------|--|---------------|--|--|
| 1 2 | 3 | 4 | 5 | 6 | 7 | | | | | |
| | Name chara in En | cteristics | Nom o carac frança | tère en | Name des Merkmals auf Deutsch | Nombre del carácter en español | | | | |
| | | states of expression | | d'expression | Ausprägungsstufen | tipos de expresión | | | | |

1 Characteristic number

2 (*) Asterisked characteristic – see Chapter 6.1.2

3 Type of expression

QL Qualitative characteristic – see Chapter 6.3
QN Quantitative characteristic – see Chapter 6.3
PQ Pseudo-qualitative characteristic – see Chapter 6.3

4 Method of observation (and type of plot, if applicable)
MG, MS, VG, VS - see Chapter 4.1.5

5 (+) See Explanations on the Table of Characteristics in Chapter 8.2

6 (a)-(d) See Explanations on the Table of Characteristics in Chapter 8.1

7 Not applicable

7. <u>Table of Characteristics/Tableau des caractères/Merkmalstabelle/Tabla de caracteres</u>

| | | | English | | français | deutsch | español | Example Varieties Exemples Beispielssorten Variedades ejemplo | Note/ Nota |
|----|-----|----------------------|-----------------------------------|-----|----------|---------|---------|--|---------------|
| 1. | (*) | QN | MG/MS/VG | (+) | (a) | | | | |
| | | Plant: | height | | | | | | |
| | | short | | | | | | | 3 |
| | | mediu | m | | | | | | 5 |
| | | tall | | | | | | | 7 |
| 2. | (*) | QN | MG/MS/VG | (+) | (a) | | | | |
| | | Stem: | thickness | | | | | | |
| | | thin | | | | | | | 3 |
| | | mediu | m | | | | | | 5 |
| | | thick | | | | | | | 7 |
| 3. | | QN | VG | (+) | (a) | | | , | 1 |
| | | Stem: colora | anthocyanin ition | | | | | | |
| | | absent | t or very weak | | | | | | 1 |
| | | weak | | | | | | | 3 |
| | | mediu | m | | | | | | 5 |
| | | strong | , | | , | | | | 7 |
| 4. | | QN | VG | | (a) | | | | |
| | | Stem: anthocolora | distribution of cyanin tion | | | | | | |
| | | at base | e only | | | | | | 1 |
| | | basal l | nalf only | | | | | | 2 |
| | | basal a | and apical part | | | | | | 3 |
| | | whole | stem | | | | | | 4 |

| | | English | | français | deutsch | español | Example Varieties Exemples Beispielssorten Variedades ejemplo | Note/ Nota |
|----|------------|--|-----|----------|---------|---------|--|---------------|
| 5. | QN | MG/MS/VG | | (a), (b) | | | | |
| • | Leaf: | Leaf: length | | • | | | | |
| | short | | | | | | | 3 |
| | mediu | um | | | | | | 5 |
| | long | | | | | | | 7 |
| 6. | QN | MG/MS/VG | | (a), (b) | | 1 | | |
| | Leaf: | width | | • | | | | |
| | narro | w | | | | | | 3 |
| | mediu | | | | | | | 5 |
| | broad | | | | | | | 7 |
| 7. | QN | MS/VG | (+) | (a), (b) | | | | - |
| : | Leaf: | attitude | | • | | | | |
| | semi-erect | | | | | | | 3 |
| | | horizontal | | | | | | 5 |
| | semi- | drooping | | | | | | 7 |
| 8. | QL | VG | (+) | (a), (b) | | | | • |
| · | color | blade: number of s on inner side erry colored stripe ded) | | • | | | | |
| | one | | | | | | | 1 |
| | two | | | | | | | 2 |
| | more | than two | | | | | | 3 |
| 9. | QL | VG | (+) | (a), (b) | | | 1 | |
| : | Leaf color | blade: silvery ed longitudinal | | | | | | |
| | abser | nt | | | | | | 1 |
| | prese | ent | | | | | | 9 |

| | | English | | français | deutsch | español | Example Varieties Exemples Beispielssorten Variedades ejemplo | Note/ Nota |
|---------|---------------------------|-------------------|-----|---------------|---------|---------|--|---------------|
| 10. (*) | QN | MG/MS/VG | (+) | (a) | | | | |
| | Umbe | I: length of ray | | | | | | |
| | short | | | | | | | 3 |
| | mediu | m | | | | | | 5 |
| | long | | | | | | | 7 |
| 11. (*) | <u> </u> | MG/MS/VG | | (a) | | | | |
| 117 | | I: number of rays | | 1 ' ' | | | | |
| | | | | | | | | |
| | few | | | | | | | 3 |
| | medium | | | | | | | 5 |
| 12. (*) | many | MG/MS/VG | (1) | (a), (c) | | | | 7 |
| 12. | | | (+) | (a), (c) | | | | T |
| | Flower: length of pedicel | | | | | | | |
| | short | | | | | | | 3 |
| | medium | | | | | | | 5 |
| | long | | | | | | | 7 |
| 13. (*) | PQ | VG | | (a), (c), (d) | | | | |
| : | Flowe | r: main color | | • | | | | |
| | | | | | | | | |
| | white . | | | | | | | 1 |
| | | sh yellow | | | | | | 2 |
| | light ye | | | | | | | 3 |
| | | m yellow | | | | | | 4 |
| | orange | | | | | | | 5 |
| | light pi | | | | | | | 6 |
| | | m pink | | | | | | 7 |
| | purple | | | | | | | 8 |
| | orange | e red | | | | | | 9 |
| | red | | | | | | | 10 |
| | purple | | | | | | | 11 |
| | light p | | | | | | | 12 |
| | mediu | m purple | | | | | | 13 |

English **Example Varieties** Note/ français deutsch español Exemples Nota Beispielssorten Variedades ejemplo 14. QN MG/MS/VG (+) (a), (c) Flower: length in frontal view short 3 medium 5 7 long 15. MG/MS/VG QN (+) (a), (c) Flower: width in frontal view narrow 3 5 medium broad 7 16. QN MG/MS/VG (a), (c) (+) Flower: ratio length/width low 3 5 medium 7 high 17. QN MG/MS/VG (+) (a), (c) Flower: length in side view 3 short medium 5 7 long 18. (*) PQ ۷G (+) (a), (c) Outer tepal: shape of blade moderately elliptic 1 broad elliptic 2 round 3 moderately obovate 4 broad obovate 5

| | | English | | français | deutsch | español | Example Varieties Exemples Beispielssorten Variedades ejemplo | Note/ Nota |
|---------|--|--|-----|---------------|---------|---------|---|---------------|
| 19. | QN | VG | (+) | (a), (c) | | | | • |
| | Outer tepal: emargination | | | | | | | |
| | shallo | shallow | | | | | | 3 |
| | mediu | medium | | | | | | 5 |
| | deep | | | | | | | 7 |
| 20. (*) | PQ | VG | | (a), (c), (d) | | | 1 | |
| | Outer | tepal: main of outer side | | | | | | |
| | | Colour Chart ate reference er) | | | | | | |
| 21. (*) | QN | VG | (+) | (a), (c) | | | | |
| | Outer tepal: area of green color of outer side | | | | | | | |
| | abser | nt or very small | | | | | | 1 |
| | small | | | | | | | 2 |
| | mediu | ım | | | | | | 3 |
| | large | | | | | | | 4 |
| 22. (*) | PQ | VG | | (a), (c), (d) | | | | |
| | color | r tepal: main of central zone ner side | | | | | | |
| | RHS (indication | Colour Chart ate reference er) | | | | | | |
| 23. (*) | PQ | VG | | (a), (c), (d) | | | | |
| | color | Outer tepal: main color of top zone of inner side (green tip excluded) | | | | | | |
| | RHS Colour Chart (indicate reference number) | | | | | | | |

| | | English | | français | deutsch | español | Example Varieties Exemples Beispielssorten Variedades ejemplo | Note/ Nota |
|---------|--------|--|-------|---------------|---------|---------|--|---------------|
| 24. (*) | PQ | VG | | (a), (c), (d) | | | | 1 |
| | color | tepal: main of lateral zone er side | | | | | | |
| | | Colour Chart ate reference er) | | | | | | |
| 25. (*) | PQ | VG | | (a), (c), (d) | | , | | 1 |
| | color | tepal: main of basal zone er side | | _ | | | | |
| | | Colour Chart ate reference er) | | | | | | |
| 26. (*) | QN | VG | (+) | (a), (c) | | • | • | |
| | stripe | tepal: small es on marginal of lateral zone of side | | | | | | |
| | absen | nt or very few | ••••• | | | | | 1 |
| | few | | | | | | | 3 |
| | mediu | ım | | | | | | 5 |
| | many | | | | | | | 7 |
| 27. (*) | QN | VG | (+) | (a), (c) | | | | |
| | inner | large stripes on side (marginal excluded) | | | | | | |
| | absen | ıt | | | | | | 1 |
| | few | | | | | | | 2 |
| | mediu | ım | | | | | | 3 |
| | many | | | | | | | 4 |

| | | English | | français | deutsch | español | Example Varieties Exemples Beispielssorten | Note/ Nota |
|---------|---|--|-----|---------------|---------|----------|--|---------------|
| | | | | | | | Variedades ejemplo | |
| 28. (*) | PQ | VG | (+) | (a), (c) | | | | |
| | Inner | tepal: shape | | | | | | |
| | narrov | w elliptic | • | | | | | 1 |
| | mode | rately elliptic | | | | | | 2 |
| | narrov | w obovate | | | | | | 3 |
| | mode | rately obovate | | | | | | 4 |
| 29. (*) | PQ | VG | | (a), (c), (d) | | | | |
| | Inner lateral tepal: main color of central zone of inner side | | | | | | | |
| | RHS ((indica numb | Colour Chart ate reference er) | | | | | | |
| 30. (*) | PQ | VG | | (a), (c), (d) | | | | |
| | main zone | Inner lateral tepal: main color of apical zone of inner side (tip excluded) | | | | | | |
| | | Colour Chart ate reference er) | | | | | | |
| 31. (*) | PQ | VG | | (a), (c), (d) | | <u> </u> | | - |
| | main | lateral tepal: color of basal of inner side | | | | | | |
| | | Colour Chart ate reference er) | | | | | | |

| | | English | | français | deutsch | español | Example Varieties Exemples Beispielssorten Variedades ejemplo | Note/ Nota |
|---------|--|--|-----|----------|---------|---------|--|---------------|
| 32. (*) | QN | MG/VG | (+) | (a), (c) | | | | |
| | numb | Inner lateral tepal: number of stripes on inner side | | | | | | |
| | abser | absent or very few | | | | | | 1 |
| | few | | | | | | | 3 |
| | mediu | ım | | | | | | 5 |
| | many | | | | | | | 7 |
| | very r | many | | | | | | 9 |
| 33. (*) | QN | VG | (+) | (a), (c) | | | | • |
| | tepal | lateral : area of striped on inner side | | | | | | |
| | small | | | | | | | 3 |
| | medium | | | | | | | 5 |
| | large | | | | | | | 7 |
| 34. (*) | QN | MG/MS/VG | (+) | (a), (c) | | • | | |
| | lengt stripe | lateral tepal: h of longest es on inner side e on central vein ided) | | | | | | |
| | short | | | | | | | 3 |
| | mediu | ım | | | | | | 5 |
| | long | | | | | | | 7 |
| 35. (*) | QN | MG/VG | (+) | (a), (c) | | | | |
| | Inner lateral tepal: width of widest stripes on inner side (stripe on central vein excluded) | | | | | | | |
| | narro | w | | | | | | 3 |
| | mediu | ım | | | | | | 5 |
| | broad | l | | | | | | 7 |


| | | | | | | | T | 1 |
|---------|--------------------------|---|-----|---------------|---------|---------|--|---------------|
| | | English | | français | deutsch | español | Example Varieties Exemples Beispielssorten Variedades ejemplo | Note/ Nota |
| 36. (*) | PQ | VG | | (a), (c), (d) | | • | | |
| | Inner tepal: inner | median main color of side | | | | | | |
| | | Colour Chart ate reference er) | • | | | | | |
| 37. (*) | PQ | VG | | (a), (c), (d) | | | | |
| | tepal: | median secondary color er side | | | | | | |
| | | Colour Chart ate reference er) | | | | | | |
| 38. (*) | QN | MG/VG | | (a), (c) | | | | • |
| | Inner numb inner | median tepal: er of stripes on side | | | | | | |
| | absen | absent or very few | | | | | | 1 |
| | few | | | | | | | 3 |
| | mediu | ım | | | | | | 5 |
| | many | many | | | | | | 7 |
| 39. (*) | PQ | VG | (+) | (a), (c) | | | | |
| | Anthe | er: color | | | | | | |
| | green | ish | | | | | | 1 |
| | yellow | | | | | | | 2 |
| | orang | | | | | | | 3 |
| | purplis | | | | | | | 4 |
| | brown | | | | | | | 5 |
| | blue | | | | | | | 6 |
| | mediu | ım grey | | | | | | 7 |
| | dark g | grey | | | | | | 8 |


| | | English | | français | deutsch | español | Example Varieties Exemples Beispielssorten Variedades ejemplo | Note/ Nota |
|---------|-----------------------|--|----------|---------------|---------|---------|--|---------------|
| 40. (*) | PQ | VG | | (a), (c), (d) | | | | |
| | Filam | ent: main color | | | | | | |
| | white | | | | | | | 1 |
| | yellow | ······································ | | | | | | 2 |
| | orang | e | <u> </u> | | | | | 3 |
| | orang | e red | | | | | | 4 |
| | red | | | | | | | 5 |
| | pink | | | | | | | 6 |
| | red pu | ırple | | | | | | 7 |
| | light p | ourple | | | | | | 8 |
| | mediu | ım purple | | | | | | 9 |
| 41. (*) | QL | VG | (+) | (a), (c) | | | | |
| | Filament: small spots | | | | | | | |
| | abser | nt | | | | | | 1 |
| | few | | | | | | | 2 |
| | mediu | ım | | | | | | 3 |
| | many | | | | | | | 4 |
| 42. (*) | QL | VG | (+) | (a), (c) | | | | |
| | Stign | na: spots | | | | | | |
| | abser | nt | | | | | | 1 |
| | prese | nt | | | | | | 9 |
| 43. (*) | QN | VG | (+) | (a), (c) | | 1 | | • |
| | Ovary | y: anthocyanin ation | | | | | | |
| | abser | nt or very weak | · | | | | | 1 |
| | weak | | 1 | | | | | 3 |
| | mediu | ım | | | | | | 5 |
| | strong |) | | | | | | 7 |
| | very s | strong | Ī | | | | | 9 |

- 8. <u>Explanations on the Table of Characteristics</u>
- 8.1 Explanations covering several characteristics

Characteristics containing the following key in the second column of the Table of Characteristics should be examined as indicated below:

- (a) Unless otherwise indicated, all characteristics should be observed on the first full developed stem when 50% of the flowers are open.
- (b) Observations on the leaves should be done on leaves taken from the middle third of the stem.
- (c) Observations on the flower should be made at the time of dehiscence of the first anther in an individual flower.





(d) The main color is the color with the largest surface area. In cases where the areas of the main and secondary color are too similar to reliably decide which color has the largest surface area, the darkest color is considered to be the main color.

8.2 Explanations for individual characteristics

Ad. 1: Plant: height

Plant height should be observed from soil level to the top of the plant, including the flowers.


Ad. 2: Stem: thickness

The thickness of the stem should be measured at the middle third of the stem.

Ad. 3: Stem: anthocyanin coloration



Ad. 7: Leaf: attitude



Ad. 8: Leaf blade: number of colors on inner side (silvery colored stripe excluded)









Ad. 9: Leaf blade: silvery colored longitudinal stripe





Ad. 10: Umbel: length of ray



Ad. 12: Flower: length of pedicel



Ad. 14: Flower: length in frontal view



Ad. 15: Flower: width in frontal view



Ad. 16: Flower: ratio length/width



Ad. 17: Flower: length in side view




Ad. 18: Outer tepal: shape of blade

| | ← broadest part → | | | | | |
|------------------------|-------------------|-----------------|--------------------|--|--|--|
| | below middle | at middle | above middle | | | |
| | | 1 | 4 | | | |
| | | medium elliptic | moderately obovate | | | |
| broad ← width → narrow | | 2 | 5 | | | |
| | | broad elliptic | broad obovate | | | |
| | | 3 round | | | | |


Ad. 19: Outer tepal: emargination



Ad. 21: Outer tepal: area of green color of outer side




Ad. 26: Outer tepal: small stripes on marginal part of lateral zone of inner side



Ad. 27: Outer tepal: large stripes on inner side (marginal zone excluded)



Ad. 28: Inner tepal: shape



Ad. 32: Inner lateral tepal: number of stripes on inner side




Ad. 33: Inner lateral tepal: area of striped zone on inner side



Ad. 34: Inner lateral tepal: length of longest stripes on inner side (stripe on central vein excluded)




Ad. 35: Inner lateral tepal: width of widest stripes on inner side (stripe on central vein excluded)




Ad. 39: Anther: color

To be observed just before dehiscence.

Ad. 41: Filament: small spots




Ad. 42: Stigma: spots



TG/29/8(proj.2) Alstroemeria, 2017-07-28 30

Ad. 43: Ovary: anthocyanin coloration



9. <u>Literature</u>

Grunert, C, 1980: Das Blumenzwiebelbuch. Verlag Eugen Ulmer. Stuttgart, DE, x pp.

The Royal General Bulbgrowers' Association, 1991: International Checklist for Hyacinths and Miscellaneous Bulbs. Koninklijke Algemeene Vereeniging voor Bloembollencultuur. Hillegom, NL, pp. 15 to 47

10. <u>Technical Questionnaire</u>

| TECHNICAL QUESTIONNAIRE | | | | Page {x} of {y} | Reference Number: | | |
|-------------------------|--|-----------------------------|------|---|--|--|--|
| | | | | | | | |
| | | | | | Application date: (not to be filled in by the applicant) | | |
| | | | | CHNICAL QUESTIONNAtection with an application | AIRE n for plant breeders' rights | | |
| 1. | Subject of the Technical Questionnaire | | | | | | |
| | 1.1 | Botanical name | Als | stroemeria L. | | | |
| | 1.2 | Common name | Als | stroemeria, Herb Lily | | | |
| | | | | | | | |
| 2. | Applica | nt | | | | | |
| | Name | | | | | | |
| | Addres | S | | | | | |
| | Telepho | one No. | | | | | |
| | Fax No | | | | | | |
| | E-mail | address | | | | | |
| | Breede applica | r (if different from nt) | | | | | |
| 3. | Propos | ed denomination and bree | eder | 's reference | | | |
| | Propos (if avail | ed denomination able) | | | | | |
| | Breede | r's reference | | | | | |

TECHNICAL QUESTIONNAIRE Page {x} of {y} Reference Number: #4. Information on the breeding scheme and propagation of the variety 4.1 Breeding scheme Variety resulting from: 4.1.1 Crossing (a) controlled cross [] (please state parent varieties) (.....) (.....) female parent male parent (b) partially known cross [] (please state known parent variety(ies)) (.....) (.....) female parent male parent (c) unknown cross [] 4.1.2 [] Mutation (please state parent variety) [] 4.1.3 Discovery and development (please state where and when discovered and how developed) [] Other 4.1.4 (please provide details)

| TECHNICAL Q | UESTIONNAIRE | Page {x} of {y} | Reference Number | : |
|-------------|-----------------------------------|-----------------|------------------|------------|
| | | | | |
| 4.2 | Method of propagating the | variety | | |
| 4.2.1 | Vegetative propagation | | | |
| (a) (b) | Rhizomes Other (state method) | | | [] [] |
| | | | | |
| 4.2.2 | Other (Please provide details) | | | [] |
| | | | | |
| | | | | |

TECHNICAL QUESTIONNAIRE Page {x} of {y} Reference Number:

5. Characteristics of the variety to be indicated (the number in brackets refers to the corresponding characteristic in Test Guidelines; please mark the note which best corresponds).

| | Characteristics | Example Varieties | Note |
|-------------|---|-------------------|--------|
| 5.1 (1) | Plant: height | | |
| | short | | 3[] |
| | medium | | 5[] |
| | tall | | 7[] |
| 5.2 (8) | Leaf blade: number of colors on inner side (silvery colore stripe excluded) | ed | |
| | one | | 1[] |
| | two | | 2[] |
| | more than two | | 3[] |
| 5.3 (13) | Flower: main color | | |
| | white | | 1[] |
| | greenish yellow | | 2[] |
| | light yellow | | 3[] |
| | medium yellow | | 4[] |
| | orange | | 5[] |
| | light pink | | 6[] |
| | medium pink | | 7[] |
| | purple pink | | 8[] |
| | orange red | | 9[] |
| | red | | 10[] |
| | purple red | | 11 [] |
| | light purple | | 12[] |
| | medium purple | | 13 [] |
| | dark purple | | 14 [] |

| TECHNICAL QUESTIONN | NAIRE | Page {x} of | {y} | Reference Nu | ımber: | |
|--|---|----------------------------|---------------------------------|--|--|------------------------|
| 6. Similar varieties and of Please use the following tal from the variety (or varieties help the examination author) | ble and box for o | comments to post of your P | provide inform knowledge, is | (or are) most | similar. This info | |
| Denomination(s) of variety(ies) similar to your candidate variety | Characteristic your candidate from the simila | variety differs | the characte | expression of ristic(s) for the variety(ies) | Describe the e the characteris candidate | tic(s) for your |
| Example | Plant: h | eight | sl | hort | medi | ium |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| Comments: | | | | | | |

| TECHNICAL QUESTIONNAIRE | Page {x} of {y} | Reference Number: |
|-------------------------|-----------------|-------------------|

| #7. | Addition | nal information which may he | elp in the examination of the | ne variety | | | | |
|-----|---|---|-------------------------------|---|--|--|--|--|
| 7.1 | In addition to the information provided in sections 5 and 6, are there any additional characteristics which make help to distinguish the variety? | | | | | | | |
| | Yes | [] | No | [] | | | | |
| | (If yes, | please provide details) | | | | | | |
| 7.2 | Are the | Are there any special conditions for growing the variety or conducting the examination? | | | | | | |
| | Yes | [] | No | [] | | | | |
| | (If yes, | please provide details) | | | | | | |
| 7.3 | Other i | nformation | | | | | | |
| | | | | stinguishing feature(s), should accompany the tion of the candidate variety which | | | | |

supplements the information provided in the Technical Questionnaire. The key points to consider when taking a photograph of the candidate variety are:

- Indication of the date and geographic location
- Correct labeling (breeder's reference)
 Good quality printed photograph (minimum 10 cm x 15 cm) and/or sufficient resolution electronic format version (minimum 960 x 1280 pixels)"

Further guidance on providing photographs with the Technical Questionnaire is available in document TGP/7 "Development of Test Guidelines", Guidance Note 35 (http://www.upov.int/tgp/en/).
[The link provided may be deleted by members of the Union when developing authorities' own test guidelines.]

| TECH | INICA | L QUEST | TONNAIRE | Page {x} of | {y} | Reference | e Numb | er: | | |
|----------------|-------------------|---|---|-------------------------------------|---------------------------------|-------------------------------|-----------------------|----------|----------|------------|
| 8. | Autho | rization for | release | | | | | | | |
| | (a) | Does the variety require prior authorization for release under legislation concerning the protection of the environment, human and animal health? | | | | | | | | |
| | | Yes | [] | No | [] | | | | | |
| | (b) | Has such | authorization bee | n obtained? | | | | | | |
| | | Yes | [] | No | [] | | | | | |
| | If the | answer to | (b) is yes, please a | ttach a copy of the | he authorizati | ion. | | | | |
| 9. Info | ormatio | on on plant | t material to be exa | mined or submit | ted for exami | nation | | | | |
| | and o | disease, cl | on of a characterist hemical treatment en from different gr | (e.g. growth ret | tardants or p | | | | | |
| chara has u | cterist ndergo | ics of the vone such t | al should not havariety, unless the reatment, full detailedge, if the plant m | competent authors is of the treatme | orities allow on the must be gi | r request so iven. In this | uch treat respect, | ment. If | the plan | t material |
| | (a) | Micro | oorganisms (e.g. vi | rus, bacteria, ph | ytoplasma) | | Yes [|] | No [|] |
| | (b) | Cher | mical treatment (e. | g. growth retarda | nt, pesticide) | | Yes [|] | No [|] |
| | (c) | Tissu | ue culture | | | | Yes [|] | No [|] |
| | (d) | Othe | er factors | | | | Yes [|] | No [|] |
| | Plea | ase provid | e details for where | you have indicat | ed "yes". | | | | | |
| | | | | | | | | | | |
| 10. | I he | reby decla | re that, to the best | of my knowledge | e, the informa | ation provide | ed in this | form is | correct: | |
| | App | licant's na | me | | | | | | | |
| | Signature | | | | | | | | | |

[End of document]