



TG/49/8(proj.1)
ORIGINAL: English
DATE: 2006-05-18

INTERNATIONAL UNION FOR THE PROTECTION OF NEW VARIETIES OF PLANTS
 GENEVA

DRAFT

CARROT

UPOV Code: DAUCU_CAR

Daucus carota L.

*

GUIDELINES

FOR THE CONDUCT OF TESTS

FOR DISTINCTNESS, UNIFORMITY AND STABILITY

prepared by experts from France

*to be considered by the Technical Working Party for Vegetables (TWP)
 at its fortieth session to be held in Guanajuato, Guanajuato State, Mexico,
 from June 12 to 16, 2006*

Alternative Names:^{*}

<i>Botanical name</i>	<i>English</i>	<i>French</i>	<i>German</i>	<i>Spanish</i>
<i>Daucus carota L.</i>	Carrot	Carotte	Möhre	Zanahoria

The purpose of these guidelines ("Test Guidelines") is to elaborate the principles contained in the General Introduction (document TG/1/3), and its associated TGP documents, into detailed practical guidance for the harmonized examination of distinctness, uniformity and stability (DUS) and, in particular, to identify appropriate characteristics for the examination of DUS and production of harmonized variety descriptions.

ASSOCIATED DOCUMENTS

These Test Guidelines should be read in conjunction with the General Introduction and its associated TGP documents.

* These names were correct at the time of the introduction of these Test Guidelines but may be revised or updated. [Readers are advised to consult the UPOV Code, which can be found on the UPOV Website (www.upov.int), for the latest information.]

<u>TABLE OF CONTENTS</u>	<u>PAGE</u>
1. SUBJECT OF THESE TEST GUIDELINES.....	3
2. MATERIAL REQUIRED	3
3. METHOD OF EXAMINATION.....	3
3.1 Duration of Tests.....	3
3.2 Testing Place.....	3
3.3 Conditions for Conducting the Examination.....	3
3.4 Test Design	4
3.5 Number of Plants / Parts of Plants to be Examined.....	4
3.6 Additional Tests	4
4. ASSESSMENT OF DISTINCTNESS, UNIFORMITY AND STABILITY.....	4
4.1 Distinctness	4
4.2 Uniformity.....	5
4.3 Stability	5
5. GROUPING OF VARIETIES AND ORGANIZATION OF THE GROWING TRIAL.....	5
6. INTRODUCTION TO THE TABLE OF CHARACTERISTICS	6
6.1 Categories of Characteristics.....	6
6.2 States of Expression and Corresponding Notes.....	6
6.3 Types of Expression.....	6
6.4 Example Varieties	7
6.5 Legend.....	7
7. TABLE OF CHARACTERISTICS/TABLEAU DES CARACTÈRES/MERKMALSTABELLE/TABLA DE CARACTERES.....	8
8. EXPLANATIONS ON THE TABLE OF CHARACTERISTICS	18
8.1 Explanations for covering several characteristics	18
8.2 Explanations for individual characteristics	18
9. LITERATURE.....	23
10. TECHNICAL QUESTIONNAIRE.....	24

1. Subject of these Test Guidelines

These Test Guidelines apply to all varieties of *Daucus carota* L.

2. Material Required

2.1 The competent authorities decide on the quantity and quality of the plant material required for testing the variety and when and where it is to be delivered. Applicants submitting material from a State other than that in which the testing takes place must ensure that all customs formalities and phytosanitary requirements are complied with.

2.2 The material is to be supplied in the form of seed.

2.3 The minimum quantity of plant material, to be supplied by the applicant, should be:

25 g or 30,000 seeds. [FR propose 90 000 or 120 g]

2.4 The seed should meet the minimum requirements for germination, species and analytical purity, health and moisture content, specified by the competent authority.

2.5 The plant material supplied should be visibly healthy, not lacking in vigor, nor affected by any important pest or disease.

2.6 The plant material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If it has been treated, full details of the treatment must be given.

3. Method of Examination

3.1 *Duration of Tests*

The minimum duration of tests should normally be two independent growing cycles.

3.2 *Testing Place*

Tests are normally conducted at one place. In the case of tests conducted at more than one place, guidance is provided in TGP/9 "Examining Distinctness".

3.3 *Conditions for Conducting the Examination*

The tests should be carried out under conditions ensuring satisfactory growth for the expression of the relevant characteristics of the variety and for the conduct of the examination.

3.3.1 Type of observation

The recommended method of observing the characteristic is indicated by the following key in the second column of the Table of Characteristics:

MG: single measurement of a group of plants or parts of plants
MS: measurement of a number of individual plants or parts of plants
VG: visual assessment by a single observation of a group of plants or parts of plants
VS: visual assessment by observation of individual plants or parts of plants”

3.4 *Test Design*

3.4.1 Each test should be designed to result in a total of at least 200 plants, which should be divided between two or more replicates.

3.4.2 The design of the tests should be such that plants or parts of plants may be removed for measurement or counting without prejudice to the observations which must be made up to the end of the growing cycle.

3.5 *Number of Plants / Parts of Plants to be Examined*

Unless otherwise indicated, all observations should be made on 60 plants or parts taken from each of 60 plants and any other observations made on all plants in the test. [FR propose to reduce to 40 plants.]

3.6 *Additional Tests*

Additional tests, for examining relevant characteristics, may be established.

4. Assessment of Distinctness, Uniformity and Stability

4.1 *Distinctness*

4.1.1 General Recommendations

It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding distinctness. However, the following points are provided for elaboration or emphasis in these Test Guidelines.

4.1.2 Consistent Differences

The differences observed between varieties may be so clear that more than one growing cycle is not necessary. In addition, in some circumstances, the influence of the environment is not such that more than a single growing cycle is required to provide assurance that the differences observed between varieties are sufficiently consistent. One means of ensuring that a difference in a characteristic, observed in a growing trial, is sufficiently consistent is to examine the characteristic in at least two independent growing cycles.

4.1.3 Clear Differences

Determining whether a difference between two varieties is clear depends on many factors, and should consider, in particular, the type of expression of the characteristic being examined, i.e. whether it is expressed in a qualitative, quantitative, or pseudo-qualitative manner. Therefore, it is important that users of these Test Guidelines are familiar with the

recommendations contained in the General Introduction prior to making decisions regarding distinctness.

4.2 *Uniformity*

It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding uniformity. However, the following points are provided for elaboration or emphasis in these Test Guidelines:

4.2.1 Cross-pollinated varieties

The assessment of uniformity for cross-pollinated varieties should be according to the recommendations for cross-pollinated varieties in the General Introduction.

4.2.2 Single cross hybrids and inbred lines

For the assessment of uniformity of single cross hybrids and inbred lines, a population standard of 2% and an acceptance probability of at least 95% should be applied. In the case of a sample size of 200 plants, 7 off-types are allowed.

4.2.3 Hybrids

The assessment of uniformity for hybrid varieties depends on the type of hybrid and should be according to the recommendations for hybrid varieties in the General Introduction. In the case of single cross hybrids, the uniformity standards are set out in Section 4.2.2.

4.3 *Stability*

4.3.1 In practice, it is not usual to perform tests of stability that produce results as certain as those of the testing of distinctness and uniformity. However, experience has demonstrated that, for many types of variety, when a variety has been shown to be uniform, it can also be considered to be stable.

4.3.2 Where appropriate, or in cases of doubt, stability may be tested, either by growing a further generation, or by testing a new seed or plant stock to ensure that it exhibits the same characteristics as those shown by the previous material supplied.

4.3.3 Where appropriate, or in cases of doubt, the stability of a hybrid variety may, in addition to an examination of the hybrid variety itself, also be assessed by examination of the uniformity and stability of its parent lines

5. Grouping of Varieties and Organization of the Growing Trial

5.1 The selection of varieties of common knowledge to be grown in the trial with the candidate varieties and the way in which these varieties are divided into groups to facilitate the assessment of distinctness are aided by the use of grouping characteristics.

5.2 Grouping characteristics are those in which the documented states of expression, even where produced at different locations, can be used, either individually or in combination with other such characteristics: (a) to select varieties of common knowledge that can be excluded

from the growing trial used for examination of distinctness; and (b) to organize the growing trial so that similar varieties are grouped together.

5.3 The following have been agreed as useful grouping characteristics:

- (a) Leaf: length (including petiole) (characteristic 3)
- (b) Root: length (characteristic 7)
- (c) Root: width (characteristic 8)
- (d) Root: shape in longitudinal section (characteristic 10)
- (e) Root: tip (when fully developed) (characteristic 12)
- (f) Root: external color (characteristic 13)
- (g) Root: time of coloration of tip in longitudinal section (characteristic 28) – [FR propose to delete]
[FR propose to add male stérile plants (characteristic 31) and Type of male sterility (characteristic 32).]

5.4 Guidance for the use of grouping characteristics, in the process of examining distinctness, is provided through the General Introduction.

6. Introduction to the Table of Characteristics

6.1 *Categories of Characteristics*

6.1.1 Standard Test Guidelines Characteristics

Standard Test Guidelines characteristics are those which are approved by UPOV for examination of DUS and from which members of the Union can select those suitable for their particular circumstances.

6.1.2 Asterisked Characteristics

Asterisked characteristics (denoted by *) are those included in the Test Guidelines which are important for the international harmonization of variety descriptions and should always be examined for DUS and included in the variety description by all members of the Union, except when the state of expression of a preceding characteristic or regional environmental conditions render this inappropriate.

6.2 *States of Expression and Corresponding Notes*

States of expression are given for each characteristic to define the characteristic and to harmonize descriptions. Each state of expression is allocated a corresponding numerical note for ease of recording of data and for the production and exchange of the description.

6.3 *Types of Expression*

An explanation of the types of expression of characteristics (qualitative, quantitative and pseudo-qualitative) is provided in the General Introduction.

6.4 Example Varieties

Where appropriate, example varieties are provided to clarify the states of expression of each characteristic.

6.5 Legend

(*) Asterisked characteristic – see Chapter 6.1.2

QL: Qualitative characteristic – see Chapter 6.3

QN: Quantitative characteristic – see Chapter 6.3

PQ: Pseudo-qualitative characteristic – see Chapter 6.3

MG: single measurement of a group of plants or parts of plants – see Chapter 3.3.1

MS: measurement of a number of individual plants or parts of plants – see Chapter 3.3.1

VG: visual assessment by a single observation of a group of plants or parts of plants – Chapter 3.3.1

VS: visual assessment by observation of individual plants or parts of plants” –see Chapter 3.3.1

(a)-(c) See Explanations on the Table of Characteristics in Chapter 8.1

(+) See Explanations on the Table of Characteristics in Chapter 8.2

7. Table of Characteristics/Tableau des caractères/Merkmalstabelle/Tabla de caracteresticas

		English	français	deutsch	español	Example Varieties/ Exemples/ Beispielssorten/ Variedades ejemplos	Note/ Nota
1.	VG (+)	Foliage: width of crown	Feuillage: largeur de la couronne	Laub: Breite des Blattansatzes	Follaje: anchura del cuello		
QN	(a)	narrow	étroite	schmal	estrecho	Amsterdam 2	3
		medium	moyenne	mittel	medio	Nantaise améliorée 2, Rothild	5
		broad	large	breit	ancho	Chantenay à cœur rouge 2	7
2.	VG	Leaf: attitude	Feuille: port	Blatt: Stellung	Hoja: porte		
QN	(a)	erect	dressé	aufrecht	erecto	Touchon	1
		semi-erect	demi-dressé	halbaufrecht	semierecto	Nantaise améliorée 2	3
		prostrate	étalé	liegend	postrado		5
3.	VG/ MS (*)	Leaf: length (including petiole)	Feuille: longueur (pétiole compris)	Blatt: Länge (einschließlich Stiel)	Hoja: longitud (incluido el peciolo)		
QN	(a)	very short	très courte	sehr kurz	muy corta	Mokum, Mignon	1
		short	courte	kurz	corta	Amsterdam 2, Amsterdam 3	3
		medium	moyenne	mittel	media	Juwarot, Nantaise améliorée 2	5
		long	longue	lang	larga	Chantenay, Chantenay à cœur rouge 2	7
		very long	très longue	sehr lang	muy larga	De Colmar à cœur rouge 2, Rothild	9
4.	VG (*)	Leaf: division	Feuille: division	Blatt: Fiederung	Hoja: división		
QN	(a)	fine	fine	fein	fina	Amsterdam 2, Amsterdam 3	3
		medium	moyenne	mittel	mediana	Nantaise améliorée 2, Nantaise améliorée 3	5
		coarse	grossière	grob	grosera	Hytop	7

					Example Varieties/ Exemples/ Beispielssorten/ Variedades ejemplo	Note/ Nota
	English	français	deutsch	español		
5. (*)	VG	Leaf: intensity of green color	Feuille: intensité de la couleur verte	Blatt: Intensität der Grünfärbung	Hoja: intensidad del color verde	
QN	(a)	light	claire	hell	claro	Adelaide, Leonor 3
		medium	moyenne	mittel	medio	Amsterdam 2, Amsterdam 3 5
		dark	foncée	dunkel	oscuro	Rothild 7
6. (*)	VG	Leaf: anthocyanin coloration of petiole	Feuille: pigmentation anthocyanique du pétiole	Blatt: Anthocyan-färbung des Blattstiels	Hoja: pigmentación antociánica del pecíolo	
QL	(a)	absent	absente	fehlend	ausente	Amsterdam 2 1
		present	présente	vorhanden	presente	Tareno 9
7. (*)	VG/ MS	Root: length	Racine: longueur	Rübe: Länge	Raíz: longitud	
QN	(b)	very short	très courte	sehr kurz	muy corta	Parijse Markt 2, Parijse Markt 3 1
		short	courte	kurz	corta	Chantenay 3
		medium	moyenne	mittel	media	Nantaise améliorée 2, Nantaise améliorée 3 5
		long	longue	lang	larga	Berlikumer 2, Berlikumer 3 7
		very long	très longue	sehr lang	muy larga	Lange Stompe Winter 9
8. (*)	VG/ MS	Root: width	Racine: largeur	Rübe: Breite	Raíz: anchura	
QN	(b)	narrow	étroite	schmal	estrecha	Amsterdam 2, Amsterdam 3 3
		medium	moyenne	mittel	media	Nantaise améliorée 2, Nantaise améliorée 3 5
		broad	large	breit	ancha	De Colmar à cœur rouge 2, Parijse Markt 2, Parijse Markt 3 7

		English	français	deutsch	español	Example Varieties/ Exemples/ Beispielssorten/ Variedades ejemplo	Note/ Nota
9. <small>(*)</small>	VG/ MS	Root: ratio length/ width	Racine: rapport longueur/largeur	Rübe: Verhältnis Länge/Breite	Raíz: relación longitud y anchura		
QN	(b)	very small	très petit	sehr klein	muy pequeña	Parijse Markt 2, Parijse Markt 3, Parmex	1
		small	petit	klein	pequeña	Courte améliorée à forcer	3
		medium	moyen	mittel	media	Chantenay	5
		large	grand	groß	grande	Nantaise améliorée 2, Nantaise améliorée 3	7
		very large	très grand	sehr groß	muy grande	Amsterdam 2	9
10. <small>(*) (+)</small>	VG	Root: shape in longitudinal section	Racine: forme en section longitudinale	Rübe: Form im Längsschnitt	Raíz: forma en sección longitudinal		
PQ	(b)	circular	arrondie	kreisförmig	circular	Parijse Markt 2, Parijse Markt 3	1
		obovate	obovale	verkehrt eiförmig	oboval		2
		obtriangular	obtriangulaire	verkehrt dreieckig	obtriangular	Chantenay, De Colmar à cœur rouge 2	3
		narrow obtriangular	obtriangulaire étroite	schmal verkehrt dreieckig	obtriangular estrecha	Imperator, De Colmar à cœur rouge 3	4
		narrow obtriangular to narrow oblong	obtriangulaire étroite à rectangulaire étroite	schmal verkehrt dreieckig bis schmal rechteckig	obtriangular estrecha a oblonga estrecha	Maestro	5
		narrow oblong	rectangulaire étroite	schmal rechteckig	oblonga estrecha	Amsterdam 2, Berlikumer 2, Berlikumer 3, Nantaise améliorée 5, Touchon	6
11. <small>(*) (+)</small>	VG	Root: shape of shoulder	Racine: forme de l'épaulement	Rübe: Form des Kopfes	Raíz: forma del hombro		
PQ	(b)	flat	plat	flach	plana	De Colmar à cœur rouge 2	1
		flat to rounded	plat à arrondi	flach bis abgerundet	plana a redondeada	Parijse Markt 2	2
		rounded	arrondi	abgerundet	redondeada		3
		rounded to conical	arrondi à conique	abgerundet bis kegelförmig	redondeada a cónica		4
		conical	conique	kegelförmig	cónica	Touchon	5

					Example Varieties/ Exemples/ Beispielssorten/ Variedades ejemplo	Note/ Nota
English		français	deutsch	español		
12. (*)	VG	Root: tip (when fully developed)	Racine: extrémité (à plein développement)	Rübe: Ende (bei voller Entwicklung)	Raíz: extremo (en pleno desarrollo)	
PQ	(b)	blunt	arrondie	stumpf	romo	Berlikumer 3
		slightly pointed	légèrement pointue	leicht spitz	ligeramente puntiagudo	Mello Yello
		strongly pointed	fortement pointue	sehr spitz	muy puntiagudo	Allred, Orbit
13. (*)	VG	Root: external color	Racine: couleur externe	Rübe: äußere Farbe	Raíz: color externo	
PQ	(b)	white	blanche	weiß	blanco	White Satin
		yellow	jaune	gelb	amarillo	Mello Yello
		orange	orange	orange	naranja	Bingo, Tancar, Goliath, Karotan, Pinocchio
		pinkish red	rouge rosâtre	rosarot	rojo rosado	Nutri-red
		red	rouge	rot	rojo	Pulsor
		purple	pourpre	purpurn	púrpura	Purple Haze
14.	VG	Root: intensity of external color	Racine: intensité de la couleur externe	Rübe: Intensität der äußeren Farbe	Raíz: intensidad el color externo	
		[For non white varieties]				
QN	(b)	light	claire	hell	claro	Mello Yello, Bingo, Tancar
		medium	moyenne	mittel	medio	Nutri-red, Goliath
		dark	foncée	dunkel	oscuro	Purple Haze, Karotan, Pinocchio
15.	VG	Root: anthocyanin coloration of skin of shoulder	Racine: pigmentation anthocyanique de la peau du collet	Rübe: Anthocyanfärbung der Haut des Kopfes	Raíz: pigmentación antociánica de la epidermis del hombro	
QL	(b)	absent	absente	fehlend	ausente	Trevor
		present	présente	vorhanden	presente	Touchon

					Example Varieties/ Exemples/ Beispielssorten/ Variedades ejemplo	Note/ Nota
16.	VG (*) (+)	Root: extent of green color of skin of shoulder	Racine: extension de la coloration verte de la peau du collet	Rübe: Ausdehnung der Grünfärbung der Haut des Kopfes	Raíz: extensión del color verde de la epidermis del hombro	
		[FR propose to deletes *]				
QN	(b)	absent or very small small medium large very large	nulle ou très petite petite moyenne grande très grande	fehlend oder sehr gering gering mittel groß sehr groß	ausente o muy pequeño pequeño medio grande muy grande	Karotan Scarla De Colmar à cœur rouge 2 Touchon Lange Stompe Winter
17.	VG	Root: ridging of surface	Racine: annelure de la surface	Rübe: Ringelung der Oberfläche	Raíz: anillado de la superficie	
QN	(b)	absent or very weak weak medium strong very strong	absente ou très faible faible moyenne forte très forte	fehlend oder sehr gering gering mittel stark sehr stark	ausente o muy débil débil medio fuerte muy fuerte	Favor, Sytan Major Chantenay De Colmar à cœur rouge 2 9
18.	VG (*)	Root: diameter of core relative to total diameter	Racine: diamètre du cœur par rapport au diamètre total	Rübe: Durchmesser des Herzens im Verhältnis zum gesamten Durchmesser	Raíz: diámetro del corazón en relación con el diámetro total	
QN	(b)	very small small medium large very large	très petit petit moyen grand très grand	sehr klein klein mittel groß sehr groß	muy pequeño pequeño medio grande muy grande	Amsterdam 2, Amsterdam 3, Tourino Nantaise améliorée 2, Nantaise améliorée 3 Berlikumer 2, Berlikumer 3 De Colmar à cœur rouge 2 Giganta

		English	français	deutsch	español	Example Varieties/ Exemples/ Beispielssorten/ Variedades ejemplo	Note/ Nota
19. (*)	VG	Root: color of core	Racine: couleur du cœur	Rübe: Farbe des Herzens	Raíz: color del corazón		
PQ	(b)	white	blanc	weiß	blanco	White Satin	1
		yellow	jaune	gelb	amarillo	Jaune de Lobberich, Pariser Markt	2
		orange	orange	orange	naranja	Nantaise améliorée 2, Nantaise améliorée 3	3
		pinkish red	rouge rosâtre	rosarot	rojo rosado	Nutri-red	4
		red	rouge	rot	rojo		5
20.	VG	Root: intensity of color of core	Racine: intensité de la couleur du cœur	Rübe: Intensität der Farbe des Herzens	Raíz: intensidad del color del corazón		
		[For non white varieties]					
QN	(b)	light	claire	hell	claro		3
		medium	moyenne	mittel	medio		5
		dark	foncée	dunkel	oscuro		7
21. (*)	VG	Root: color of cortex	Racine: couleur du cortex	Rübe: Farbe der Rinde	Raíz: color de la corteza		
PQ	(b)	white	blanc	weiß	blanco	White Satin	1
		yellow	jaune	gelb	amarillo	Mellow Yello	2
		orange	orange	orange	naranja	Allred, Carlo	3
		pinkish red	rouge rosâtre	rosarot	rojo rosado	Nutri-red	4
		red	rouge	rot	rojo		5
22.	VG	Root: intensity of color of cortex	Racine: intensité de la couleur du cortex	Rübe: Intensität der Farbe der Rinde	Raíz: intensidad del color de la corteza		
		[For non white varieties]					
QN	(b)	light	claire	hell	claro		3
		medium	moyenne	mittel	medio		5
		dark	foncée	dunkel	oscuro		7

					Example Varieties/ Exemples/ Beispielssorten/ Variedades ejemplo	Note/ Nota
English	français	deutsch	español			
23. VG	Root: color of core compared to color of cortex	Racine: couleur du cœur par rapport à la couleur du cortex	Rübe: Farbe des Herzens im Verhältnis zur Rinde	Raíz: color del corazón en relación con la corteza		
QN (b)	lighter	plus claire	heller	más claro		1
	same	même couleur	gleichfarbig	mismo color		2
	darker	plus foncée	dunkler	más oscuro		3
24. VG (*)	Root: extent of green coloration of interior (in longitudinal section)	Racine: extension de la coloration verte à l'intérieur (en section longitudinale)	Rübe: Ausdehnung der Grünfärbung im Inneren (im Längsschnitt)	Raíz: extensión del color verde del interior (en sección longitudinal)		
QN (b)	absent or very small	nulle ou très petite	fehlend oder sehr klein	ausente o muy pequeña	Major	1
	small	petite	klein	pequeño	Meaux	3
	medium	moyenne	mittel	medio	Chantenay à cœur rouge 2, De Colmar à cœur rouge 3	5
	large	grande	groß	grande	Touchon	7
	very large	très grande	sehr groß	muy grande	Muscade	9
25. VG	Root: protrusion above soil	Racine: partie hors terre	Rübe: Sitz über dem Boden	Raíz: parte fuera del suelo		
QN (b)	absent or very slight	nulle ou très faible	fehlend oder sehr gering	ausente o muy pequeña	Karotan, Parijse Markt 3	1
	slight	faible	gering	pequeña	Amsterdam 2, Amsterdam 3, Nantaise améliorée 2, Nantaise améliorée 3	3
	medium	moyenne	mittel	media	Tancar, Toudo	5
	much	importante	hoch	grande	Lange Stompe Winter, Touchon	7
	very much	très importante	sehr hoch	muy grande	Blanche à collet vert hors terre	9

	English	français	deutsch	español	Example Varieties/ Exemples/ Beispielssorten/ Variedades ejemplo	Note/ Nota
26.	MS	Root: weight	Racine: poids	Rübe: Gewicht	Raíz: peso	
		[This characteristic is deleted from CPVO protocole]				
QN	(b)	small	faible	gering	pequeño	Amsterdam 2 3
		medium	moyen	mittel	medio	Nantaise améliorée 2, Nantaise améliorée 3 5
		high	élevé	hoch	grande	Giganta 7
27.	VG	Varieties with blunt tip only: Root: time of development of rounded tip	Variétés avec extrémité arrondie seulement: Racine: époque de boutage	Nur Sorten mit stumpfem Ende: Rübe: Zeitpunkt der Bildung eines runden Endes	Solo variedades con extremo romo: Raíz: época de desarrollo del extremo redondeado	
(+)						
QN	(b)	early	précoce	früh	temprana	Touchon 3
		medium	moyenne	mittel	media	Tiana, Nantaise améliorée 2, Nantaise améliorée 3 5
		late	tardive	spät	tardía	Bureau, Tancar, Nantaise améliorée 7 7
28.	VG	Root: time of coloration of tip in longitudinal section	Racine: époque de coloration de l'extrémité en section longitudinale	Rübe: Zeitpunkt der Färbung der Spitze im Längsschnitt	Raíz: época de coloración del extremo en sección longitudinal	
(*)						
(+)						
		[FR propose to delete *]				
QN	(b)	very early	très précoce	sehr früh	muy temprana	Parise Markt 3 1
		early	précoce	früh	temprana	Amsterdam 2, Amsterdam 3 3
		medium	moyenne	mittel	media	Nantaise améliorée 2, Nantaise améliorée 3 5
		late	tardive	spät	tardía	De Colmar à cœur rouge 2, Touchon 7
		very late	très tardive	sehr spät	muy tardía	Goliath 9

		English	français	deutsch	español	Example Varieties/ Exemples/ Beispielssorten/ Variedades ejemplo	Note/ Nota
29.	VG	Plant: tendency to bolting	Plante: tendance à la montaison	Pflanze: Neigung zum Schossen	Planta: tendencia a la subida a flor		
		weak	faible	gering	débil	Molene, Tancar	3
		medium	moyenne	mittel	media	Nantaise améliorée 2, Nantaise améliorée 3	5
		strong	forte	stark	fuerte	Touchon	7
30.	VG	Plant: height of primary umbel at time of its flowering	Plante: hauteur de l'ombelle primaire à l'époque de sa floraison	Pflanze: Höhe der ersten Dolde zum Zeitpunkt seiner Blüte	Planta: altura de la umbela primaria en la época de floración		
		short	basse	niedrig	baja		3
		medium	moyenne	mittel	media		5
		tall	haute	hoch	alta		7
31.	VS	Plants: proportion of male sterile plants	Plantes: proportion de plantes mâles stériles	Pflanzen: Anteil männlich steriler Pflanzen	Plantas: proporción de plantas androestériles		
		[FR propose to add *]					
		absent or very low	nulle ou très faible	fehlend oder sehr gering	ausente o muy baja	Nantaise améliorée 2, Touchon	1
		low	faible	gering	baja		3
		medium	moyenne	mittel	media	Nanco, Tino	5
		high	forte	hoch	alta	Nandor, Tancar	7
		very high	très forte	sehr hoch	muy alta		9

		English	français	deutsch	español	Example Varieties/ Exemples/ Beispielssorten/ Variedades ejemplo	Note/ Nota
32.	VS	Plant: type of male sterility	Plante: type de stérilité mâle	Pflanze: Typ der männlichen Sterilität	Planta: tipo de línea androestéril		
		[FR propose to add *]					
QL	brown anther	anthère brune	braune Antheren	antera: marrón	Nanco	1	
	petaloid anther	anthère pétaloïde	petaloide Antheren	antera petaloide	Tino	2	
33.	(+)	[FR propose to add a new characteristic, without *]					
		Resistance to <i>Alternaria dauci</i>					
		absent			Presto	1	
		intermediate			Tino	2	
		present			Bolero	3	

8. Explanations on the Table of Characteristics


8.1 *Explanations for covering several characteristics*

Characteristics containing the following key in the second column of the Table of Characteristics should be examined as indicated below:



- (a) Foliage and leaf: All observations on the foliage and the leaf should be made at the time of full development of the foliage.
- (b) Root: All observations on the root should be made when the root is fully developed.

8.2 *Explanations for individual characteristics*

Ad. 1: Foliage: width of crown



Ad. 10: Root: shape in longitudinal section



1
circular

2
obovate

3
obtriangular



4
narrow obtriangular




5
narrow obtriangular to
narrow oblong




6
narrow oblong

Ad. 11: Root: shape of shoulder



Ad. 16: Root: extent of green color of skin of shoulder



Ad. 27: Varieties with blunt tip only: Root: time of development of rounded tip

Ad. 28: Root: time of coloration of tip in longitudinal section

The earliness of carrot varieties can be judged according to two criteria, characteristic 27, time of development of "rounded tip" for the varieties with a blunt tip at maturity and characteristic 28, time of coloration of the tip in longitudinal section.

Three weeks before the normal maturity date of the varieties (where the variety 'Touchon' has a blunt tip): pull up of part of the test roots in order to judge the shape of the tip, characteristic 27 (early: blunt tip: variety 'Touchon', medium: varieties 'Tiana', 'Nantaise améliorée 2', 'Nantaise améliorée 3', late: pointed tip: varieties 'Bureau', 'Tancar', 'Nantaise améliorée 7').

Following longitudinal cutting of the roots: examination of the coloration of the tip, characteristic 28 (early: colored tip: varieties 'Amsterdam 2', 'Amsterdam 3', late: whitish tip: varieties 'De Colmar à coeur rouge 2', 'Touchon').

A good example is the variety 'Touchon' which is early for characteristic 27 and late for characteristic 28.

Ad. 33: Resistance to *Alternaria dauci*

Resistance to leaf blight (*Alternaria dauci* (Kühn) Groves & Skolko)

Method

Maintenance of strains

Strains: no differences in virulence have been described for *Alternaria dauci*, so one aggressive isolates or a mix of isolates can be used. For example 7 isolates (185, 187, 188, 189, 190, 191, 193) can be used

:

growth conditions: on V8 plates, 7 days in darkness, at 20°C ; afterward, 12 h UV- 12 h darkness for 7 days at 20°C to stimulate sporulation

Execution of test

Sowing: in open field, with a direct sowing (mid-May). No fungicides applied.

Specific design: randomized complete-block design, with 4 replications and 2 lines per cultivar per replication. In each replication, either a susceptible cultivar (Presto) or a less susceptible (Valor) are regularly planted in order to enable a uniform repartition of the fungus.

Temperature: summer temperature in field. Optimum temperature: 24-28 °C. Temperature above 12°C during night and a high humidity on leaves are necessary for disease development (Strandberg, 1988)

Humidity: high humidity. Irrigation by sprinkler 24 h before inoculation and 48 after inoculation; afterwards, at least twice a week.

Inoculum: A suspension of spores in water with tween is made with a mix of the isolates and the suspension is adjusted to about 10^3 to 4.10^3 spores/ml

Method of inoculation: By spraying a suspension of conidia on foliage with a sprayer

Growth stage of plants: 5-7 leaves for the first inoculation

Duration of test (2 inoculations):

- from sowing to the first inoculation: 2 months (mid-July)
- from sowing to the second inoculation: 2.5 months (end of July-beginning of August)

- from inoculation to reading (at least 2 readings):
 - * first reading : 5-7 weeks after the 2nd inoculation
 - * second reading: about 15 days after the first reading.

Standard varieties:

Susceptible: PRESTO

Intermediate resistant (moderate level): TINO

Intermediate resistant (high level): BOLERO

Quantity of plants tested: 4 replicates of 2 lines of 4.3 meters per cultivar

Scale of notes: modified Aguilar scale, enlarged from 1 to 9 (Cadot V, 2000).

- 1: **no disease**
- 2: very few lesions on lower leaves
- 3: **few lesions on lower leaves,**
- 4: few lesions on the upper leaves and lower leaves moderately attacked
- 5: **few lesions on the upper leaves and lower leaves heavily attacked**
- 6: moderate lesions on the upper leaves and lower leaves heavily attacked
- 7: **upper leaves heavily attacked and lower leaves completely blighted**
- 8 : upper leaves very heavily attacked and lower leaves completely blighted
- 9: **all leaves blighted**

Cadot Valérie, 9 mai 2006.

Bibliography :

Aguilar J.A.E. et al. 1986. Nivel de resistencia de cenoura a *Alternaria dauci* e interação com tratamento químico. Hort. Bras. 4(2), 19-22.

Cadot V. et al . 2002. Setting up a resistance test to *Alternaria dauci* of carrot by inoculation in the open field, as part of registering varieties in the French Catalogue of vegetable species. Journées J. Chevaugeon Mycologie, Aussois, mars 2002. Poster.

Cadot V. et al. 2000. Développement d'un test de résistance en plein champ : *Alternaria dauci* de la carotte. In Mise au point de tests de résistance à des maladies chez les espèces potagères. Séminaire protection sanitaire des semences, plants et variétés, 30 mars 1999, INRA d'Angers.78-88.

Strandberg J.O. 1988. Establishment of *Alternaria* leaf blight on carrots in controlled environments. Plant Disease / Vol 72 n°6, 522-526.

9. Literature

There is no specific literature.

10. Technical Questionnaire

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
		Application date: (not to be filled in by the applicant)
<p style="text-align:center">TECHNICAL QUESTIONNAIRE to be completed in connection with an application for plant breeders' rights</p> <p>In the case of hybrid varieties which are the subject of an application for plant breeders' rights, and where the parent lines are to be submitted as a part of the examination of the hybrid variety, this Technical Questionnaire should be completed for each of the parent lines, in addition to being completed for the hybrid variety.</p>		
1. Subject of the Technical Questionnaire		
1.1 Latin Name	Daucus carota L.	
1.2 Common Name	Carrot	
2. Applicant		
Name		
Address		
Telephone No.		
Fax No.		
E-mail address		
Breeder (if different from applicant)		
3. Proposed denomination and breeder's reference		
Proposed denomination (if available)		
Breeder's reference		

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
<p>4. Information on the breeding scheme and propagation of the variety</p> <p>4.1 Breeding scheme</p> <p>Variety resulting from:</p> <p>4.1.1 Crossing</p> <p>(a) controlled cross [] (please state parent varieties)</p> <p>(b) partially known cross [] (please state known parent variety(ies))</p> <p>(c) unknown cross []</p> <p>4.1.2 Mutation [] (please state parent variety)</p> <p>4.1.3 Discovery and development [] (please state where and when discovered and how developed)</p> <p>4.1.4 Other [] (please provide details)</p> <p>4.2 Method of propagating the variety</p> <p>4.2.1 Seed-propagated varieties</p> <p>(a) Self-pollination []</p> <p>(b) Cross-pollination</p> <p>(i) population []</p> <p>(ii) synthetic variety []</p> <p>(c) Hybrid []</p> <p>(d) Other [] (please provide details)</p> <p>4.2.2 Other [] (please provide details)</p>		

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
5. Characteristics of the variety to be indicated (the number in brackets refers to the corresponding characteristic in Test Guidelines; please mark the note which best corresponds).		
Characteristics	Example Varieties	Note
5.1 Leaf: length (including petiole) (3)		
very short	Mokum, Mignon	1[]
short	Amsterdam 2, Amsterdam 3	3[]
medium	Juwarot, Nantaise améliorée 2	5[]
long	Chantenay, Chantenay à cœur rouge 2	7[]
very long	De Colmar à cœur rouge 2, Rothild	9[]
5.2 Leaf: intensity of green color (5)		
light	Adelaide, Leonor	3[]
medium	Amsterdam 2, Amsterdam 3	5[]
dark	Rothild	7[]
5.3 Root: length (7)		
very short	Parijse Markt 2, Parijse Markt 3	1[]
short	Chantenay	3[]
medium	Nantaise améliorée 2, Nantaise améliorée 3	5[]
long	Berlikumer 2, Berlikumer 3	7[]
very long	Lange Stompe Winter	9[]

TECHNICAL QUESTIONNAIRE		Page {x} of {y}	Reference Number:
Characteristics		Example Varieties	Note
5.4	Root: width (8)	narrow	Amsterdam 2, Amsterdam 3 3[]
		medium	Nantaise améliorée 2, Nantaise améliorée 3 5[]
		broad	De Colmar à cœur rouge 2, Parijse Markt 2, Parijse Markt 3 7[]
5.5	Root: shape in longitudinal section (10)	circular	Parijse Markt 2, Parijse Markt 3 1[]
		obovate	2[]
		obtriangular	Chantenay, De Colmar à cœur rouge 2 3[]
		narrow obtriangular	Imperator, De Colmar à cœur rouge 3 4[]
		narrow obtriangular to narrow oblong	Maestro 5[]
		narrow oblong	Amsterdam 2, Berlikumer 2, Berlikumer 3, Nantaise améliorée 5, Touchon 6[]
5.6	Root: shape of shoulder (11)	flat	De Colmar à cœur rouge 2 1[]
		flat to rounded	Parijse Markt 2 2[]
		rounded	3[]
		rounded to conical	4[]
		conical	Touchon 5[]
5.7	Root: tip (when fully developed) (12)	blunt	Berlikumer 3 1[]
		slightly pointed	Mello Yello 2[]
		strongly pointed	Allred, Orbit 3[]

TECHNICAL QUESTIONNAIRE		Page {x} of {y}	Reference Number:
Characteristics		Example Varieties	Note
5.8	Root: external color		
(13)			
	white	White Satin	1[]
	yellow	Mello Yello	2[]
	orange	Bingo, Tancar, Goliath, Karotan, Pinocchio	3[]
	pinkish red	Nutri-red	4[]
	red	Pulsor	5[]
	purple	Purple Haze	6[]
5.9	Root: intensity of external color		
(14)			
	light	Mello Yello, Bingo, Tancar	3[]
	medium	Nutri-red, Goliath	5[]
	dark	Purple Haze, Karotan, Pinocchio	7[]
5.10	Root: color of core		
(19)			
	white	White Satin	1[]
	yellow	Jaune de Lobberich, Pariser Markt	2[]
	orange	Nantaise améliorée 2, Nantaise améliorée 3	3[]
	pinkish red	Nutri-red	4[]
	red		5[]

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:	
Characteristics	Example Varieties	Note	
5.11 Root: time of coloration of tip in longitudinal section (28) FR propose to delete			
very early	Parijse Markt 3	1[]	
early	Amsterdam 2, Amsterdam 3	3[]	
medium	Nantaise améliorée 2, Nantaise améliorée 3	5[]	
late	De Colmar à cœur rouge 2, Touchon	7[]	
very late	Goliath	9[]	
FR propose to add 2 characteristics for male sterility, 31 and 32.			
6. Similar varieties and differences from these varieties			
<i>Please use the table, and space provided for comments, below to provide information on how your candidate variety differs from the variety (or varieties) which, to the best of your knowledge, is (or are) most similar. This information may help the examination authority to conduct its examination of distinctness in a more efficient way.</i>			
Denomination(s) of variety(ies) similar to your candidate variety	Characteristic(s) in which your candidate variety differs from the similar variety(ies)	Describe the expression of the characteristic(s) for the similar variety(ies)	Describe the expression of the characteristic(s) for your candidate variety
<i>Example</i>	<i>Root: external color</i>	<i>orange</i>	<i>pinkish red</i>
Comments:			

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
<p>7. Additional information which may help in the examination of the variety</p> <p>7.1 In addition to the information provided in sections 5 and 6, are there any additional characteristics which may help to distinguish the variety?</p> <p>Yes [] No []</p> <p>(If yes, please provide details)</p> <p>7.2 Special conditions for the examination of the variety</p> <p>7.2.1 Are there any special conditions for growing the variety or conducting the examination?</p> <p>Yes [] No []</p> <p>7.2.2 If yes, please give details:</p> <p>7.3 Other information</p>		
<p>8. Authorization for release</p> <p>(a) Does the variety require prior authorization for release under legislation concerning the protection of the environment, human and animal health?</p> <p>Yes [] No []</p> <p>(b) Has such authorization been obtained?</p> <p>Yes [] No []</p> <p>If the answer to (b) is yes, please attach a copy of the authorization.</p>		

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

9. Information on plant material to be examined.

9.1 The expression of a characteristic or several characteristics of a variety may be affected by factors, such as pests and disease, chemical treatment (e.g. growth retardants or pesticides), effects of tissue culture, different rootstocks, scions taken from different growth phases of a tree, etc.

9.2 The plant material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If the plant material has undergone such treatment, full details of the treatment must be given. In this respect, please indicate below, to the best of your knowledge, if the plant material to be examined has been subjected to:

- | | | |
|---|---------|--------|
| (a) Microorganisms (e.g. virus, bacteria, phytoplasma) | Yes [] | No [] |
| (b) Chemical treatment (e.g. growth retardant or pesticide) | Yes [] | No [] |
| (c) Tissue culture | Yes [] | No [] |
| (d) Other factors | Yes [] | No [] |

Please provide details of where you have indicated “yes”:

.....

10. I hereby declare that, to the best of my knowledge, the information provided in this form is correct:

Applicant's name

Signature

Date

[End of document]