



TG/BRASS\_JUN(proj.4)

ORIGINAL: English

DATE: 2016-05-20

## INTERNATIONAL UNION FOR THE PROTECTION OF NEW VARIETIES OF PLANTS

Geneva

DRAFT

## BROWN MUSTARD

UPOV Code(s): BRASS\_JUN

*Brassica juncea* (L.) Czern.

## GUIDELINES

## FOR THE CONDUCT OF TESTS

## FOR DISTINCTNESS, UNIFORMITY AND STABILITY

*prepared by experts from Japan**to be considered by the*

*Technical Working Party for Vegetables  
at its fiftieth session, to be held in Brno, Czech Republic,  
from 2016-06-27 to 2016-07-01*

*Disclaimer: this document does not represent UPOV policies or guidance*

Alternative names:\*

<i>Botanical name</i>	<i>English</i>	<i>French</i>	<i>German</i>	<i>Spanish</i>
<i>Brassica juncea</i> (L.) Czern.	Brown mustard, India mustard, Indian mustard, Oriental mustard	Moutarde brune	Sareptasenf	Mostaza de Sarepta, Mostaza india

The purpose of these guidelines ("Test Guidelines") is to elaborate the principles contained in the General Introduction (document TG/1/3), and its associated TGP documents, into detailed practical guidance for the harmonized examination of distinctness, uniformity and stability (DUS) and, in particular, to identify appropriate characteristics for the examination of DUS and production of harmonized variety descriptions.

## ASSOCIATED DOCUMENTS

These Test Guidelines should be read in conjunction with the General Introduction and its associated TGP documents.

\* These names were correct at the time of the introduction of these Test Guidelines but may be revised or updated. [Readers are advised to consult the UPOV Code, which can be found on the UPOV Website ([www.upov.int](http://www.upov.int)), for the latest information.]

<u>TABLE OF CONTENTS</u>	<u>PAGE</u>
1. SUBJECT OF THESE TEST GUIDELINES.....	<a href="#">3</a>
2. MATERIAL REQUIRED.....	<a href="#">3</a>
3. METHOD OF EXAMINATION.....	<a href="#">4</a>
3.1 Number of Growing Cycles.....	<a href="#">4</a>
3.2 Testing Place.....	<a href="#">4</a>
3.3 Conditions for Conducting the Examination.....	<a href="#">4</a>
3.4 Test Design.....	<a href="#">4</a>
3.5 Additional Tests.....	<a href="#">4</a>
4. ASSESSMENT OF DISTINCTNESS, UNIFORMITY AND STABILITY.....	<a href="#">5</a>
4.1 Distinctness.....	<a href="#">5</a>
4.2 Uniformity.....	<a href="#">5</a>
4.3 Stability.....	<a href="#">5</a>
5. GROUPING OF VARIETIES AND ORGANIZATION OF THE GROWING TRIAL.....	<a href="#">6</a>
6. INTRODUCTION TO THE TABLE OF CHARACTERISTICS.....	<a href="#">7</a>
6.1 Categories of Characteristics.....	<a href="#">7</a>
6.2 States of Expression and Corresponding Notes.....	<a href="#">7</a>
6.3 Types of Expression.....	<a href="#">7</a>
6.4 Example Varieties.....	<a href="#">7</a>
6.5 Legend.....	<a href="#">7</a>
7. TABLE OF CHARACTERISTICS/TABLEAU DES CARACTÈRES/MERKMALSTABELLE/TABLA DE CARACTERES.....	<a href="#">8</a>
8. EXPLANATIONS ON THE TABLE OF CHARACTERISTICS.....	<a href="#">9</a>
8.1 Explanations covering several characteristics.....	<a href="#">9</a>
8.2 Explanations for individual characteristics.....	<a href="#">9</a>
9. LITERATURE.....	<a href="#">9</a>
10. TECHNICAL QUESTIONNAIRE.....	<a href="#">11</a>

1. Subject of these Test Guidelines

These Test Guidelines apply to all varieties of *Brassica juncea* (L.) Czern.

2. Material Required

2.1 The competent authorities decide on the quantity and quality of the plant material required for testing the variety and when and where it is to be delivered. Applicants submitting material from a State other than that in which the testing takes place must ensure that all customs formalities and phytosanitary requirements are complied with.

2.2 The material is to be supplied in the form of seed.

2.3 The minimum quantity of plant material, to be supplied by the applicant, should be:

3,000 seeds for single spaced plants. 20,000 seeds for drilled plants.

The seed should meet the minimum requirements for germination, species and analytical purity, health and moisture content, specified by the competent authority. In cases where the seed is to be stored, the germination capacity should be as high as possible and should, be stated by the applicant.

2.4 The plant material supplied should be visibly healthy, not lacking in vigor, nor affected by any important pest or disease.

2.5 The plant material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If it has been treated, full details of the treatment must be given.

### 3. Method of Examination

#### 3.1 *Number of Growing Cycles*

The minimum duration of tests should normally be two independent growing cycles.

#### 3.2 *Testing Place*

Tests are normally conducted at one place. In the case of tests conducted at more than one place, guidance is provided in TGP/9 "Examining Distinctness".

#### 3.3 *Conditions for Conducting the Examination*

3.3.1 The tests should be carried out under conditions ensuring satisfactory growth for the expression of the relevant characteristics of the variety and for the conduct of the examination.

3.3.2 The optimum stage of development for the assessment of each characteristic is indicated by a number in the second column of the Table of Characteristics. The stages of development denoted by each number are described in Chapter 8.

#### 3.4 *Test Design*

3.4.1 single spaced plants: Each test should be designed to result in a total of at least 60 Plants, which should be divided between at least 2 replicates.

3.4.2 drilled plots: Each test should be design to result in a total of at least 200 Plants which should be divided between at least 2 replicates.

#### 3.5 *Additional Tests*

Additional tests, for examining relevant characteristics, may be established.

#### 4. Assessment of Distinctness, Uniformity and Stability

##### 4.1 *Distinctness*

###### 4.1.1 General Recommendations

It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding distinctness. However, the following points are provided for elaboration or emphasis in these Test Guidelines.

###### 4.1.2 Consistent Differences

The differences observed between varieties may be so clear that more than one growing cycle is not necessary. In addition, in some circumstances, the influence of the environment is not such that more than a single growing cycle is required to provide assurance that the differences observed between varieties are sufficiently consistent. One means of ensuring that a difference in a characteristic, observed in a growing trial, is sufficiently consistent is to examine the characteristic in at least two independent growing cycles.

###### 4.1.3 Clear Differences

Determining whether a difference between two varieties is clear depends on many factors, and should consider, in particular, the type of expression of the characteristic being examined, i.e. whether it is expressed in a qualitative, quantitative, or pseudo-qualitative manner. Therefore, it is important that users of these Test Guidelines are familiar with the recommendations contained in the General Introduction prior to making decisions regarding distinctness.

###### 4.1.4 Number of plants or parts of plants to be Examined

Unless otherwise indicated, for the purposes of distinctness, all observations on single plants should be made on 30 plants or parts of plants taken from each of 30 plants and any other observations made on all plants in the test, disregarding any off-type plants.

###### 4.1.5 Method of Observation

The recommended method of observing the characteristic for the purposes of distinctness is indicated by the following key in the second column of the Table of Characteristics (see document TGP/9 "Examining Distinctness", Section 4 "Observation of characteristics"):

MG: single measurement of a group of plants or parts of plants

MS: measurement of a number of individual plants or parts of plants

VG: visual assessment by a single observation of a group of plants or parts of plants

VS: visual assessment by observation of individual plants or parts of plants

Type of observation: visual (V) or measurement (M)

"Visual" observation (V) is an observation made on the basis of the expert's judgment. For the purposes of this document, "visual" observation refers to the sensory observations of the experts and, therefore, also includes smell, taste and touch. Visual observation includes observations where the expert uses reference points (e.g. diagrams, example varieties, side-by-side comparison) or non-linear charts (e.g. color charts). Measurement (M) is an objective observation against a calibrated, linear scale e.g. using a ruler, weighing scales, colorimeter, dates, counts, etc.

Type of record: for a group of plants (G) or for single, individual plants (S)

For the purposes of distinctness, observations may be recorded as a single record for a group of plants or parts of plants (G), or may be recorded as records for a number of single, individual plants or parts of plants (S). In most cases, "G" provides a single record per variety and it is not possible or necessary to apply statistical methods in a plant-by-plant analysis for the assessment of distinctness.

In cases where more than one method of observing the characteristic is indicated in the Table of Characteristics (e.g. VG/MG), guidance on selecting an appropriate method is provided in document TGP/9, Section 4.2.

## 4.2 *Uniformity*

- 4.2.1 It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding uniformity. However, the following points are provided for elaboration or emphasis in these Test Guidelines:
- 4.2.2 The assessment of uniformity for cross-pollinated varieties should be according to the recommendations for cross-pollinated varieties in the General Introduction.
- 4.2.3 For the assessment of uniformity of self-pollinated varieties, a population standard of 2% and an acceptance probability of at least 95% should be applied. In the case of a sample size of 60 plants, 3 off-types are allowed. In the case of a sample size of 200 plants for drilled plants, 7 off-types are allowed.

## 4.3 *Stability*

- 4.3.1 In practice, it is not usual to perform tests of stability that produce results as certain as those of the testing of distinctness and uniformity. However, experience has demonstrated that, for many types of variety, when a variety has been shown to be uniform, it can also be considered to be stable.
- 4.3.2 Where appropriate, or in cases of doubt, stability may be further examined by testing a new seed stock to ensure that it exhibits the same characteristics as those shown by the initial material supplied.

5. Grouping of Varieties and Organization of the Growing Trial

5.1 The selection of varieties of common knowledge to be grown in the trial with the candidate varieties and the way in which these varieties are divided into groups to facilitate the assessment of distinctness are aided by the use of grouping characteristics.

5.2 Grouping characteristics are those in which the documented states of expression, even where produced at different locations, can be used, either individually or in combination with other such characteristics: (a) to select varieties of common knowledge that can be excluded from the growing trial used for examination of distinctness; and (b) to organize the growing trial so that similar varieties are grouped together.

5.3 The following have been agreed as useful grouping characteristics:

- (a) Seed: color (characteristic 1)
- (b) Leaf blade: anthocyanin coloration (characteristic 14)
- (c) Leaf blade: density of incisions of margin (excluding type2) (characteristic 17)
- (d) Leaf blade: blistering (excluding type2) (characteristic 18)
- (e) Plant: head formation (characteristic 20)

In the first place, the collection should be divided according to leaf types in Table 1 of Chapter 8.1.

5.4 Guidance for the use of grouping characteristics, in the process of examining distinctness, is provided through the General Introduction and document TGP/9 "Examining Distinctness".

## 6. Introduction to the Table of Characteristics

### 6.1 *Categories of Characteristics*

#### 6.1.1 Standard Test Guidelines Characteristics

Standard Test Guidelines characteristics are those which are approved by UPOV for examination of DUS and from which members of the Union can select those suitable for their particular circumstances.

#### 6.1.2 Asterisked Characteristics

Asterisked characteristics (denoted by \*) are those included in the Test Guidelines which are important for the international harmonization of variety descriptions and should always be examined for DUS and included in the variety description by all members of the Union, except when the state of expression of a preceding characteristic or regional environmental conditions render this inappropriate.

### 6.2 *States of Expression and Corresponding Notes*

6.2.1 States of expression are given for each characteristic to define the characteristic and to harmonize descriptions. Each state of expression is allocated a corresponding numerical note for ease of recording of data and for the production and exchange of the description.

6.2.2 In the case of qualitative and pseudo-qualitative characteristics (see Chapter 6.3), all relevant states of expression are presented in the characteristic. However, in the case of quantitative characteristics with 5 or more states, an abbreviated scale may be used to minimize the size of the Table of Characteristics. For example, in the case of a quantitative characteristic with 9 states, the presentation of states of expression in the Test Guidelines may be abbreviated as follows:

State	Note
small	3
medium	5
large	7

However, it should be noted that all of the following 9 states of expression exist to describe varieties and should be used as appropriate:

State	Note
very small	1
very small to small	2
small	3
small to medium	4
medium	5
medium to large	6
large	7
large to very large	8
very large	9

6.2.3 Further explanation of the presentation of states of expression and notes is provided in document TGP/7 "Development of Test Guidelines".

### 6.3 *Types of Expression*

An explanation of the types of expression of characteristics (qualitative, quantitative and pseudo-qualitative) is provided in the General Introduction.

### 6.4 *Example Varieties*

Where appropriate, example varieties are provided to clarify the states of expression of each characteristic.



6.5 Legend

		English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
1	2	3	4	5	6	7	
		Name of characteristics in English	Nom du caractère en français	Name des Merkmals auf Deutsch	Nombre del carácter en español		
		states of expression	types d'expression	Ausprägungsstufen	tipos de expresión		

1 Characteristic number

2 (\*) Asterisked characteristic – see Chapter 6.1.2

3 Type of expression

QL Qualitative characteristic – see Chapter 6.3

QN Quantitative characteristic – see Chapter 6.3

PQ Pseudo-qualitative characteristic – see Chapter 6.3

4 Method of observation (and type of plot, if applicable)

MG, MS, VG, VS – see Chapter 4.1.5

5 (+) See Explanations on the Table of Characteristics in Chapter 8.2

6 (a)-(f) See Explanations on the Table of Characteristics in Chapter 8.1

7 Growth stage key See Explanations on the Table of Characteristics in Chapter 8

7. Table of Characteristics/Tableau des caractères/Merkmalstabelle/Tabla de caracteres

	English		français		deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
<b>1.</b>	<b>(*)</b>	<b>PQ</b>	<b>VG</b>	<b>(+)</b>	<b>(a)</b>	<b>00</b>		
		<b>Seed: color</b>						
		yellow					Kigarashina	1
		brown					Akaoba Takana(Red Giant), Esperance, Miike Takana, Terrafit, Terraplus	2
		black					Hagarashina, TTK456(Chaplin)	3
<b>2.</b>		<b>QN</b>	<b>VG</b>		<b>(a)</b>	<b>10</b>		
		<b>Hypocotyl: anthocyanin coloration</b>						
		absent or weak					Zasai FM-58	1
		medium					Shinkoku Seisai	2
		strong					TTK456(Chaplin)	3
<b>3.</b>		<b>QN</b>	<b>MS/VG</b>		<b>(a), (c)</b>	<b>10</b>		
		<b>Cotyledon: length</b>						
		short					Junkei Yamashiona, Vittasso	3
		medium					Katsuona, Terraplus	5
		long					Scala	7
<b>4.</b>		<b>QN</b>	<b>MS/VG</b>		<b>(a), (c)</b>	<b>10</b>		
		<b>Cotyledon: width</b>						
		narrow					Junkei Yamashiona, Vittasso	3
		medium					Katsuona, Pacific Gold, Terraplus	5
		broad					Minaret, Terminator	7

	English		français		deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
<b>5.</b>	<b>PQ</b>	<b>VG</b>	<b>(+)</b>	<b>(a), (e)</b>	<b>19</b>			
	<b>Leaf: shape</b>							
		ovate					Serihon	1
		circular					Kekkyu Takana	2
		elliptic					Akariasu	3
		oblong					Etamine, Zasai FM-58	4
		obovate					Esperance, Katsuona	5
		spatulate					Kigarashina	6
<b>6.</b>	<b>QN</b>	<b>VG</b>	<b>(+)</b>	<b>(a), (e)</b>	<b>19</b>			
	<b>Leaf: attitude</b>							
		erect					Wasabina	1
		semi-erect					Esperance, Shinkoku Seisai	3
		horizontal					Etamine, Miike Takana	5
<b>7.</b>	<b>QN</b>	<b>MS/VG</b>	<b>(a), (d), (e)</b>		<b>19</b>			
	<b>Leaf: length</b>							
		short					Chirimen Hakarashina	3
		medium					Miike Takana	5
		long					Akaoba Takana(Red Giant)	7
<b>8.</b>	<b>QN</b>	<b>MS/VG</b>	<b>(a), (d), (e)</b>		<b>19</b>			
	<b>Leaf: width</b>							
		narrow					Chirimen Hakarashina	3
		medium					Miike Takana	5
		broad					Katsuona	7
<b>9.</b>	<b>QN</b>	<b>MS/VG</b>	<b>(a), (d), (e)</b>		<b>19</b>			
	<b>Leaf: length of petiole</b>							
		absent or very short					Serihon	1
		short					Miike Takana	3
		medium					Junkei Yamashiona	5
		long					Kigarashina	7

	English		français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
<b>10.</b>	<b>QN</b>	<b>MS/VG</b>	<b>(a), (d), (e)</b>	<b>19</b>			
	<b>Leaf: width of petiole</b>						
	narrow					Kigarashina	3
	medium					Katsuona	5
	broad					Shinkoku Seisai	7
<b>11.</b>	<b>QN</b>	<b>MS/VG</b>	<b>(+)</b>	<b>(a), (b), (e)</b>	<b>19</b>		
	<b>Leaf blade: size of terminal lobe (total surface)</b>						
	small					Chirimen Hakarashina, Etamine	3
	medium					Kigarashina	5
	large					Pacific Gold, Perm Green	7
<b>12.</b>	<b>QN</b>	<b>VG</b>	<b>(a), (b), (e)</b>	<b>19</b>			
	<b>Leaf blade: number of lateral lobes</b>						
	absent or very few						1
	few					Minaret	3
	medium					Esperance, Kigarashina	5
	many					Akariasu, Etamine, TTK456(Chaplin)	7
<b>13.</b>	<b>QN</b>	<b>VG</b>	<b>(a), (e)</b>	<b>19</b>			
	<b>Leaf blade: pubescence on lower side</b>						
	absent or weak					Miike Takana	1
	medium					Oba Takana	2
	strong					Kigarashina	3
<b>14.</b>	<b>QN</b>	<b>VG</b>	<b>(+)</b>	<b>(a), (e)</b>	<b>19</b>		
	<b>Leaf blade: anthocyanin coloration</b>						
	absent or very weak					Kekkyu Takana, Vitamine	1
	weak						3
	medium					Miike Takana	5
	strong					Akaoba Takana(Red Giant)	7

	English		français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
<b>15.</b>	<b>QN</b>	<b>VG</b>	<b>(a), (e)</b>	<b>19</b>			
	<b>Only varieties with anthocyanin coloration absent or very weak: Leaf blade: intensity of green color</b>						
	light					Wasabina	3
	medium					Etamine, Golden Streaks, Katsuona	5
	dark					Minaret	7
<b>16.</b>	<b>QN</b>	<b>VG</b>	<b>(a), (e)</b>	<b>19</b>			
	<b>Leaf blade: undulation of margin (excluding type2)</b>						
	absent or very weak						1
	weak					Akaoba Takana(Red Giant)	3
	medium					Katsuona	5
	strong					Chirimen Hakarashina	7
<b>17.</b>	<b>QN</b>	<b>VG</b>	<b>(+)</b>	<b>(a), (e)</b>	<b>19</b>		
	<b>Leaf blade: density of incisions of margin (excluding type2)</b>						
	absent or very sparse						1
	sparse					Etamine, Katsuona	3
	medium					Opaleska	5
	dense					Oportuna	7
<b>18.</b>	<b>QN</b>	<b>VG</b>	<b>(+)</b>	<b>(a), (b), (e)</b>	<b>19</b>		
	<b>Leaf blade: blistering (excluding type2)</b>						
	absent or weak					Etamine, Kigarashina	1
	medium					Akaoba Takana(Red Giant)	2
	strong					Katsuona	3

	English		français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
19.	QN	MS/VG	(a), (b), (c), (e)	19			
	<b>Only varieties with leaf: type: type3 and 4: Leaf blade: width of midrib at widest point</b>						
	narrow					Sagami Green	3
	medium					Katsuona	5
	broad					Shinkoku Seisai	7
20. (*)	QL	VG	(+)	(a), (b)	19		
	<b>Plant: head formation</b>						
	absent					Kigarashina	1
	present					Kekkyu Takana	9
21.	QN	MS/VG	(a)	19			
	<b>Head: height</b>						
	short						1
	medium					Unzen Kekkyu Takana	2
	tall						3
22.	QN	MS/VG	(a)	19			
	<b>Head: width</b>						
	narrow						1
	medium					Kekkyu Takana	2
	broad						3
23.	QN	MS/VG	(+)	(a)	19		
	<b>Head: number of leaves</b>						
	few						3
	medium					Kekkyu Takana	5
	many						7
24.	PQ	VG	(a)	19			
	<b>Head: internal color</b>						
	yellowish white					Unzen Kekkyu Takana	1
	light green						2
	medium green					Kekkyu Takana	3

	English		français		deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
<b>25.</b>	<b>PQ</b>	<b>VG</b>	<b>(+)</b>	<b>(a)</b>	<b>20-29</b>			
	<b>Stem: type of main stem (excluding heading type)</b>							
	narrow type						Kigarashina	1
	round type						Umino	2
	oblong type						Zasai FM-58	3
	budding type						FE-K226	4
<b>26.</b>	<b>QN</b>	<b>MG</b>	<b>(a)</b>		<b>31</b>			
	<b>Time of beginning of bolting</b>							
	early						Junkei Yamashiona, Scala	3
	medium						Katsuona	5
	late						Akaoba Takana(Red Giant)	7
<b>27.</b>	<b>QN</b>	<b>MG</b>	<b>(a)</b>		<b>50</b>			
	<b>Time of flowering</b>							
	early						Terrafit	3
	medium						Terraplus	5
	late							7
	very late						Vitasso	9
<b>28.</b>	<b>QL</b>	<b>VG</b>	<b>(a)</b>		<b>50-59</b>			
	<b>Only varieties with head formation: absent: Production of pollen</b>							
	absent							1
	present							9
<b>29.</b>	<b>QN</b>	<b>MS/VG</b>	<b>(a)</b>		<b>61</b>			
	<b>Only varieties with head formation: absent: Plant: length</b>							
	short						Pacific Gold, Terminator	3
	medium						Terraplus	5
	long						Minaret	7
	very long						Vitasso	9

	English		français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
<b>30.</b>	<b>QN</b>	<b>MS/VG</b>	<b>(a), (f)</b>	<b>79</b>			
	<b>Only varieties with head formation: absent: siliqua: length</b>						
	short					Terraplus, Vittasso	3
	medium					Pacific Gold	5
	long					Minaret	7
<b>31.</b>	<b>QN</b>	<b>MS/VG</b>	<b>(a), (f)</b>	<b>61</b>			
	<b>Only varieties with head formation: absent: Siliqua: length of beak</b>						
	short					Terraplus, Vittasso	3
	medium					Terrafit	5
	long						7
<b>32.</b>	<b>QN</b>	<b>MS/VG</b>	<b>(a), (f)</b>	<b>61</b>			
	<b>Only varieties with head formation: absent: Siliqua: width</b>						
	narrow					Vittasso	3
	medium					Energy, Terrafit	5
	broad						7
<b>33.</b>	<b>QN</b>	<b>MS/VG</b>	<b>(a), (f)</b>	<b>61</b>			
	<b>Only varieties with head formation: absent: Siliqua: length of peduncle</b>						
	short					Vittasso	3
	medium					Energy	5
	long						7



	English		français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
34.	QN	MS/VG	(a)				
	<b>Generative development in the year of sowing for late summer sown trials</b>						
	absent or very weak					Vitasso	1
	weak						3
	medium					Terraplus	5
	strong					Energy, Terrafit, Terratop	7
	very strong						9

8. Explanations on the Table of Characteristics

8.1 *Explanations covering several characteristics*

Characteristics containing the following key in the second column of the Table of Characteristics should be examined as indicated below:


(a) Table1: KEY FOR THE STAGE OF DEVELOPMENT

KEY	GENERAL DESCRIPTION
0	<u>Principal growth stage 0: Germination</u>
01	Beginning of seed imbibition
03	Seed imbibition complete
05	Radicle emerged from seed
07	Hypocotyl with cotyledons emerged from seed
08	Hypocotyl with cotyledons growing towards soil
09	surface Emergence: cotyledons emerge through soil surface
1	<u>Principal growth stage 1: Leaf development</u>
10	Cotyledons completely unfolded
11	First leaf unfolded
12	2 leaves unfolded
13	3 leaves unfolded
14	4 leaves unfolded
15	5 leaves unfolded
16	6 leaves unfolded
17	7 leaves unfolded
18	8 leaves unfolded
19	9 or more leaves unfolded
2	<u>Principal growth stage 2: Formation of side shoots</u>
20	No side shoots
21	First side shoot detectable
22	2 side shoots detectable
23	3 side shoots detectable
24	4 side shoots detectable
25	5 side shoots detectable
26	6 side shoots detectable
27	7 side shoots detectable
28	8 side shoots detectable
29	9 or more side shoots detectable
3	<u>Principal growth stage 3: Stem elongation</u>
30	no internodes("rosette")
31	1 visibly extended internode
32	2 visibly extended internodes
33	3 visibly extended internodes
34	4 visibly extended internodes
35	5 visibly extended internodes
36	6 visibly extended internodes
37	7 visibly extended internodes
38	8 visibly extended internodes
39	9 or more visibly extended internodes
4	<u>Principal growth stage 5: Inflorescence emergence</u>
40	Flower buds present, still enclosed by leaves
41	Flower buds visible from above ("green bud")
42	Flower buds free, level with the youngest leaves
43	Flower buds raised above the youngest leaves
44	Individual flower buds (main inflorescence)
45	visible but still closed
46	Individual flower buds (secondary inflorescence)
47	


49	visible but still closed First petals visible, flower buds still closed by ("yellow bud")
5	<u>Principal growth stage 6: Formation of side shoots</u>
50	First flowers open
51	10% of flowers on main raceme open, main raceme elongating
52	20% of flowers on main raceme open
53	30% of flowers on main raceme open
54	40% of flowers on main raceme open
55	Full flowering: 50% flowers on main raceme open, older petals falling
57	Flowering declining: majority of petals fallen
59	End of flowering
6	<u>Principal growth stage 7: Development of fruit</u>
61	10% of pods have reached final size
62	20% of pods have reached final size
63	30% of pods have reached final size
64	40% of pods have reached final size
65	50% of pods have reached final size
66	60% of pods have reached final size
67	70% of pods have reached final size
68	80% of pods have reached final size
69	Nearly all pods have reached final size
7	<u>Principal growth stage 8: Ripening</u>
70	seed green, filling pod cavity
71	10% of pods ripe, seeds dark and hard
72	20% of pods ripe, seeds dark and hard
73	30% of pods ripe, seeds dark and hard
74	40% of pods ripe, seeds dark and hard
75	50% of pods ripe, seeds dark and hard
76	60% of pods ripe, seeds dark and hard
77	70% of pods ripe, seeds dark and hard
78	80% of pods ripe, seeds dark and hard
79	Fully ripe: nearly all pods ripe, seeds dark and hard
8	<u>Principal growth stage 2: Formation of side shoots</u>
87	Plant dead and dry
89	Harvested product

(b) Table 2: LEAF TYPE


Leaf: type	Example varieties	Leaf blade: size of terminal lobe (12)	Leaf blade: number of lateral lobes (13)	Leaf blade: blistering (excluding type2) (20)	Leaf blade: width of midrib at widest point(21)	Plant: head formation (22)
type1	Hagarashina, Kigarashina	medium to very large	few to medium	absent or weak to medium	-	absent
type2	Akariasu, Riasu, Karashina	very small to small	many to very many	-	-	absent
type3	Oba Takana, Sagami Green	-	absent or very few	medium to strong	narrow	absent
type4	Kekkyu, Takana, Miike, Takana, Shinkoku, Seisai	-	absent or very few	medium to strong	medium to broad	absent or present




1  
type1  
(lyrate)



2  
type2  
(division)




3  
type3  
(entire and narrow midrib)




4  
type4  
(entire and broad midrib)

- (c) The measurements should be taken in the glasshouse on cotyledons of 30 seedlings. If the two cotyledons differ in size, the biggest one should be measured. The length is defined as distance between the inclination at top of the cotyledon and the point where the width of the petiole is about 4 mm. The width of the cotyledon should be measured at the widest point of the cotyledons.




- (d)



- (e) observations on the leaves should be made on the largest fully developed (non-senescent) leaf.

(f)




## 8.2 Explanations for individual characteristics

### Ad. 1: Seed: color



1  
yellow




2  
brown



3  
black


Ad. 5: Leaf: shape



Ad. 6: Leaf: attitude




Ad. 11: Leaf blade: size of terminal lobe (total surface)



Ad. 14: Leaf blade: anthocyanin coloration

the strongest intensity of anthocyanin should be observed (not the extension).

Ad. 17: Leaf blade: density of incisions of margin (excluding type2)





Ad. 18: Leaf blade: blistering (excluding type2)



1  
absent or weak



2  
medium



3  
strong

Ad. 20: Plant: head formation



1  
absent




9  
present

Ad. 23: Head: number of leaves

Observation on the leaves which form the head.

Ad. 25: Stem: type of main stem (excluding heading type)

Observation on type of main stem should be made on shape of main stem without lateral stem which locate at the base of main stem..



8.3 *Other Names of the Example Varieties*

TTK456 <sup>(1)</sup>	Chaplin <sup>(2)</sup>
Akaoba Takana <sup>(3)</sup>	Red Giant <sup>(4)</sup>

(1) official denomination registered under the law in Japan in 2011.

(2) official denomination of TTK456 registered under the law in European Union in 2014.

(3) commercial name in Japan.

(4) commercial name of Akaoba Takana in European Union.

9. Literature

- Fujishiro, T., 1996: Breeding processes and characteristics of a newly bred leaf mustard (*Brassica Juncea* Coss.), Kanagawa, Japan.
- Joy Larkcom., 1991: Oriental Vegetables (The Complete guide for Garden and Kitchen) London, United Kingdom. pp. 39 to 45
- Ministry of Agriculture, Forestry & Fisheries of Japan., 1994: National Test Guideline for Karashina.
- Roger Phillips, Martyn Rix., 1993: VEGETABLES (The Pan Garden Plants Series) p.44
- Tsukamoto, Y., 1994: The Grand Dictionary of Horticulture Volume 1. The Shogakukan Ltd., Tokyo, Japan, pp. 520 to 522.
- Takasi A., 2004: Yasai-engei-daihyakka 17. Shadanhoin Nousan-gyoson-bunkakyokai. Tokyo, Japan. 169-233
- Uwe Meier. Federal Biological Research Centre for Agriculture and Forestry, 2001: Growth stages of mono- and dicotyledonous plants, BBCH Monograph,

10. Technical Questionnaire

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
		Application date: (not to be filled in by the applicant)
TECHNICAL QUESTIONNAIRE to be completed in connection with an application for plant breeders' rights		
1. Subject of the Technical Questionnaire		
1.1	Botanical name	<input type="text" value="Brassica juncea (L.) Czern."/>
1.2	Common name	<input type="text" value="Brown mustard, India mustard, Indian mustard, Oriental mustard"/>
2. Applicant		
	Name	<input type="text"/>
	Address	<input type="text"/>
	Telephone No.	<input type="text"/>
	Fax No.	<input type="text"/>
	E-mail address	<input type="text"/>
	Breeder (if different from applicant)	<input type="text"/>
3. Proposed denomination and breeder's reference		
	Proposed denomination (if available)	<input type="text"/>
	Breeder's reference	<input type="text"/>

#4. Information on the breeding scheme and propagation of the variety

4.1 Breeding scheme

Variety resulting from:

4.1.1 Crossing

(a) controlled cross [ ]  
(please state parent varieties)

(.....) x (.....)  
female parent male parent

(b) partially known cross [ ]  
(please state known parent variety(ies))

(.....) x (.....)  
female parent male parent

(c) unknown cross [ ]

4.1.2 Mutation [ ]  
(please state parent variety)

4.1.3 Discovery and development [ ]  
(please state where and when discovered and how developed)

4.1.4 Other [ ]  
(please provide details)

# Authorities may allow certain of this information to be provided in a confidential section of the Technical Questionnaire.

4.2 Method of propagating the variety

4.2.1 Seed-propagated varieties

- (a) Cross-pollination [ ]  
(b) Other (please provide details) [ ]

- 4.2.2 Other [ ]  
(Please provide details)

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

5. Characteristics of the variety to be indicated (the number in brackets refers to the corresponding characteristic in Test Guidelines; please mark the note which best corresponds).

Characteristics	Example Varieties	Note
<b>5.1 Seed: color</b>		
<b>(1)</b>		
yellow	Kigarashina	1 [ ]
brown	Akaoba Takana(Red Giant), Esperance, Miike Takana, Terrafit, Terraplus	2 [ ]
black	Hagarashina, TTK456(Chaplin)	3 [ ]
<b>5.2 Leaf: shape</b>		
<b>(5)</b>		
ovate	Serihon	1 [ ]
circular	Kekkyu Takana	2 [ ]
elliptic	Akariasu	3 [ ]
oblong	Etamine, Zasai FM-58	4 [ ]
obovate	Esperance, Katsuona	5 [ ]
spatulate	Kigarashina	6 [ ]
<b>5.3 Leaf: attitude</b>		
<b>(6)</b>		
erect	Wasabina	1 [ ]
semi-erect	Esperance, Shinkoku Seisai	3 [ ]
horizontal	Etamine, Miike Takana	5 [ ]
<b>5.4 Leaf blade: anthocyanin coloration</b>		
<b>(14)</b>		
absent or very weak	Kekkyu Takana, Vitamine	1 [ ]
weak		3 [ ]
medium	Miike Takana	5 [ ]
strong	Akaoba Takana(Red Giant)	7 [ ]

Characteristics	Example Varieties	Note
<b>5.5 Leaf blade: undulation of margin (excluding type2)</b>		
<b>(16)</b>		
absent or very weak		1 [ ]
very weak to weak		2 [ ]
weak	Akaoba Takana(Red Giant)	3 [ ]
weak to medium		4 [ ]
medium	Katsuona	5 [ ]
medium to strong		6 [ ]
strong	Chirimem Hakarashina	7 [ ]
strong to very strong		8 [ ]
very strong		9 [ ]
<b>5.6 Leaf blade: density of incisions of margin (excluding type2)</b>		
<b>(17)</b>		
absent or very sparse		1 [ ]
sparse	Etamine, Katsuona	3 [ ]
medium	Opaleska	5 [ ]
dense	Oportuna	7 [ ]
<b>5.7 Leaf blade: blistering (excluding type2)</b>		
<b>(18)</b>		
absent or weak	Etamine, Kigarashina	1 [ ]
medium	Akaoba Takana(Red Giant)	2 [ ]
strong	Katsuona	3 [ ]
<b>5.8 Plant: head formation</b>		
<b>(20)</b>		
absent	Kigarashina	1 [ ]
present	Kekkyu Takana	9 [ ]



TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

6. Similar varieties and differences from these varieties

*Please use the following table and box for comments to provide information on how your candidate variety differs from the variety (or varieties) which, to the best of your knowledge, is (or are) most similar. This information may help the examination authority to conduct its examination of distinctness in a more efficient way.*

Denomination(s) of variety(ies) similar to your candidate variety	Characteristic(s) in which your candidate variety differs from the similar variety(ies)	Describe the expression of the characteristic(s) for the <b>similar</b> variety(ies)	Describe the expression of the characteristic(s) for <b>your</b> candidate variety
<i>Example</i>	<i>Leaf: shape</i>	<i>ovate</i>	<i>oblong</i>
Comments:			

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

#7. Additional information which may help in the examination of the variety

7.1 In addition to the information provided in sections 5 and 6, are there any additional characteristics which may help to distinguish the variety?

Yes  No

(If yes, please provide details)

7.2 Are there any special conditions for growing the variety or conducting the examination?

Yes  No

(If yes, please provide details)

7.3 Other information

(a) Main use

Vegetable   
Seed i oil ii condiment   
Green manure   
other

(b) Leaf type (see chapter 8.1 (b) for explanation)

type 1   
type 2   
type 3   
type 4

8. Authorization for release			
(a)	Does the variety require prior authorization for release under legislation concerning the protection of the environment, human and animal health?		
Yes	<input type="checkbox"/>	No	<input type="checkbox"/>
(b)	Has such authorization been obtained?		
Yes	<input type="checkbox"/>	No	<input type="checkbox"/>
If the answer to (b) is yes, please attach a copy of the authorization.			
9. Information on plant material to be examined or submitted for examination			
9.1 The expression of a characteristic or several characteristics of a variety may be affected by factors, such as pests and disease, chemical treatment (e.g. growth retardants or pesticides), effects of tissue culture, different rootstocks, scions taken from different growth phases of a tree, etc.			
9.2 The plant material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If the plant material has undergone such treatment, full details of the treatment must be given. In this respect, please indicate below, to the best of your knowledge, if the plant material to be examined has been subjected to:			
(a)	Microorganisms (e.g. virus, bacteria, phytoplasma)	Yes <input type="checkbox"/>	No <input type="checkbox"/>
(b)	Chemical treatment (e.g. growth retardant, pesticide)	Yes <input type="checkbox"/>	No <input type="checkbox"/>
(c)	Tissue culture	Yes <input type="checkbox"/>	No <input type="checkbox"/>
(d)	Other factors	Yes <input type="checkbox"/>	No <input type="checkbox"/>
Please provide details for where you have indicated "yes".			
.....			
10. I hereby declare that, to the best of my knowledge, the information provided in this form is correct:			
Applicant's name	<input type="text"/>		
Signature	<input type="text"/>		
Date	<input type="text"/>		

[End of document]