



TG/NASTU(proj.2)

ORIGINAL: English

DATE: 2017-05-12

INTERNATIONAL UNION FOR THE PROTECTION OF NEW VARIETIES OF PLANTS

Geneva

DRAFT

WATERCRESS

UPOV Code(s):

NASTU_STE; NASTU_MIC; NASTU_OFF

Nasturtium microphyllum Boenn. ex Rchb.;
Nasturtium officinale R. Br.;
Nasturtium xsterile (Airy Shaw) Oefelein

GUIDELINES

FOR THE CONDUCT OF TESTS

FOR DISTINCTNESS, UNIFORMITY AND STABILITY

*prepared by experts from United Kingdom
to be considered by the
Technical Working Party for Vegetables
at its fifty-first session, to be held in Roelofarendsveen, Netherlands,
from 2017-07-03 to 2017-07-07*

Disclaimer: this document does not represent UPOV policies or guidance

Alternative names:*

Botanical name	English	French	German	Spanish
<i>Nasturtium microphyllum</i> Boenn. ex Rchb.	One-row watercress			
<i>Nasturtium officinale</i> R. Br., <i>Rorippa nasturtium-aquaticum</i> (L.) Hayek	Watercress	cresson de fontaine; cresson d'eau	Brunnenkresse	berro
<i>Nasturtium xsterile</i> (Airy Shaw) Oefelein, <i>Nasturtium microphyllum</i> x <i>Nasturtium officinale</i> , <i>Rorippa microphylla</i> x <i>Rorippa nasturtium-aquaticum</i>				

* These names were correct at the time of the introduction of these Test Guidelines but may be revised or updated. [Readers are advised to consult the UPOV Code, which can be found on the UPOV Website (www.upov.int), for the latest information.]

The purpose of these guidelines (“Test Guidelines”) is to elaborate the principles contained in the General Introduction (document TG/1/3), and its associated TGP documents, into detailed practical guidance for the harmonized examination of distinctness, uniformity and stability (DUS) and, in particular, to identify appropriate characteristics for the examination of DUS and production of harmonized variety descriptions.

ASSOCIATED DOCUMENTS

These Test Guidelines should be read in conjunction with the General Introduction and its associated TGP documents.

<u>TABLE OF CONTENTS</u>	<u>PAGE</u>
1. SUBJECT OF THESE TEST GUIDELINES.....	4
2. MATERIAL REQUIRED.....	4
3. METHOD OF EXAMINATION.....	5
3.1 Number of Growing Cycles.....	5
3.2 Testing Place.....	5
3.3 Conditions for Conducting the Examination.....	5
3.4 Test Design.....	5
3.5 Additional Tests.....	5
4. ASSESSMENT OF DISTINCTNESS, UNIFORMITY AND STABILITY.....	6
4.1 Distinctness.....	6
4.2 Uniformity.....	7
4.3 Stability.....	7
5. GROUPING OF VARIETIES AND ORGANIZATION OF THE GROWING TRIAL.....	8
6. INTRODUCTION TO THE TABLE OF CHARACTERISTICS.....	9
6.1 Categories of Characteristics.....	9
6.2 States of Expression and Corresponding Notes.....	9
6.3 Types of Expression.....	9
6.4 Example Varieties.....	10
6.5 Legend.....	11
7. TABLE OF CHARACTERISTICS/TABLEAU DES CARACTÈRES/MERKMALSTABELLE/TABLA DE CARACTERES.....	12
8. EXPLANATIONS ON THE TABLE OF CHARACTERISTICS.....	20
8.1 Explanations covering several characteristics.....	20
8.2 Explanations for individual characteristics.....	22
9. LITERATURE.....	22
10. TECHNICAL QUESTIONNAIRE.....	28

1. Subject of these Test Guidelines

These Test Guidelines apply to all varieties of *Nasturtium microphyllum* Boenn. ex Rchb., *Nasturtium officinale* R. Br and *Nasturtium xsterile* (Airy Shaw) Oefelein.

2. Material Required

2.1 The competent authorities decide on the quantity and quality of the plant material required for testing the variety and when and where it is to be delivered. Applicants submitting material from a State other than that in which the testing takes place must ensure that all customs formalities and phytosanitary requirements are complied with.

2.2 The material is to be supplied in the form of seed or plants.

2.3 The minimum quantity of plant material, to be supplied by the applicant, should be:

10 g for seed-propagated varieties
80 plants for vegetatively-propagated varieties

In the case of seed, the seed should meet the minimum requirements for germination, species and analytical purity, health and moisture content, specified by the competent authority. In cases where the seed is to be stored, the germination capacity should be as high as possible and should, be stated by the applicant.

2.4 The plant material supplied should be visibly healthy, not lacking in vigor, nor affected by any important pest or disease.

2.5 The plant material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If it has been treated, full details of the treatment must be given.

3. Method of Examination

3.1 *Number of Growing Cycles*

3.1.1 The minimum duration of tests should normally be two independent growing cycles.

3.1.2 The two independent growing cycles should be in the form of two separate plantings.

3.2 *Testing Place*

Tests are normally conducted at one place. In the case of tests conducted at more than one place, guidance is provided in TGP/9 "Examining Distinctness".

3.3 *Conditions for Conducting the Examination*

The tests should be carried out under conditions ensuring satisfactory growth for the expression of the relevant characteristics of the variety and for the conduct of the examination.

3.4 *Test Design*

3.4.1 In the case of seed-propagated varieties.: Each test should be designed to result in a total of at least 60 plants, which should be divided between at least 2 replicates.

3.4.2 In the case of vegetatively propagated varieties: Each test should be design to result in a total of at least 30 plants which should be divided between at least 2 replicates.

3.4.3 The design of the tests should be such that plants or parts of plants may be removed for measurement or counting without prejudice to the observations which must be made up to the end of the growing cycle.

3.5 *Additional Tests*

Additional tests, for examining relevant characteristics, may be established.

4. Assessment of Distinctness, Uniformity and Stability

4.1 *Distinctness*

4.1.1 General Recommendations

It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding distinctness. However, the following points are provided for elaboration or emphasis in these Test Guidelines.

4.1.2 Consistent Differences

The differences observed between varieties may be so clear that more than one growing cycle is not necessary. In addition, in some circumstances, the influence of the environment is not such that more than a single growing cycle is required to provide assurance that the differences observed between varieties are sufficiently consistent. One means of ensuring that a difference in a characteristic, observed in a growing trial, is sufficiently consistent is to examine the characteristic in at least two independent growing cycles.

4.1.3 Clear Differences

Determining whether a difference between two varieties is clear depends on many factors, and should consider, in particular, the type of expression of the characteristic being examined, i.e. whether it is expressed in a qualitative, quantitative, or pseudo-qualitative manner. Therefore, it is important that users of these Test Guidelines are familiar with the recommendations contained in the General Introduction prior to making decisions regarding distinctness.

4.1.4 Number of plants or parts of plants to be Examined

In the case of seed propagated varieties, unless otherwise indicated, for the purposes of distinctness, all observations on single plants should be made on 40 plants or parts taken from each of 40 plants and any other observation made on all plants in the test, disregarding any off-type plants.

In the case of observations of parts taken from single fruit bodies, the number of parts to be taken from each of the fruit bodies should be 1.

In the case of vegetatively propagated varieties, unless otherwise indicated, for the purposes of distinctness, all observations on single plants should be made on 20 plants or parts taken from each of 20 plants and any other observation made on all plants in the test, disregarding any off-type plants.

In the case of observations of parts taken from single fruit bodies, the number of parts to be taken from each of the fruit bodies should be 1.

4.1.5 Method of Observation

The recommended method of observing the characteristic for the purposes of distinctness is indicated by the following key in the second column of the Table of Characteristics (see document TGP/9 "Examining Distinctness", Section 4 "Observation of characteristics"):

MG: single measurement of a group of plants or parts of plants

MS: measurement of a number of individual plants or parts of plants

VG: visual assessment by a single observation of a group of plants or parts of plants

VS: visual assessment by observation of individual plants or parts of plants

Type of observation: visual (V) or measurement (M)

“Visual” observation (V) is an observation made on the basis of the expert’s judgment. For the purposes of this document, “visual” observation refers to the sensory observations of the experts and, therefore, also includes smell, taste and touch. Visual observation includes observations where the expert uses reference points (e.g. diagrams, example varieties, side-by-side comparison) or non-linear charts (e.g. color charts). Measurement (M) is an objective observation against a calibrated, linear scale e.g. using a ruler, weighing scales, colorimeter, dates, counts, etc.

Type of record: for a group of plants (G) or for single, individual plants (S)

For the purposes of distinctness, observations may be recorded as a single record for a group of plants or parts of plants (G), or may be recorded as records for a number of single, individual plants or parts of plants (S). In most cases, “G” provides a single record per variety and it is not possible or necessary to apply statistical methods in a plant-by-plant analysis for the assessment of distinctness.

In cases where more than one method of observing the characteristic is indicated in the Table of Characteristics (e.g. VG/MG), guidance on selecting an appropriate method is provided in document TGP/9, Section 4.2.

4.2 *Uniformity*

- 4.2.1 It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding uniformity. However, the following points are provided for elaboration or emphasis in these Test Guidelines:
- 4.2.2 The assessment of uniformity for cross-pollinated varieties should be according to the recommendations for cross-pollinated varieties in the General Introduction.
- 4.2.3 For the assessment of uniformity of seed-propagated varieties, a population standard of 2% and an acceptance probability of at least 95% should be applied. In the case of a sample size of 40 plants, 2 off-types are allowed.
- 4.2.4 For the assessment of vegetatively-propagated varieties, a population standard of 1% and an acceptance probability of at least 95 % will be applied. In the case of a sample size of 20 plants, 1 off-type is allowed.

4.3 *Stability*

- 4.3.1 In practice, it is not usual to perform tests of stability that produce results as certain as those of the testing of distinctness and uniformity. However, experience has demonstrated that, for many types of variety, when a variety has been shown to be uniform, it can also be considered to be stable.
- 4.3.2 Where appropriate, or in cases of doubt, stability may be further examined by testing a new seed or plant stock to ensure that it exhibits the same characteristics as those shown by the initial material supplied.

5. Grouping of Varieties and Organization of the Growing Trial
- 5.1 The selection of varieties of common knowledge to be grown in the trial with the candidate varieties and the way in which these varieties are divided into groups to facilitate the assessment of distinctness are aided by the use of grouping characteristics.
- 5.2 Grouping characteristics are those in which the documented states of expression, even where produced at different locations, can be used, either individually or in combination with other such characteristics: (a) to select varieties of common knowledge that can be excluded from the growing trial used for examination of distinctness; and (b) to organize the growing trial so that similar varieties are grouped together.
- 5.3 The following have been agreed as useful grouping characteristics:
 - (a) Plant: growth habit (characteristic 2)
 - (b) Time of beginning of flowering (characteristic 22)
- 5.4 Guidance for the use of grouping characteristics, in the process of examining distinctness, is provided through the General Introduction and document TGP/9 "Examining Distinctness".

6. Introduction to the Table of Characteristics

6.1 *Categories of Characteristics*

6.1.1 Standard Test Guidelines Characteristics

Standard Test Guidelines characteristics are those which are approved by UPOV for examination of DUS and from which members of the Union can select those suitable for their particular circumstances.

6.1.2 Asterisked Characteristics

Asterisked characteristics (denoted by *) are those included in the Test Guidelines which are important for the international harmonization of variety descriptions and should always be examined for DUS and included in the variety description by all members of the Union, except when the state of expression of a preceding characteristic or regional environmental conditions render this inappropriate.

6.2 *States of Expression and Corresponding Notes*

6.2.1 States of expression are given for each characteristic to define the characteristic and to harmonize descriptions. Each state of expression is allocated a corresponding numerical note for ease of recording of data and for the production and exchange of the description.

6.2.2 In the case of qualitative and pseudo-qualitative characteristics (see Chapter 6.3), all relevant states of expression are presented in the characteristic. However, in the case of quantitative characteristics with 5 or more states, an abbreviated scale may be used to minimize the size of the Table of Characteristics. For example, in the case of a quantitative characteristic with 9 states, the presentation of states of expression in the Test Guidelines may be abbreviated as follows:

<i>State</i>	<i>Note</i>
small	3
medium	5
large	7

However, it should be noted that all of the following 9 states of expression exist to describe varieties and should be used as appropriate:

<i>State</i>	<i>Note</i>
very small	1
very small to small	2
small	3
small to medium	4
medium	5
medium to large	6
large	7
large to very large	8
very large	9

6.2.3 Further explanation of the presentation of states of expression and notes is provided in document TGP/7 "Development of Test Guidelines".

6.3 *Types of Expression*

An explanation of the types of expression of characteristics (qualitative, quantitative and pseudo-qualitative) is provided in the General Introduction.

6.4 Example Varieties

Where appropriate, example varieties are provided to clarify the states of expression of each characteristic.

6.5 Legend

		English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
1	2	3	4	5	6	7	
		Name of characteristics in English	Nom du caractère en français	Name des Merkmals auf Deutsch	Nombre del carácter en español		
		states of expression	types d'expression	Ausprägungsstufen	tipos de expresión		

1 Characteristic number

2 (*) Asterisked characteristic – see Chapter 6.1.2

3 Type of expression
 QL Qualitative characteristic – see Chapter 6.3
 QN Quantitative characteristic – see Chapter 6.3
 PQ Pseudo-qualitative characteristic – see Chapter 6.3

4 Method of observation (and type of plot, if applicable)
 MG, MS, VG, VS – see Chapter 4.1.5

5 (+) See Explanations on the Table of Characteristics in Chapter 8.2

6 (a)-(f) See Explanations on the Table of Characteristics in Chapter 8.1

7 Not applicable

7. Table of Characteristics/Tableau des caractères/Merkmalstabelle/Tabla de caracteres

	English		français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
1.	QN	MG/VG	(a)				
	Plant: height						
	short						3
	medium					John Hurd's 98 Special	5
	tall						7
2. (*)	QN	VG	(a)				
	Plant: growth habit						
	erect						1
	semi erect					John Hurd's 98 Special	2
	prostrate						3
3. (*)	QN	MS/VG	(+)	(a)			
	Plant: number of primary lateral shoots						
	few						3
	medium						5
	many						7
4.	QN	MS/VG	(+)	(a)			
	Stem: internode length						
	short						3
	medium					John Hurd's 98 Special	5
	long						7

	English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
5.	QN MS/VG	(a), (f)				
	Stem: Thickness					
	thin					3
	medium					5
	thick					7
6.	QN VG	(a)				
	Stem: intensity of anthocyanin coloration					
	weak					3
	medium				John Hurd's 98 Special	5
	strong					7
7.	QN VG	(a)				
	Stem: intensity of green color					
	light					1
	medium					2
	dark					3
8.	QN VG	(+) (a)				
	Stem: Number of adventitious roots					
	few					3
	medium				Emerald	5
	many					7

	English		français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
9.	QN	VG	(a)				
	Stem: hairiness						
	absent or very weak					John Hurd's 98 Special	1
	weak						3
	medium						5
	strong						7
10.	QN	VG	(a)				
	Foliage: glossiness						
	weak					Boldrewood	3
	medium						5
	strong						7
11. (*)	QN	VG	(a)				
	Leaf: profile of terminal leaflet in cross-section						
	concave						1
	flat					Emerald	2
	convex						3
12. (*)	QN	MS/VG	(d), (e)				
	Leaf: length						
	short						3
	medium						5
	long						7

	English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
13. (*)	QN MS/VG	(d), (e)				
	Leaf: width					
	narrow					3
	medium					5
	broad					7
14.	QN VG	(a)				
	Leaf: intensity of anthocyanin coloration					
	absent or very weak					1
	weak					3
	medium					5
	strong				Emerald	7
15.	QN VG	(a)				
	Leaf: intensity of green color					
	light					3
	medium					5
	dark					7
16. (*)	QN MS/VG	(d), (e)				
	Leaf: length of terminal leaflet					
	short					3
	medium					5
	long					7

	English		français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
17. (*)	QN	MS/VG	(d), (e)				
	Leaf: width of terminal leaflet						
	narrow						3
	medium						5
	broad						7
18. (*)	PQ	VG	(+)	(d)			
	Leaf: shape of terminal leaflet						
	lanceolate						1
	ovate						2
	narrow elliptic						3
	medium elliptic						4
	circular						5
19. (*)	PQ	VG	(+)	(d)			
	Leaf: shape of apex of terminal leaflet						
	acute						1
	obtuse						2
	rounded						3
20. (*)	PQ	VG	(+)	(d)			
	Leaf: shape of base of terminal leaflet						
	obtuse						1
	truncate						2
	cordate						3

	English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
21. (*)	QN MS/VG	(d), (e)				
	Petiole: length from axil to first leaflet					
	short					3
	medium					5
	long					7
22. (*)	QN MS/VG	(+)	(b)			
	Time of beginning of flowering					
	early				Aqua	3
	medium				Emerald	5
	late					7
23. (*)	QN MS/VG	(+)	(b)			
	Proportion of plants with flowers					
	low				John Hurd's 98 Special	1
	medium				Emerald	3
	high				Aqua	5
24. (*)	QN MS/VG	(b)				
	Flower: diameter					
	small					3
	medium					5
	large					7


	English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
25. (*)	QN MS/VG	(c), (f)				
	Pedicel: length					
	short					3
	medium					5
	long					7
26. (*)	QN MS/VG	(c), (f)				
	Siliqua: length					
	short					3
	medium					5
	long					7
27. (*)	QN MS/VG	(c), (f)				
	Siliqua: width					
	narrow					3
	medium					5
	broad					7
28. (*)	QN VG	(c)				
	Siliqua: tendency to set seed					
	absent or very weak					1
	weak					3
	medium					5
	strong					7
29. (*)	QN VG	(+)	(c)			
	Seed: reticulation of surface					
	weak					3
	medium					5
	strong					7

8. Explanations on the Table of Characteristics

8.1 *Explanations covering several characteristics*


Characteristics containing the following key in the second column of the Table of Characteristics should be examined as indicated below:

- (a) observations should be made before flowering when leaves are fully developed
- (b) observations should be made on fully developed, fresh flowers if inflorescences develop.
- (c) observations should be made on fully developed pods at early stages of senescence if inflorescences have developed.
- (d) observations should be made before flowering when leaves are fully developed, on plants with excised axillary branches.
- (e) Characteristics for leaf and petiole length and width:



- Ad. 12: Leaf: length (A – B)
- Ad. 13: Leaf: width (C – D)
- Ad. 16: Leaf: length of terminal leaflet (A – E)
- Ad. 17: Leaf: width of terminal leaflet (F – G)
- Ad. 21: Petiole: length from axil to first leaflet (B – I)

(f) Characteristics for pedicel and siliqua lengths and widths:



Ad. 25: Pedicel: length (A – B)

Ad. 26: Siliqua: length (B – C)

Ad. 27: Siliqua: width (D)

8.2 Explanations for individual characteristics

Ad. 3: Plant: number of primary lateral shoots

Illustration to be added when available.

Ad. 4: Stem: internode length

If stem internode length is to be measured, this should be done in the central region of the stem where internode lengths are consistent.

Ad. 8: Stem: Number of adventitious roots




3 few




7 many


Ad. 18: Leaf: shape of terminal leaflet



Ad. 19: Leaf: shape of apex of terminal leaflet



Ad. 20: Leaf: shape of base of terminal leaflet



Ad. 22: Time of beginning of flowering

Time of beginning of flowering is defined as when 10% of the plants in a plot have fully open flowers.

Ad. 23: Proportion of plants with flowers

Assess at completion of flowering when no new flower buds are visible.

<i>Proportion</i>	<i>Note</i>	<i>Ranges (percentage)</i>
low	1	$\leq 5\%$
low to medium	2	6-35 %
medium	3	36-65 %
medium to high	4	66-95 %
high	5	$\geq 96\%$

Ad. 29: Seed: reticulation of surface

Observations should be done when no new flower buds develop



3 weak



7 strong

9. Literature

Bleasdale J.K.A. (1964). The flowering and growth of watercress (*Nasturtium officinale* R. Br.). J. Hort Sci. 39, 277-83.

Bleeker, W., Huthmann., Marion and Hurka, H. (1999). Evolution of hybrid tax in *Nasturtium* R. Br. (*Brassicaceae*). Folia Geobotanica. 34. 421 – 433.

Clapham, A.R., Tutin, T.G. and Warburg, E.F. (1981). Flora of the British Isles. Cambridge University Press. 3rd Edition 60 – 64.

Howard, H.W. and Manton, I. (1946). Autopolyploid and Allopolyploid Watercress with the description of a new species. Annals of Botany N.S. Vol. 10 No.37 1-16

Howard, H.W. and Lyon, A.G. (1952). Biological Flora of the British Isles. Journal of Ecology 40. 228-245.

Sheridan, G.E.C. (1996) Molecular studies of Watercress Phylogeny and the Crook-Root Pathogen. PhD thesis University of Bath (British Library Ref DX 205310).

Sheridan G.E.C., Claxton J.R., Clarkson J.M. and Blakesley D. (2001). Genetic diversity within commercial populations of watercress (*Rorippa nasturtium-aquaticum*), and between allied Brassicaceae inferred from RAPD-PCR. Euphytica 122 (2), 319-325.

Stevens, C.P. (1983). Watercress: production of the cultivated crop. ADAS/MAFF Reference Book 136. Grower Books. London.

10. Technical Questionnaire

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
		Application date: (not to be filled in by the applicant)
TECHNICAL QUESTIONNAIRE to be completed in connection with an application for plant breeders' rights		
1. Subject of the Technical Questionnaire		
1.1.1	Botanical name	<input type="text" value="Nasturtium xsterile (Airy Shaw) Oefelein"/> []
1.1.2	Common name	<input type="text"/>
1.2.1	Botanical name	<input type="text" value="Nasturtium microphyllum Boenn. ex Rchb."/> []
1.2.2	Common name	<input type="text" value="One-row watercress"/>
1.3.1	Botanical name	<input type="text" value="Nasturtium officinale R. Br."/> []
1.3.2	Common name	<input type="text" value="Watercress"/>
1.4.1	Botanical name	<input type="text" value="Nasturtium xsterile (Airy Shaw) Oefelein"/> []
1.4.2	Common name	<input type="text" value="Watercress"/>
2. Applicant		
	Name	<input type="text"/>
	Address	<input type="text"/>
	Telephone No.	<input type="text"/>
	Fax No.	<input type="text"/>
	E-mail address	<input type="text"/>
	Breeder (if different from applicant)	<input type="text"/>
3. Proposed denomination and breeder's reference		
	Proposed denomination (if available)	<input type="text"/>
	Breeder's reference	<input type="text"/>

#4. Information on the breeding scheme and propagation of the variety

4.1 Breeding scheme

Variety resulting from:

4.1.1 Crossing

(a) controlled cross []
(please state parent varieties)

(.....) x (.....)
female parent male parent

(b) partially known cross []
(please state known parent variety(ies))

(.....) x (.....)
female parent male parent

(c) unknown cross []

4.1.2 Mutation []
(please state parent variety)

4.1.3 Discovery and development []
(please state where and when discovered and how developed)

4.1.4 Other []
(please provide details)

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

4.2 Method of propagating the variety

4.2.1 Seed-propagated varieties

- (a) Self-pollination []
- (b) Cross-pollination []
- (c) []
- (d) Other (please provide details) []

4.2.2 Vegetative propagation

- (a) Cuttings []
- (b) *In vitro* propagation []
- (c) []
- (d) Other (state method) []

4.2.3 Other []
(Please provide details)

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

5. Characteristics of the variety to be indicated (the number in brackets refers to the corresponding characteristic in Test Guidelines; please mark the note which best corresponds).

Characteristics	Example Varieties	Note
5.1 Plant: growth habit (2)		
erect		1 []
semi erect	John Hurd's 98 Special	2 []
prostrate		3 []
5.2 Leaf: shape of terminal leaflet (18)		
lanceolate		1 []
ovate		2 []
narrow elliptic		3 []
medium elliptic		4 []
circular		5 []
5.3 Time of beginning of flowering (22)		
early	Aqua	3 []
medium	Emerald	5 []
late		7 []
5.4 Proportion of plants with flowers (23)		
low	John Hurd's 98 Special	1 []
medium	Emerald	3 []
high	Aqua	5 []
5.5 Siliqua: tendency to set seed (28)		
absent or very weak		1 []
weak		3 []
medium		5 []
strong		7 []

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

6. Similar varieties and differences from these varieties

Please use the following table and box for comments to provide information on how your candidate variety differs from the variety (or varieties) which, to the best of your knowledge, is (or are) most similar. This information may help the examination authority to conduct its examination of distinctness in a more efficient way.

Denomination(s) of variety(ies) similar to your candidate variety	Characteristic(s) in which your candidate variety differs from the similar variety(ies)	Describe the expression of the characteristic(s) for the similar variety(ies)	Describe the expression of the characteristic(s) for your candidate variety
<i>Example</i>	<i>Plant: growth habit</i>	<i>erect</i>	<i>prostrate</i>
Comments:			

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

#7. Additional information which may help in the examination of the variety

7.1 In addition to the information provided in sections 5 and 6, are there any additional characteristics which may help to distinguish the variety?

Yes No

(If yes, please provide details)

7.2 Are there any special conditions for growing the variety or conducting the examination?

Yes No

(If yes, please provide details)

7.3 Other information

A representative color photograph of the variety displaying its main distinguishing feature(s), should accompany the Technical Questionnaire. The photograph will provide a visual illustration of the candidate variety which supplements the information provided in the Technical Questionnaire.

The key points to consider when taking a photograph of the candidate variety are:

- Indication of the date and geographic location
- Correct labeling (breeder's reference)
- Good quality printed photograph (minimum 10 cm x 15 cm) and/or sufficient resolution electronic format version (minimum 960 x 1280 pixels)

Further guidance on providing photographs with the Technical Questionnaire is available in document TGP/7 "Development of Test Guidelines", Guidance Note 35 (<http://www.upov.int/tgp/en/>).

[The link provided may be deleted by members of the Union when developing authorities' own test guidelines.]

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

8. Authorization for release

(a) Does the variety require prior authorization for release under legislation concerning the protection of the environment, human and animal health?

Yes No

(b) Has such authorization been obtained?

Yes No

If the answer to (b) is yes, please attach a copy of the authorization.

9. Information on plant material to be examined or submitted for examination

9.1 The expression of a characteristic or several characteristics of a variety may be affected by factors, such as pests and disease, chemical treatment (e.g. growth retardants or pesticides), effects of tissue culture, different rootstocks, scions taken from different growth phases of a tree, etc.

9.2 The plant material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If the plant material has undergone such treatment, full details of the treatment must be given. In this respect, please indicate below, to the best of your knowledge, if the plant material to be examined has been subjected to:

(a)	Microorganisms (e.g. virus, bacteria, phytoplasma)	Yes <input type="checkbox"/>	No <input type="checkbox"/>
(b)	Chemical treatment (e.g. growth retardant, pesticide)	Yes <input type="checkbox"/>	No <input type="checkbox"/>
(c)	Tissue culture	Yes <input type="checkbox"/>	No <input type="checkbox"/>
(d)	Other factors	Yes <input type="checkbox"/>	No <input type="checkbox"/>

Please provide details for where you have indicated "yes".

.....

9.3 Has the plant material to be examined been tested for the presence of virus or other pathogens?

Yes

(please provide details as specified by the Authority)

No

10. I hereby declare that, to the best of my knowledge, the information provided in this form is correct:

Applicant's name

Signature Date

[End of document]