

Technical Working Party for Vegetables

TWV/51/8

Fifty-First Session

Roelofarendsveen, Netherlands, July 3 to 7, 2017

Original: English

Date: June 14, 2017

PARTIAL REVISION OF THE TEST GUIDELINES FOR SPINACH (DOCUMENT TG/55/7 REV. 4)

Document prepared by the Netherlands

Disclaimer: this document does not represent UPOV policies or guidance

1. The purpose of this document is to present a proposal for a partial revision of the Test Guidelines for Spinach (document TG/55/7 Rev. 4).
2. The Technical Working Party for Vegetables (TWV), at its fiftieth session, held in Brno, Czech Republic, from June 27 to July 1, 2016, agreed that the Test Guidelines for Spinach (document TG/55/7 Rev. 4) be partially revised for Characteristic 18 “Resistance to Peronospora farinosa f. sp. spinaciae (Pfs) (see document TWV/50/25 “Report”, Annex IV).
3. The following changes are proposed:
 - (a) Addition of “Race Pfs: 16” to Characteristic 18 “Resistance to Peronospora farinosa f. sp. spinaciae (Pfs)”, including example varieties;
 - (b) Revision of explanation Ad. 18 in Chapter 8.2 “Explanations for individual characteristics”
 - (c) Addition of “Race Pfs: 16” to TQ 7.3 “Other information”
4. The proposed changes are presented below in highlight and underline (insertion) and ~~strikethrough~~ (deletion).

Proposal to add “Race Pfs: 16” to Characteristic 18 “Resistance to Peronospora farinosa f. sp. spinaciae (Pfs)”

5.	VG	Resistance to <u>Peronospora farinosa</u> f. sp. <u>spinaciae</u> (Pfs)	Résistance à <u>Peronospora farinosa</u> f. sp. <u>spinaciae</u> (Pfs)	Resistenz gegen <u>Peronospora farinosa</u> f. sp. <u>spinaciae</u> (Pfs)	Resistencia a <u>Peronospora farinosa</u> f. sp. <u>spinaciae</u> (Pfs)		
18.1	Race Pfs: 1		Race Pfs: 1	Pathotyp Pfs: 1	Raza Pfs: 1		
QL	absent	absente	fehrend	ausente	Viroflay, Winterreuzen		1
	present	présente	vorhanden	presente	Califlay, Resistoflay		9
18.2	Race Pfs: 2		Race Pfs: 2	Pathotyp Pfs: 2	Raza Pfs: 2		
QL	absent	absente	fehrend	ausente	Califlay		1
	present	présente	vorhanden	presente	Resistoflay		9

18.3	Race Pfs: 3	Race Pfs: 3	Pathotyp Pfs: 3	Raza Pfs: 3		
QL	absent	absente	fehlend	ausente	Resistoflay	1
	present	présente	vorhanden	presente	Califlay, Clermont	9

18.4	Race Pfs: 4	Race Pfs: 4	Pathotyp Pfs: 4	Raza Pfs: 4		
QL	absent	absente	fehlend	ausente	Califlay	1
	present	présente	vorhanden	presente	Clermont	9

18.5	Race Pfs: 5	Race Pfs: 5	Pathotyp Pfs: 5	Raza Pfs: 5		
QL	absent	absente	fehlend	ausente	Clermont	1
	present	présente	vorhanden	presente	Califlay, Campania	9

18.6	Race Pfs: 6	Race Pfs: 6	Pathotyp Pfs: 6	Raza Pfs: 6		
QL	absent	absente	fehlend	ausente	Califlay, Campania	1
	present	présente	vorhanden	presente	Boeing	9

18.7	Race Pfs: 7	Race Pfs: 7	Pathotyp Pfs: 7	Raza Pfs: 7		
QL	absent	absente	fehlend	ausente	Califlay	1
	present	présente	vorhanden	presente	Campania	9

18.8	Race Pfs: 8	Race Pfs: 8	Pathotyp Pfs: 8	Raza Pfs: 8		
QL	absent	absente	fehlend	ausente	Boeing, Campania	1
	present	présente	vorhanden	presente	Lazio, Lion	9

18.9	Race Pfs: 10	Race Pfs: 10	Pathotyp Pfs:10	Raza Pfs:10		
QL	absent	absente	fehlend	ausente	Boeing, Campania, Lion	1
	present	présente	vorhanden	presente	Lazio	9

18.10	Race Pfs: 11	Race Pfs: 11	Pathotyp Pfs: 11	Raza Pfs: 11		
QL	absent	absente	fehlend	ausente	Lazio	1
	present	présente	vorhanden	presente	Boeing, Califlay, Campania, Lion	9

18.11	Race Pfs: 12	Race Pfs: 12	Pathotyp Pfs: 12	Raza Pfs: 12		
QL	absent	absente	fehlend	ausente	Boeing, Campania	1
	present	présente	vorhanden	presente	Finch, Pigeon, Red Kitten, Zebu	9
18.12	Race Pfs: 13	Race Pfs: 13	Pathotyp Pfs: 13	Raza Pfs: 13		
QL	absent	absente	fehlend	ausente	Campania	1
	present	présente	vorhanden	presente	Boeing, Lion	9
18.13	Race Pfs: 14	Race Pfs: 14	Pathotyp Pfs: 14	Raza Pfs: 14		
QL	absent	absente	fehlend	ausente	Campania, Pigeon	1
	present	présente	vorhanden	presente	Califlay, Lion	9
18.14	Race Pfs: 15	Race Pfs: 15	Pathotyp Pfs: 15	Raza Pfs: 15		
QL	absent	absente	fehlend	ausente	Caladonia	1
	present	présente	vorhanden	presente	Pigeon	9
<u>18.15</u>	<u>Race Pfs: 16</u>	<u>Race Pfs: 16</u>	<u>Pathotyp Pfs: 16</u>	<u>Raza Pfs: 16</u>		
<u>QL</u>	<u>absent</u>	<u>absente</u>	<u>fehlend</u>	<u>ausente</u>	<u>Meerkat</u>	<u>1</u>
	<u>present</u>	<u>présente</u>	<u>vorhanden</u>	<u>presente</u>	<u>Caladonia</u>	<u>9</u>

Proposal to revise explanation Ad. 18 in Chapter 8.2 "Explanations for individual characteristics"

Ad. 18: Resistance to *Peronospora farinosa* f. sp. *spinaciae*

Maintenance of races

Type of medium: Living host plants, obtainable from:
Naktuinbouw
P.O. Box 40
NL-2370 AA Roelofarendsveen
Netherlands
www.naktuinbouw.com
or plant material with spores stored at -20° C for a maximum of one year

Execution of test

Growth stage of plants: First cotyledons/leaf, eleven-day-old plants

Temperature: 15°C during day/12°C during night

Light: 15 hours per day, after emergence

Growing method: In soil in pots or trays in a glasshouse or growth chamber

Method of inoculation: Sporulating leaves, taken from host plants that were infected seven days before, are thoroughly rinsed with sterile tap water (maximum 150 ml water per 224 plants). The spore suspension is filtered through cheesecloth and sprayed on test plants until the inoculum covers the leaves but does not run off. 150 ml of suspension is enough for up to 3 x 224 plants. Spore density should be 20,000 to 100,000 conidia/ml water. The spore suspension should be used fresh.

Remarks: Spinach downy mildew is wind-borne. Sporulating plants should be kept in closed containers or isolated chambers to prevent any cross-contamination. Resistant controls are needed in each multiplication and in each test to ensure the race identity.

Light and humidity conditions during seedling development and incubation are critical. Optimal humidity of approximately 80-90% RH allows plant growth and fungal growth; strong light inhibits spore germination and infection.

The test should be carried out in wintertime with protection against direct sunshine. After inoculation, the plants should remain under plastic for three days. After this time, the plastic should be slightly raised during the daytime.

Duration of test

- Multiplication harvest spores 7 days after inoculation
- Sowing to inoculation: 11 days
- Inoculation to reading: 10 days

Number of plants tested ~~56 plants~~ at least 20 plants

Evaluation of infection: Resistance is usually complete; sometimes necrotic spots are visible as a result of infection. Susceptible plants show varying degrees of sporulation. Sporulation is visible as a grey covering on leaves, starting on the more humid abaxial side.

Differential varieties to identify races

Races Pfs: 1-8 and 10-16 of *Peronospora farinosa* f. sp. *spinaciae* are defined with a standard set of “differential varieties” according to the following table:

Differential variety	Pfs: 1	Pfs: 2	Pfs: 3	Pfs: 4	Pfs: 5	Pfs: 6	Pfs: 7	Pfs: 8	Pfs: 10	Pfs: 11	Pfs: 12	Pfs: 13	Pfs: 14	Pfs: 15	Pfs: 16
Vioflay	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S
Resistoflay	R	R	S	S	S	S	S	S	S	S	S	S	S	S	S
Califlay	R	S	R	S	R	S	S	R	S	R	R	S	R	S	R
Clermont	R	R	R	R	S	S	S	S	S	S	S	S	S	R	S
Campania	R	R	R	R	R	S	R	S	S	R	S	S	S	R	R
Boeing	R	R	R	R	R	R	R	S	S	R	S	R	S	R	R
Lion	R	R	R	R	R	R	R	R	S	R	R	R	R	R	R
Lazio	R	R	R	R	R	R	R	R	R	S	S	S	S	R	S
Whale	R	R	R	R	R	R	R	R	S	R	R	S	R	S	R
Pigeon	R	R	R	R	R	R	R	R	R	R	R	R	S	R	S
Caladonia	R	R	R	R	R	R	R	R	R	R	R	R	R	S	R
Meerkat	R	R	R	R	R	R	R	R	R	R	R	R	R	R	S

Legend: R= resistance present; S = resistance absent, susceptible

Proposal to add “Race Pfs: 16” to TQ 7.3 “Other information”

“7.3 Other information

[...]

“(b) Resistance to pests and diseases (specify)

“(i) Resistance to *Peronospora farinosa* f. sp. *spinaciae*

Race Pfs: 1	[] absent	[] present
Race Pfs: 2	[] absent	[] present
Race Pfs: 3	[] absent	[] present
Race Pfs: 4	[] absent	[] present
Race Pfs: 5	[] absent	[] present
Race Pfs: 6	[] absent	[] present
Race Pfs: 7	[] absent	[] present
Race Pfs: 8	[] absent	[] present
Race Pfs: 10	[] absent	[] present
Race Pfs: 11	[] absent	[] present
Race Pfs: 12	[] absent	[] present
Race Pfs: 13	[] absent	[] present
Race Pfs: 14	[] absent	[] present
Race Pfs: 15	[] absent	[] present
Race Pfs: 16	[] absent	[] present

[...]”