



TG/136/6(proj.2)

ORIGINAL: English

DATE: 2024-03-09

INTERNATIONAL UNION FOR THE PROTECTION OF NEW VARIETIES OF PLANTS

Geneva

DRAFT

PARSLEY

UPOV Code(s): PETRO_CRI

Petroselinum crispum (Mill.) Fuss

GUIDELINES

FOR THE CONDUCT OF TESTS

FOR DISTINCTNESS, UNIFORMITY AND STABILITY

*prepared by experts from Germany
to be considered by the
Technical Working Party for Vegetables
at its fifty-eighth session, to be held virtually
from 2024-04-22 to 2024-04-25*

Disclaimer: this document does not represent UPOV policies or guidance

Alternative names:*

<i>Botanical name</i>	<i>English</i>	<i>French</i>	<i>German</i>	<i>Spanish</i>
<i>Petroselinum crispum</i> (Mill.) Fuss	Parsley	Persil	Petersilie	Perejil

The purpose of these guidelines ("Test Guidelines") is to elaborate the principles contained in the General Introduction (document TG/1/3), and its associated TGP documents, into detailed practical guidance for the harmonized examination of distinctness, uniformity and stability (DUS) and, in particular, to identify appropriate characteristics for the examination of DUS and production of harmonized variety descriptions.

ASSOCIATED DOCUMENTS

These Test Guidelines should be read in conjunction with the General Introduction and its associated TGP documents.

* These names were correct at the time of the introduction of these Test Guidelines but may be revised or updated. [Readers are advised to consult the UPOV Code, which can be found on the UPOV Website (www.upov.int), for the latest information.]

TABLE OF CONTENTS	PAGE
1. SUBJECT OF THESE TEST GUIDELINES.....	3
2. MATERIAL REQUIRED.....	3
3. METHOD OF EXAMINATION.....	3
3.1 Number of Growing Cycles.....	3
3.2 Testing Place.....	3
3.3 Conditions for Conducting the Examination.....	3
3.4 Test Design.....	4
3.5 Additional Tests.....	4
4. ASSESSMENT OF DISTINCTNESS, UNIFORMITY AND STABILITY.....	4
4.1 Distinctness.....	4
4.2 Uniformity.....	5
4.3 Stability.....	5
5. GROUPING OF VARIETIES AND ORGANIZATION OF THE GROWING TRIAL.....	6
6. INTRODUCTION TO THE TABLE OF CHARACTERISTICS.....	6
6.1 Categories of Characteristics.....	6
6.2 States of Expression and Corresponding Notes.....	6
6.3 Types of Expression.....	6
6.4 Example Varieties.....	7
6.5 Legend.....	8
7. TABLE OF CHARACTERISTICS/TABLEAU DES CARACTÈRES/MERKMALSTABELLE/TABLA DE CARACTERES.....	9
8. EXPLANATIONS ON THE TABLE OF CHARACTERISTICS.....	19
8.1 Explanations covering several characteristics.....	19
8.2 Explanations for individual characteristics.....	20
9. LITERATURE.....	24
10 TECHNICAL QUESTIONNAIRE.....	25

1. Subject of these Test Guidelines

These Test Guidelines apply to all varieties of *Petroselinum crispum* (Mill.) Fuss.

2. Material Required

2.1 The competent authorities decide on the quantity and quality of the plant material required for testing the variety and when and where it is to be delivered. Applicants submitting material from a State other than that in which the testing takes place must ensure that all customs formalities and phytosanitary requirements are complied with.

2.2 The material is to be supplied in the form of seed.

2.3 The minimum quantity of plant material, to be supplied by the applicant, should be:

Leaf parsley: 12,000 seeds
Root parsley: 24,000 seeds.

The seed should meet the minimum requirements for germination, species and analytical purity, health and moisture content, specified by the competent authority. In cases where the seed is to be stored, the germination capacity should be as high as possible and should, be stated by the applicant.

2.4 The plant material supplied should be visibly healthy, not lacking in vigor, nor affected by any important pest or disease.

2.5 The plant material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If it has been treated, full details of the treatment must be given.

3. Method of Examination

3.1 *Number of Growing Cycles*

3.1.1 The minimum duration of tests should normally be two independent growing cycles.

3.1.2 The two independent growing cycles should be in the form of two separate plantings.

3.1.3 The testing of a variety may be concluded when the competent authority can determine with certainty the outcome of the test.

3.2 *Testing Place*

Tests are normally conducted at one place. In the case of tests conducted at more than one place, guidance is provided in TGP/9 "Examining Distinctness".

3.3 *Conditions for Conducting the Examination*

The tests should be carried out under conditions ensuring satisfactory growth for the expression of the relevant characteristics of the variety and for the conduct of the examination.

3.4 *Test Design*

3.4.1 Each test should be designed to result in at least 60 plants for leaf parsley and 160 plants for root parsley, which should be divided between at least 2 replicates.

3.4.2 The design of the tests should be such that plants or parts of plants may be removed for measurement or counting without prejudice to the observations which must be made up to the end of the growing cycle.

3.5 *Additional Tests*

Additional tests, for examining relevant characteristics, may be established.

4. Assessment of Distinctness, Uniformity and Stability

4.1 *Distinctness*

4.1.1 General Recommendations

It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding distinctness. However, the following points are provided for elaboration or emphasis in these Test Guidelines.

4.1.2 Consistent Differences

The differences observed between varieties may be so clear that more than one growing cycle is not necessary. In addition, in some circumstances, the influence of the environment is not such that more than a single growing cycle is required to provide assurance that the differences observed between varieties are sufficiently consistent. One means of ensuring that a difference in a characteristic, observed in a growing trial, is sufficiently consistent is to examine the characteristic in at least two independent growing cycles.

4.1.3 Clear Differences

Determining whether a difference between two varieties is clear depends on many factors, and should consider, in particular, the type of expression of the characteristic being examined, i.e. whether it is expressed in a qualitative, quantitative, or pseudo-qualitative manner. Therefore, it is important that users of these Test Guidelines are familiar with the recommendations contained in the General Introduction prior to making decisions regarding distinctness.

4.1.4 Number of Plants or Parts of Plants to be Examined

Unless otherwise indicated, for the purposes of distinctness, all observations on single plants should be made on 40 plants or parts of plants taken from each of 40 plants and any other observations made on all plants in the test, disregarding any off-type plants.

In the case of observations of parts taken from single plants, the number of parts to be taken from each of the plants should be 1.

4.1.5 Method of Observation

The recommended method of observing the characteristic for the purposes of distinctness is indicated by the following key in the Table of Characteristics (see document TGP/9 "Examining Distinctness", Section 4 "Observation of characteristics"):

MG: single measurement of a group of plants or parts of plants

MS: measurement of a number of individual plants or parts of plants

VG: visual assessment by a single observation of a group of plants or parts of plants

VS: visual assessment by observation of individual plants or parts of plants

Type of observation: visual (V) or measurement (M)

"Visual" observation (V) is an observation made on the basis of the expert's judgment. For the purposes of this document, "visual" observation refers to the sensory observations of the experts and, therefore, also includes smell, taste and touch. Visual observation includes observations where the expert uses reference points (e.g. diagrams, example varieties, side-by-side comparison) or non-linear charts (e.g. color charts). Measurement (M) is an objective observation against a calibrated, linear scale e.g. using a ruler, weighing scales, colorimeter, dates, counts, etc.

Type of record: for a group of plants (G) or for single, individual plants (S)

For the purposes of distinctness, observations may be recorded as a single record for a group of plants or parts of plants (G), or may be recorded as records for a number of single, individual plants or parts of plants (S). In most cases, "G" provides a single record per variety and it is not possible or necessary to apply statistical methods in a plant-by-plant analysis for the assessment of distinctness.

In cases where more than one method of observing the characteristic is indicated in the Table of Characteristics (e.g. VG/MG), guidance on selecting an appropriate method is provided in document TGP/9, Section 4.2.

4.2 *Uniformity*

- 4.2.1 It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding uniformity. However, the following points are provided for elaboration or emphasis in these Test Guidelines:
- 4.2.2 These Test Guidelines have been developed for the examination of seed-propagated varieties. For varieties with other types of propagation, the recommendations in the General Introduction and document TGP/13 "Guidance for new types and species" Section 4.5 "Testing Uniformity" should be followed.
- 4.2.3 The assessment of uniformity should be according to the recommendations for cross-pollinated varieties in the General Introduction.
- 4.2.4 For the assessment of uniformity of seed-propagated varieties, a population standard of 2% and an acceptance probability of at least 95% should be applied. In the case of a sample size of 40 plants, 2 off-types are allowed.

4.3 *Stability*

- 4.3.1 In practice, it is not usual to perform tests of stability that produce results as certain as those of the testing of distinctness and uniformity. However, experience has demonstrated that, for many types of variety, when a variety has been shown to be uniform, it can also be considered to be stable.
- 4.3.2 Where appropriate, or in cases of doubt, stability may be further examined by testing a new seed stock to ensure that it exhibits the same characteristics as those shown by the initial material supplied.

5. Grouping of Varieties and Organization of the Growing Trial

- 5.1 The selection of varieties of common knowledge to be grown in the trial with the candidate varieties and the way in which these varieties are divided into groups to facilitate the assessment of distinctness are aided by the use of grouping characteristics.
- 5.2 Grouping characteristics are those in which the documented states of expression, even where produced at different locations, can be used, either individually or in combination with other such characteristics: (a) to select varieties of common knowledge that can be excluded from the growing trial used for examination of distinctness; and (b) to organize the growing trial so that similar varieties are grouped together.
- 5.3 The following have been agreed as useful grouping characteristics:
- (a) Leaf blade: curling (characteristic 6)
 - (b) Root: thickening of main root (characteristic 20)
- Grouping for varieties of Root and Leaf Parsley is based on characteristic 20: Root: thickening of main root:
- absent: Leaf parsley
 - present: Root parsley
- 5.4 Guidance for the use of grouping characteristics, in the process of examining distinctness, is provided through the General Introduction and document TGP/9 "Examining Distinctness".

6. Introduction to the Table of Characteristics

6.1 *Categories of Characteristics*

6.1.1 Standard Test Guidelines Characteristics

Standard Test Guidelines characteristics are those which are approved by UPOV for examination of DUS and from which members of the Union can select those suitable for their particular circumstances.

6.1.2 Asterisked Characteristics

Asterisked characteristics (denoted by *) are those included in the Test Guidelines which are important for the international harmonization of variety descriptions and should always be examined for DUS and included in the variety description by all members of the Union, except when the state of expression of a preceding characteristic or regional environmental conditions render this inappropriate.

6.2 States of Expression and Corresponding Notes

6.2.1 States of expression are given for each characteristic to define the characteristic and to harmonize descriptions. Each state of expression is allocated a corresponding numerical note for ease of recording of data and for the production and exchange of the description.

6.2.2 All relevant states of expression are presented in the characteristic.

6.2.3 Further explanation of the presentation of states of expression and notes is provided in document TGP/7 "Development of Test Guidelines".

6.3 Types of Expression

An explanation of the types of expression of characteristics (qualitative, quantitative and pseudo-qualitative) is provided in the General Introduction.

6.4 Example Varieties

Where appropriate, example varieties are provided to clarify the states of expression of each characteristic.

6.5 Legend

English		français		deutsch		español		Example Varieties Exemples Beispielssorten Variedades ejemplo		Note/ Nota
1	2	3	4	5	6	7				
Name of characteristics in English		Nom du caractère en français		Name des Merkmals auf Deutsch		Nombre del carácter en español				
states of expression		types d'expression		Ausprägungsstufen		tipos de expresión				

- | | | | |
|---|---|---|---------------------|
| 1 | Characteristic number | | |
| 2 | (*) | Asterisked characteristic | – see Chapter 6.1.2 |
| 3 | Type of expression | | |
| | QL | Qualitative characteristic | – see Chapter 6.3 |
| | QN | Quantitative characteristic | – see Chapter 6.3 |
| | PQ | Pseudo-qualitative characteristic | – see Chapter 6.3 |
| 4 | Method of observation (and type of plot, if applicable) | | |
| | MG, MS, VG, VS | | – see Chapter 4.1.5 |
| 5 | (+) | See Explanations on the Table of Characteristics in Chapter 8.2 | |
| 6 | (a)-(b) | See Explanations on the Table of Characteristics in Chapter 8.1 | |
| 7 | Not applicable | | |

7. Table of Characteristics/Tableau des caractères/Merkmaltabelle/Tabla de caracteres

	English		français		deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
1.	QN	VG	(+)	(a)				
	Plant: growth habit							
		erect					Laura (FL), Titan (CL)	1
		erect to semi erect					Gigante d'Italia (FL)	2
		semi erect					Grüne Perle (CL)	3
		semi erect to prostrate						4
		prostrate						5
2. (*)	QN	MS/VG		(a)				
	Plant: height							
		very short						1
		very short to short						2
		short						3
		short to medium					Grüne Perle (CL)	4
		medium					Kudrnka (CL), Titan (CL)	5
		medium to tall						6
		tall					Natalka (FL)	7
		tall to very tall					Monica (R)	8
		very tall					Gigante d'Italia (FL)	9
3.	QN	MS/VG		(a)				
	Plant: width							
		very narrow						1
		very narrow to narrow						2
		narrow					Afrodite (CL)	3
		narrow to medium					Grüne Perle (CL)	4
		medium					Titan (CL)	5
		medium to broad					Laura (FL)	6
		broad						7
		broad to very broad						8
		very broad						9

	English		français		deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
4.	(*)	QN	VG	(+)	(a)			
		Plant: density of foliage						
			very loose					1
			very loose to loose					2
			loose				Gigante d'Italia (FL), Titan (CL)	3
			loose to medium				Laica (FL), Laura (FL)	4
			medium				Ines (CL)	5
			medium to dense				Bravour (CL), Kudrnka (CL)	6
			dense					7
			dense to very dense				Grüne Perle (CL), Lisette (CL)	8
			very dense				Afrodite (CL)	9
5.		QN	VG	(+)				
		Plant: number of petioles						
			very few					1
			few				Bravour (CL), Ines (CL), Starlett (CL)	2
			medium					3
			many				Gigante d'Italia (FL)	4
			very many					5
6.	(*)	QL	VG		(a)			
		Leaf blade: curling						
			absent				Gigante d'Italia (FL), Greenso (FL)	1
			present				Grüne Perle (CL), Titan (CL)	9

	English		français		deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
7.	(*)	QN	VG	(+)	(a)			
		<u>Only varieties with leaf blade curling:</u> Leaf blade: intensity of curling						
			very weak				Titan (CL)	1
			very weak to weak					2
			weak				Kudrnka (CL)	3
			weak to medium					4
			medium				Ines (CL)	5
			medium to strong					6
			strong				Emma (CL)	7
			strong to very strong				Grüne Perle (CL)	8
			very strong					9
8.		QN	VG		(a)			
		<u>Only varieties with leaf blade curling:</u> Leaf: appearance of surface of canopy						
			very open					1
			very open to open					2
			open					3
			open to medium					4
			medium				Ines (CL)	5
			medium to close				Emma (CL)	6
			close				Gusti (CL), Starlett (CL)	7
			close to very close					8
			very close					9
9.		QL	VG	(+)	(a)			
		<u>Only varieties with leaf blade curling:</u> Leaf blade: upward reflexing of lobes						
			absent					1
			present				Grüne Perle (CL), Lisette (CL), Thujade (CL), Titan (CL)	9

	English		français		deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
10	(*)	QN	MS/VG	(+)	(a)			
		Leaf blade: length						
		very short						1
		very short to short						2
		short				Afrodite (CL)		3
		short to medium				Lisette (CL), Titan (CL)		4
		medium				Laura (FL), Lion (FL)		5
		medium to long				Gigante d'Italia (FL)		6
		long				Laica (FL)		7
		long to very long						8
		very long						9
11	(*)	QN	MS/VG	(+)	(a)			
		Leaf blade: width						
		very narrow						1
		very narrow to narrow						2
		narrow						3
		narrow to medium				Titan (CL)		4
		medium				Emma (CL), Lion (FL)		5
		medium to broad				Laura (FL)		6
		broad				Laica (FL)		7
		broad to very broad						8
		very broad						9
12		QN	MS/VG		(a)			
		Leaf blade: ratio length/width						
		very low						1
		very low to low						2
		low						3
		low to medium				Grüne Perle (CL)		4
		medium				Laica (FL), Laura (FL), Starlett (CL)		5
		medium to high						6
		high				Gigante d'Italia (FL)		7
		high to very high						8
		very high						9

	English		français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
13	(*)	QN	VG	(a)			
		Leaf blade: intensity of green color					
		very light					1
		very light to light					2
		light					3
		light to medium				Bravour (CL)	4
		medium				Lisette (CL)	5
		medium to dark				Emma (CL), Starlett (CL)	6
		dark				Ines (CL), Laica (FL), Menuette (CL)	7
		dark to very dark				Lion (FL)	8
		very dark				Greenso (FL), Titan (CL)	9
14		QN	MS/VG	(+)	(a)		
		Leaf blade: distance between 1st and 2nd pair of leaflets					
		very short					1
		very short to short					2
		short				Afrodite (CL)	3
		short to medium				Grüne Perle (CL), Ines (CL)	4
		medium				Lisette (CL), Titan (CL)	5
		medium to long					6
		long				Laura (FL)	7
		long to very long					8
		very long					9
15	(*)	QN	VG	(+)	(a)		
		Only varieties with leaf blade curling; leaf blade: depth of incision					
		shallow				Grüne Perle (CL), Lisette (CL)	1
		medium					2
		deep				Menuette (CL)	3

	English		français		deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
16	QN	VG	(+)	(a)				
	<u>Only varieties without leaf blade curling:</u> leaflet: shape							
		narrow triangular					Gigante d'Italia (FL)	1
		medium triangular						2
		broad triangular						3
17	QN	MS/VG	(+)	(a)				
	Petiole: length							
		very short						1
		very short to short						2
		short					Ines (CL)	3
		short to medium					Emma (CL), Grüne Perle (CL)	4
		medium					Laura (FL)	5
		medium to long						6
		long						7
		long to very long					Monica (R)	8
		very long						9
18	QN	MS/VG	(+)	(a)				
	Petiole: thickness							
		thin						1
		thin to medium						2
		medium					Emma (CL), Menuette (CL)	3
		medium to thick						4
		thick						5
19 (*)	QN	VG		(a)				
	Petiole: anthocyanin coloration							
		absent or very weak					Grüne Perle (CL)	1
		weak					Natalka (FL)	2
		medium						3
		strong						4
		very strong						5

	English		français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
20	(*)	QL	VG	(b)			
		Root: thickening of main root					
		absent				Menuette (CL)	1
		present				Halblange (R)	9
21	(*)	QN	MS/VG	(b)			
		Only root parsley varieties: Root: length					
		very short				Halblange (R)	1
		very short to short					2
		short					3
		short to medium					4
		medium				A grosse racine gros hatif (R)	5
		medium to long				Arat (R)	6
		long				Artica (R)	7
		long to very long					8
		very long					9
22	(*)	QN	MS/VG	(b)			
		Only root parsley varieties: Root: width					
		narrow					1
		narrow to medium				Artica (R)	2
		medium				A grosse racine gros hatif (R), Arat (R)	3
		medium to broad					4
		broad					5
23	(*)	QN	MS/VG	(b)			
		Only root parsley varieties: Root: ratio length/width					
		small					1
		small to medium					2
		medium					3
		medium to large				Artica (R)	4
		large					5

	English		français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
24	QN	VG	(b)				
	<u>Only root parsley varieties: Root: branching</u>						
		very weak					1
		very weak to weak					2
		weak				A grosse racine gros hatif (R)	3
		weak to medium				Arat (R), Artica (R)	4
		medium					5
		medium to strong					6
		strong					7
		strong to very strong					8
		very strong					9

8. Explanations on the Table of Characteristics

8.1 *Explanations covering several characteristics*

Characteristics containing the following key in the Table of Characteristics should be examined as indicated below:


- (a) Observations should be made at the time of full development of the foliage.
- (b) Observations should be made at the time of full development of the roots.

8.2 *Explanations for individual characteristics*

Ad. 1: Plant: growth habit

Observations should be made visually of the angle between the outer petioles and the soil surface.

- 1 = erect
- 2 = semi-erect
- 3 = intermediate
- 4 = semi-prostrate
- 5 = prostrate




Ad. 4: Plant: density of foliage

Observations should be made from above, looking at the foliage in total.


Ad. 5: Plant: number of petioles

Observation should be made at the end of the trial by cutting the whole plant below the first branches.


Ad. 7: Only varieties with leaf blade curling: Leaf blade: intensity of curling



Ad. 9: Only varieties with leaf blade curling: Leaf blade: upward reflexing of lobes



Ad. 10: Leaf blade: length



Ad. 11: Leaf blade: width

See Ad. 10

Ad. 14: Leaf blade: distance between 1st and 2nd pair of leaflets

See Ad. 10

Ad. 15: Only varieties with leaf blade curling: leaf blade: depth of incision

Observation on the depth of incision can be made at any leaflet of the leaf blade.



1
shallow



2
medium



3
deep

Ad. 16: Only varieties without leaf blade curling: leaflet: shape



1
narrow triangular



2
medium triangular



3
broad triangular

Ad. 17: Petiole: length

See Ad. 10

Ad. 18: Petiole: thickness

See Ad. 10

9. Literature

Vogel, G., 1996: Handbuch des speziellen Gemüsebaus. Ulmer Verlag, Stuttgart, Seiten 1009 - 1026.

10. Technical Questionnaire

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

	Application date: (not to be filled in by the applicant)
--	---

TECHNICAL QUESTIONNAIRE
to be completed in connection with an application for plant breeders' rights

1. Subject of the Technical Questionnaire

1.1 Botanical name

1.2 Common name

2. Applicant

Name

Address

Telephone No.

Fax No.

E-mail address

Breeder (if different from applicant)

3. Proposed denomination and breeder's reference

Proposed denomination (if available)

Breeder's reference

#4. Information on the breeding scheme and propagation of the variety

4.1 Breeding scheme

Variety resulting from:

4.1.1 Crossing

(a) controlled cross []
(please state parent variety)
(.....) x (.....)
female parent male parent

(b) partially known cross []
(please state known parent variety(ies))
(.....) x (.....)
female parent male parent

(c) unknown cross []

4.1.2 Mutation []
(please state parent variety)

4.1.3 Discovery and development []
(please state where and when discovered and how developed)

4.1.4 Other []
(Please provide details)

Authorities may allow certain of this information to be provided in a confidential section of the Technical Questionnaire.

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

4.2 Method of propagating the variety	
4.2.1 Seed-propagated varieties	
(a) Cross-pollination	[]
(b) Other (please provide details)	[]
<input type="text"/>	
4.2.2 Other (Please provide details)	[]
<input type="text"/>	

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

5. Characteristics of the variety to be indicated (the number in brackets refers to the corresponding characteristic in Test Guidelines; please mark the note which best corresponds).

Characteristics	Example Varieties	Note
5.1 Plant: height (2)		
very short		1 []
very short to short		2 []
short		3 []
short to medium	Grüne Perle (CL)	4 []
medium	Kudrnka (CL), Titan (CL)	5 []
medium to tall		6 []
tall	Natalka (FL)	7 []
tall to very tall	Monica (R)	8 []
very tall	Gigante d'Italia (FL)	9 []
5.2 Leaf blade: curling (6)		
absent	Gigante d'Italia (FL), Greenso (FL)	1 []
present	Grüne Perle (CL), Titan (CL)	9 []
5.3 Leaf blade: intensity of green color (13)		
very light		1 []
very light to light		2 []
light		3 []
light to medium	Bravour (CL)	4 []
medium	Lisette (CL)	5 []
medium to dark	Emma (CL), Starlett (CL)	6 []
dark	Ines (CL), Laica (FL), Menuette (CL)	7 []
dark to very dark	Lion (FL)	8 []
very dark	Greenso (FL), Titan (CL)	9 []
5.4 <u>Only varieties with leaf blade curling:</u> leaf blade: depth of incision (15)		
shallow	Grüne Perle (CL), Lisette (CL)	1 []
medium		2 []
deep	Menuette (CL)	3 []

Characteristics	Example Varieties	Note
5.5 Root: thickening of main root (20)		
absent	Menuette (CL)	1 []
present	Halblange (R)	9 []
5.6 <u>Only root parsley varieties:</u> Root: length (21)		
very short	Halblange (R)	1 []
very short to short		2 []
short		3 []
short to medium		4 []
medium	A grosse racine gros hatif (R)	5 []
medium to long	Arat (R)	6 []
long	Artica (R)	7 []
long to very long		8 []
very long		9 []
5.7 <u>Only root parsley varieties:</u> Root: ratio length/width (23)		
small		1 []
small to medium		2 []
medium		3 []
medium to large	Artica (R)	4 []
large		5 []

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

6. Similar varieties and differences from these varieties

Please use the following table and box for comments to provide information on how your candidate variety differs from the variety (or varieties) which, to the best of your knowledge, is (or are) most similar. This information may help the examination authority to conduct its examination of distinctness in a more efficient way.

Denomination(s) of variety(ies) similar to your candidate variety	Characteristic(s) in which your candidate variety differs from the similar variety(ies)	Describe the expression of the characteristic(s) for the similar variety(ies)	Describe the expression of the characteristic(s) for your candidate variety
<i>Example</i>	<i>Leaf blade: width</i>	<i>medium</i>	<i>broad</i>

Comments:

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

8. Authorization for release

(a) Does the variety require prior authorization for release under legislation concerning the protection of the environment, human and animal health?

Yes [] No []

(b) Has such authorization been obtained?

Yes [] No []

If the answer to (b) is yes, please attach a copy of the authorization.

9. Information on plant material to be examined or submitted for examination

9.1 The expression of a characteristic or several characteristics of a variety may be affected by factors, such as pests and disease, chemical treatment (e.g. growth retardants or pesticides), effects of tissue culture, different rootstocks, scions taken from different growth phases of a tree, etc.

9.2 The plant material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If the plant material has undergone such treatment, full details of the treatment must be given. In this respect, please indicate below, to the best of your knowledge, if the plant material to be examined has been subjected to:

- | | | | |
|-----|---|---------|--------|
| (a) | Microorganisms (e.g. virus, bacteria, phytoplasma) | Yes [] | No [] |
| (b) | Chemical treatment (e.g. growth retardant, pesticide) | Yes [] | No [] |
| (c) | Tissue culture | Yes [] | No [] |
| (d) | Other factors | Yes [] | No [] |

Please provide details for where you have indicated "yes".

.....

10. I hereby declare that, to the best of my knowledge, the information provided in this form is correct:

Applicant's name

Signature

Date

[End of document]