



These Test Guidelines have been superseded by a later version. The latest adopted version of Test Guidelines can be found at http://www.upov.int/test_guidelines/en/list.jsp

Ces principes directeurs d'examen ont été remplacés par une version ultérieure. La version adoptée la plus récente des principes directeurs d'examen figure à l'adresse suivante : http://www.upov.int/test_guidelines/fr/list.jsp

Diese Prüfungsrichtlinien wurden durch eine neuere Fassung ersetzt. Die neueste angenommene Fassung von Prüfungsrichtlinien ist unter http://www.upov.int/test_guidelines/de/list.jsp zu finden.

Las presentes directrices de examen han sido reemplazadas por una versión posterior. La versión de las directrices de examen de más reciente aprobación está disponible en http://www.upov.int/test_guidelines/es/list.jsp.



TG/106/4

ORIGINAL: English

DATE: 2004-03-31

INTERNATIONAL UNION FOR THE PROTECTION OF NEW VARIETIES OF PLANTS
GENEVA

LEAF BEET, SWISS CHARD

(Beta vulgaris L. var. *cicla* L. (Ulrich))

GUIDELINES

FOR THE CONDUCT OF TESTS

FOR DISTINCTNESS, UNIFORMITY AND STABILITY

Alternative Names:*

| <i>Latin</i> | <i>English</i> | <i>French</i> | <i>German</i> | <i>Spanish</i> |
|---|---------------------------|---------------------------|---------------|----------------|
| <i>Beta vulgaris</i> L. var. <i>cicla</i> L. (Ulrich), <i>Beta vulgaris</i> L. var. <i>vulgaris</i> L. | Leaf Beet, Swiss Chard | Poirée, Bette à cardes | Mangold | Acelga |

ASSOCIATED DOCUMENTS

These guidelines should be read in conjunction with document TG/1/3, “General Introduction to the Examination of Distinctness, Uniformity and Stability and the Development of Harmonized Descriptions of New Varieties of Plants” (hereinafter referred to as the “General Introduction”) and its associated “TGP” documents.

* These names were correct at the time of the introduction of these Test Guidelines but may be revised or updated. [Readers are advised to consult the UPOV Code, which can be found on the UPOV Website (www.upov.int), for the latest information.]

| <u>TABLE OF CONTENTS</u> | <u>PAGE</u> |
|---|-------------|
| 1. SUBJECT OF THESE TEST GUIDELINES | 3 |
| 2. MATERIAL REQUIRED | 3 |
| 3. METHOD OF EXAMINATION..... | 3 |
| 3.1 Duration of Tests | 3 |
| 3.2 Testing Place | 3 |
| 3.3 Conditions for Conducting the Examination | 3 |
| 3.4 Test Design..... | 4 |
| 3.5 Number of Plants / Parts of Plants to be Examined..... | 4 |
| 3.6 Additional Tests | 4 |
| 4. ASSESSMENT OF DISTINCTNESS, UNIFORMITY AND STABILITY | 4 |
| 4.1 Distinctness..... | 4 |
| 4.2 Uniformity | 5 |
| 4.3 Stability..... | 5 |
| 5. GROUPING OF VARIETIES AND ORGANIZATION OF THE GROWING TRIAL | 5 |
| 6. INTRODUCTION TO THE TABLE OF CHARACTERISTICS..... | 6 |
| 6.1 Categories of Characteristics | 6 |
| 6.2 States of Expression and Corresponding Notes..... | 6 |
| 6.3 Types of Expression | 6 |
| 6.4 Example Varieties | 6 |
| 6.5 Legend | 7 |
| 7. TABLE OF CHARACTERISTICS/TABLEAU DES CARACTÈRES/MERKMALSTABELLE/TABLA DE CARACTERES | 8 |
| 8. EXPLANATIONS ON THE TABLE OF CHARACTERISTICS | 13 |
| 8.1 Explanations covering several characteristics | 13 |
| 8.2 Explanations for individual characteristics..... | 13 |
| 9. LITERATURE..... | 15 |
| 10. TECHNICAL QUESTIONNAIRE | 16 |

1. Subject of these Test Guidelines

These Test Guidelines apply to all varieties of *Beta vulgaris* L. var. *cicla* L. (Ulrich) (Syn. *Beta vulgaris* L. var. *vulgaris* L.).

2. Material Required

2.1 The competent authorities decide on the quantity and quality of the plant material required for testing the variety and when and where it is to be delivered. Applicants submitting material from a State other than that in which the testing takes place must ensure that all customs formalities and phytosanitary requirements are complied with.

2.2 The material is to be supplied in the form of seed.

2.3 The minimum quantity of plant material, to be supplied by the applicant, should be:

100 g or at least 6,600 seeds.

2.4 The seed should meet the minimum requirements for germination, species and analytical purity, health and moisture content, specified by the competent authority.

2.5 The plant material supplied should be visibly healthy, not lacking in vigor, nor affected by any important pest or disease.

2.6 The plant material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If it has been treated, full details of the treatment must be given.

3. Method of Examination

3.1 *Duration of Tests*

The minimum duration of tests should normally be two independent growing cycles.

3.2 *Testing Place*

The tests should normally be conducted at one place. If any characteristics of the variety, which are relevant for the examination of DUS, cannot be observed at that place, the variety may be tested at an additional place.

3.3 *Conditions for Conducting the Examination*

The tests should be carried out under conditions ensuring satisfactory growth for the expression of the relevant characteristics of the variety and for the conduct of the examination.

3.3.1 Type of observation

The recommended method of observing the characteristic is indicated by the following key in the second column of the Table of Characteristics:

- MG: single measurement of a group of plants or parts of plants;
- MS: measurement of a number of individual plants or parts of plants;
- VG: visual assessment by a single observation of a group of plants or parts of plants;
- VS: visual assessment by observation of individual plants or parts of plants.

3.4 *Test Design*

3.4.1 Each test should be designed to result in a total of at least 100 plants in the open or 60 plants in the greenhouse or plastic tunnel, which should be divided between two or more replicates.

3.4.2 The design of the tests should be such that plants or parts of plants may be removed for measurement or counting without prejudice to the observations which must be made up to the end of the growing cycle.

3.5 *Number of Plants / Parts of Plants to be Examined*

Unless otherwise indicated, all observations on single plants should be made on 20 plants or parts taken from each of 20 plants and any other observations made on all plants in the test.

3.6 *Additional Tests*

Additional tests, for examining relevant characteristics, may be established.

4. Assessment of Distinctness, Uniformity and Stability

4.1 *Distinctness*

4.1.1 General Recommendations

It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding distinctness. However, the following points are provided for elaboration or emphasis in these Test Guidelines.

4.1.2 Consistent Differences

The minimum duration of tests recommended in Section 3.1 reflects, in general, the need to ensure that any differences in a characteristic are sufficiently consistent.

4.1.3 Clear Differences

Determining whether a difference between two varieties is clear depends on many factors, and should consider, in particular, the type of expression of the characteristic being

examined, i.e. whether it is expressed in a qualitative, quantitative, or pseudo-qualitative manner. Therefore, it is important that users of these Test Guidelines are familiar with the recommendations contained in the General Introduction prior to making decisions regarding distinctness.

4.2 *Uniformity*

It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding uniformity. However, the following points are provided for elaboration or emphasis in these Test Guidelines:

4.2.1 Cross-pollinated varieties

The assessment of uniformity for cross-pollinated varieties should be according to the recommendations for cross-pollinated varieties in the General Introduction.

4.2.2 Hybrids

The assessment of uniformity for hybrid varieties depends on the type of hybrid and should be according to the recommendations for hybrid varieties in the General Introduction.

4.3 *Stability*

4.3.1 In practice, it is not usual to perform tests of stability that produce results as certain as those of the testing of distinctness and uniformity. However, experience has demonstrated that, for many types of variety, when a variety has been shown to be uniform, it can also be considered to be stable.

4.3.2 Where appropriate, or in cases of doubt, stability may be tested, either by growing a further generation, or by testing a new seed stock to ensure that it exhibits the same characteristics as those shown by the previous material supplied.

4.3.3 Where appropriate, or in cases of doubt, the stability of a hybrid variety may, in addition to an examination of the hybrid variety itself, also be assessed by examination of the uniformity and stability of its parent lines.

5. Grouping of Varieties and Organization of the Growing Trial

5.1 The selection of varieties of common knowledge to be grown in the trial with the candidate varieties and the way in which these varieties are divided into groups to facilitate the assessment of distinctness is aided by the use of grouping characteristics.

5.2 Grouping characteristics are those in which the documented states of expression, even where produced at different locations, can be used, either individually or in combination with other such characteristics: (a) to select varieties of common knowledge that can be excluded from the growing trial used for examination of distinctness; and (b) to organize the growing trial so that similar varieties are grouped together.

5.3 The following have been agreed as useful grouping characteristics:

- (a) Leaf blade: intensity of green color (characteristic 7);
- (b) Leaf blade: anthocyanin coloration (characteristic 11);
- (c) Petiole: width (characteristic 14);
- (d) Petiole: color (characteristic 16).

5.4 Guidance for the use of grouping characteristics, in the process of examining distinctness, is provided through the General Introduction.

6. Introduction to the Table of Characteristics

6.1 *Categories of Characteristics*

6.1.1 Standard Test Guidelines Characteristics

Standard Test Guidelines characteristics are those which are approved by UPOV for examination of DUS and from which members of the Union can select those suitable for their particular circumstances.

6.1.2 Asterisked Characteristics

Asterisked characteristics (denoted by *) are those included in the Test Guidelines which are important for the international harmonization of variety descriptions and should always be examined for DUS and included in the variety description by all members of the Union, except when the state of expression of a preceding characteristic or regional environmental conditions render this inappropriate.

6.2 *States of Expression and Corresponding Notes*

States of expression are given for each characteristic to define the characteristic and to harmonize descriptions. Each state of expression is allocated a corresponding numerical note for ease of recording of data and for the production and exchange of the description.

6.3 *Types of Expression*

An explanation of the types of expression of characteristics (qualitative, quantitative and pseudo-qualitative) is provided in the General Introduction.

6.4 *Example Varieties*

Where appropriate, example varieties are provided to clarify the states of expression of each characteristic.

6.5 *Legend*

(*) Asterisk characteristic – see Section 6.1.2

QL Qualitative characteristic – see Section 6.3

QN Quantitative characteristic – see Section 6.3

PQ Pseudo-qualitative characteristic – see Section 6.3

MG: Single measurement of a group of plants or parts of plants – see Section 3.3.1

MS: Measurement of a number of individual plants or parts of plants – see Section 3.3.1

VG: Visual assessment by a single observation of a group of plants or parts of plants – see Section 3.3.1

VS: Visual assessment by observation of individual plants or parts of plants – see Section 3.3.1

(a) See Explanations on the Table of Characteristics in Chapter 8, Section 8.1

(+) See Explanations on the Table of Characteristics in Chapter 8, Section 8.2

7. Table of Characteristics/Tableau des caractères/Merkmalstabelle/Tabla de caracteres

| | English | français | deutsch | español | Example Varieties/ Exemples/ Beispielssorten/ Variedades ejemplo | Note/ Nota |
|------------------|---|---|---|--|---|---------------|
| 1. (*) | VS Seedling: anthocyanin coloration | Plantule: pigmentation anthocyanique | Keimpflanze: Anthocyanfärbung | Plantúla: pigmentación antociánica | | |
| QL | absent | absente | fehlend | ausente | Blonde à cardé blanche | 1 |
| | present | présente | vorhanden | presente | Rhubarb Chard | 9 |
| 2. (*) | VS Seedling: intensity of anthocyanin coloration | Plantule: intensité de la pigmentation anthocyanique | Keimpflanze: Stärke der Anthocyanfärbung | Plantúla: intensidad de la pigmentación antociánica | | |
| QN | weak | faible | gering | débil | Verde de penca blanca ancha | 3 |
| | medium | moyenne | mittel | media | Amarilla de Lyon | 5 |
| | strong | forte | stark | fuerte | Rhubarb Chard | 7 |
| 3. (*)(+) | VG Leaf: length | Feuille: longueur | Blatt: Länge | Hoja: longitud | | |
| QN (a) | short | courte | kurz | corto | Groene Gewone, Verde de penca blanca ancha | 3 |
| | medium | moyenne | mittel | medio | Blonde à cardé blanche | 5 |
| | long | longue | lang | largo | Verte à cardé blanche, Paros | 7 |
| 4. (*) | VG Leaf: attitude | Feuille: port | Blatt: Haltung | Hoja: porte | | |
| QN (a) | erect | dressé | aufrecht | erecto | Paros | 1 |
| | semi erect | demi-dressé | halbaufrecht | semierecto | Blonde à cardé blanche | 3 |
| | prostrate | étalé | liegend | postrado | Groene Gewone | 5 |

| | English | français | deutsch | español | Example Varieties/ Exemples/ Beispielssorten/ Variedades ejemplo | Note/ Nota | |
|-----------|------------|---|---|---|---|------------------------------------|---|
| 5. | VG | Leaf blade: length | Limbe: longueur | Blattspreite: Länge | Limbo: longitud | | |
| (*) | | | | | | | |
| (+) | | | | | | | |
| QN | (a) | short | court | kurz | corto | Amarilla de Lyon, Groene Gewone | 3 |
| | | medium | moyen | mittel | medio | Verde de Niza | 5 |
| | | long | long | lang | largo | Blonde à cardé blanche, Paros | 7 |
| 6. | VG | Leaf blade: width | Limbe: largeur | Blattspreite: Breite | Limbo: anchura | | |
| (*) | | | | | | | |
| (+) | | | | | | | |
| QN | (a) | narrow | étroit | schmal | estrecho | Lucullus, Groene Gewone | 3 |
| | | medium | moyen | mittel | medio | Paros | 5 |
| | | broad | large | breit | ancho | Verte à cardé blanche | 7 |
| 7. | VG | Leaf blade: intensity of green color | Limbe: intensité de la couleur verte | Blattspreite: Stärke der Grünfärbung | Limbo: intensidad del color verde | | |
| (*) | | | | | | | |
| QN | (a) | very light | très claire | sehr hell | muy claro | Amarilla de Lyon | 1 |
| | | light | claire | hell | claro | Blonde à cardé blanche | 3 |
| | | medium | moyenne | mittel | medio | Verde de Niza | 5 |
| | | dark | foncée | dunkel | oscuro | Verde de penca blanca ancha | 7 |
| | | very dark | très foncée | sehr dunkel | muy oscuro | Verde de penca blanca larga | 9 |
| 8. | VG | Leaf blade: reflexing of margin | Limbe: enroulement du bord | Blattspreite: Randbiegung | Limbo: curvatura del ápice | | |
| QN | (a) | absent or very weak | nul ou très faible | fehlend oder sehr gering | ausente o muy débil | Groene Gewone | 1 |
| | | weak | faible | gering | débil | Blonde à cardé blanche | 3 |
| | | medium | moyen | mittel | media | | 5 |
| | | strong | fort | stark | fuerte | Lucullus | 7 |

| | English | français | deutsch | español | Example Varieties/ Exemples/ Beispielssorten/ Variedades ejemplo | Note/ Nota |
|------------|------------|--|--|--|---|---------------------------------|
| 9. | VG | Leaf blade: glossiness | Limbe: brillance | Blattspreite: Glanz | Limbo: brillo | |
| QN | (a) | weak | faible | gering | débil | Groene Gewone 3 |
| | | medium | moyenne | mittel | medio | 5 |
| | | strong | forte | stark | fuerte | Blonde à carde blanche, Paros 7 |
| 10. | VG | Leaf blade: blistering | Limbe: cloûre | Blattspreite: Blasigkeit | Limbo: abullonado | |
| QN | (a) | weak | faible | gering | débil | Groene Gewone 3 |
| | | medium | moyenne | mittel | medio | Blonde à carde blanche, Paros 5 |
| | | strong | forte | stark | fuerte | Lucullus 7 |
| 11. | VG | Leaf blade: anthocyanin coloration | Limbe: pigmentation anthocyanique | Blattspreite: Anthocyanfärbung | Limbo: pigmentación antociánica | |
| QL | (a) | absent | absente | fehlend | ausente | Blonde à carde blanche 1 |
| | | present | présente | vorhanden | presente | Rhubarb Chard 9 |
| 12. | VG | Leaf blade: intensity of anthocyanin coloration | Limbe: intensité de la pigmentation anthocyanique | Blattspreite: Stärke der Anthocyanfär- bung | Limbo: intensidad de la pigmentación antociánica | |
| QN | (a) | weak | faible | gering | débil | 3 |
| | | medium | moyenne | mittel | media | Rhubarb Chard 5 |
| | | strong | forte | stark | fuerte | Charlotte 7 |

| | English | français | deutsch | español | Example Varieties/ Exemples/ Beispielssorten/ Variedades ejemplo | Note/ Nota |
|------------|------------|--|--|---|---|---|
| 13. | VS | Petiole: length | Pétiolo: longueur | Blattstiel: Länge | Pecíolo: longitud | |
| | (+) | | | | | |
| QN | (a) | very short | très court | sehr kurz | muy corto | 1 |
| | | short | court | kurz | corto | Lucullus 3 |
| | | medium | moyen | mittel | medio | Paros 5 |
| | | long | long | lang | largo | Blonde à carde blanche, Verde de penca blanca larga 7 |
| | | very long | très long | sehr lang | muy largo | Groene Gewone 9 |
| 14. | VS | Petiole: width | Pétiolo: largeur | Blattstiel: Breite | Pecíolo: anchura | |
| | (*) (+) | | | | | |
| QN | (a) | very narrow | très étroit | sehr schmal | muy estrecho | Groene Gewone 1 |
| | | narrow | étroit | schmal | estrecho | Rhubarb Chard, Verde de Niza 3 |
| | | medium | moyen | mittel | medio | Lucullus, Verde de penca blanca larga 5 |
| | | broad | large | breit | ancho | Amarilla de Lyon 7 |
| | | very broad | très large | sehr breit | muy ancho | Paros, Verde de penca blanca ancha 9 |
| 15. | VS | Petiole: curvature of inner side in cross section | Pétiolo: courbure de la face interne de la section transversale | Blattstiel: Krümmung der Innenseite im Querschnitt | Pecíolo: curvatura de la cara interna en sección transversal | |
| | (+) | | | | | |
| QN | (a) | absent or very weak | absente ou très faible | fehlend oder sehr gering | ausente o muy débil | 1 |
| | | weak | faible | gering | débil | Groene Gewone 3 |
| | | medium | moyenne | mittel | media | Lucullus 5 |
| | | strong | forte | stark | fuerte | Blonde à carde blanche 7 |

| | English | français | deutsch | español | Example Varieties/ Exemples/ Beispielssorten/ Variedades ejemplo | Note/ Nota |
|----------------------|-------------------------------------|-------------------------------------|------------------------------------|--|---|---------------|
| 16. VG (*) | Petiole : color | Pétiole: couleur | Blattstiel: Farbe | Pecíolo: color | | |
| PQ | (a) white | blanc | weiß | blanco | Blonde à cardé blanche | 1 |
| | yellow | jaune | gelb | amarillo | Bright Yellow | 2 |
| | green | vert | grün | verde | Groene Gewone | 3 |
| | pink | rose | rosa | rosa | Pink Passion | 4 |
| | purple | pourpre | purpurn | púrpura | Rhubarb Chard, Ruby Red | 5 |
| 17. VG | Time of beginning of bolting | Époque de début de montaison | Zeitpunkt des Schossbeginns | Época del comienzo del espigado | | |
| QN | early | précoce | früh | temprana | Paros, Verde de Niza | 3 |
| | medium | moyenne | mittel | media | Verde de penca blanca ancha | 5 |
| | late | tardive | spät | tardía | Amarilla de Lyon | 7 |

8. Explanations on the Table of Characteristics

8.1 *Explanations covering several characteristics*

Characteristics containing the following key in the second column of the Table of Characteristics should be examined as indicated below:

- (a) All observations on the foliage, the leaf blade and the petiole should be made when the foliage has reached its maximum height.

8.2 *Explanations for individual characteristics*


Ad. 3: Leaf: length

Ad. 5: Leaf blade: length

Ad. 6: Leaf blade: width

Ad. 13: Petiole: length

Ad. 14: Petiole: width



Ad. 15: Petiole: curvature of inner side in cross section



1
absent or very weak



5
medium



7
strong

9. Literature

No specific literature

10. Technical Questionnaire

| | | |
|---|---|---|
| TECHNICAL QUESTIONNAIRE | Page {x} of {y} | Reference Number: |
| | | Application date: (not to be filled in by the applicant) |
| <p>TECHNICAL QUESTIONNAIRE to be completed in connection with an application for plant breeders' rights</p> <p>In the case of hybrid varieties which are the subject of an application for plant breeders' rights, and where the parent lines are to be submitted as a part of the examination of the hybrid variety, this Technical Questionnaire should be completed for each of the parent lines, in addition to being completed for the hybrid variety.</p> | | |
| 1. Subject of the Technical Questionnaire | | |
| 1.1 Latin Name | <input type="text" value="Beta vulgaris L. var. cicla L. (Ulrich) (Syn. Beta vulgaris L. var. vulgaris L.)"/> | |
| 1.2 Common Name | <input type="text" value="Leaf Beet, Swiss Chard"/> | |
| 2. Applicant | | |
| Name | <input type="text"/> | |
| Address | <input type="text"/> | |
| Telephone No. | <input type="text"/> | |
| Fax No. | <input type="text"/> | |
| E-mail address | <input type="text"/> | |
| Breeder (if different from applicant) | <input type="text"/> | |
| 3. Proposed denomination and breeder's reference | | |
| Proposed denomination (if available) | <input type="text"/> | |
| Breeder's reference | <input type="text"/> | |

TECHNICAL QUESTIONNAIRE

Page {x} of {y}

Reference Number:

4. Information on the breeding scheme and propagation of the variety

4.1 Breeding scheme

Variety resulting from:

4.1.1 Crossing

- (a) controlled cross []
(please state parent varieties)
- (b) partially known cross []
(please state known parent variety(ies))
- (c) unknown cross []

4.1.2 Mutation []
(please state parent variety)

4.1.3 Discovery and development []
(please state where and when discovered
and how developed)

4.1.4 Other []
(please provide details)

4.2 Method of propagating the variety

4.2.1 Seed-propagated varieties

- (a) Self-pollination []
- (b) Cross-pollination []
 - (i) population []
 - (ii) synthetic variety []
- (c) Hybrid []
- (d) Other []
(please provide details)

4.2.2 Other []
(please provide details)

| | | |
|-------------------------|-----------------|-------------------|
| TECHNICAL QUESTIONNAIRE | Page {x} of {y} | Reference Number: |
|-------------------------|-----------------|-------------------|

5. Characteristics of the variety to be indicated (the number in brackets refers to the corresponding characteristic in Test Guidelines; please mark the note which best corresponds).

| Characteristics | Example Varieties | Note |
|---|---------------------------------------|------|
| 5.1 Leaf blade : intensity of green color (7) | | |
| very light | Amarilla de Lyon | 1[] |
| light | Blonde à cardé blanche | 3[] |
| medium | Verde de Niza | 5[] |
| dark | Verde de penca blanca ancha | 7[] |
| very dark | Verde de penca blanca larga | 9[] |
| 5.2 Leaf blade: anthocyanin coloration (11) | | |
| absent | Blonde à cardé blanche | 1[] |
| present | Rhubarb Chard | 9[] |
| 5.3 Petiole: width (14) | | |
| very narrow | Groene Gewone | 1[] |
| narrow | Rhubarb Chard, Verde de Niza | 3[] |
| medium | Lucullus, Verde de penca blanca larga | 5[] |
| broad | Amarilla de Lyon | 7[] |
| very broad | Paros, Verde de penca blanca ancha | 9[] |
| 5.4 Petiole : color (16) | | |
| white | Blonde à cardé blanche | 1[] |
| yellow | Bright Yellow | 2[] |
| green | Groene Gewone | 3[] |
| pink | Pink Passion | 4[] |
| purple | Rhubarb Chard, Ruby Red | 5[] |

| | | |
|-------------------------|-----------------|-------------------|
| TECHNICAL QUESTIONNAIRE | Page {x} of {y} | Reference Number: |
|-------------------------|-----------------|-------------------|

6. Similar varieties and differences from these varieties

Please use the table, and space provided for comments, below to provide information on how your candidate variety differs from the variety (or varieties) which, to the best of your knowledge, is (or are) most similar. This information may help the examination authority to conduct its examination of distinctness in a more efficient way.

| Denomination(s) of variety(ies) similar to your candidate variety | Characteristic(s) in which your candidate variety differs from the similar variety(ies) | Describe the expression of the characteristic(s) for the similar variety(ies) | Describe the expression of the characteristic(s) for your candidate variety |
|---|---|--|--|
| <i>Example</i> | <i>Petiole: color</i> | <i>pink</i> | <i>purple</i> |
| | | | |

Comments:

7. Additional information which may help in the examination of the variety

7.1 In addition to the information provided in sections 5 and 6, are there any additional characteristics which may help to distinguish the variety?

Yes [] No []

(If yes, please provide details)

7.2 Special conditions for the examination of the variety

7.2.1 Are there any special conditions for growing the variety or conducting the examination?

Yes [] No []

7.2.2 If yes, please give details:

7.3 Other information

| | | |
|-------------------------|-----------------|-------------------|
| TECHNICAL QUESTIONNAIRE | Page {x} of {y} | Reference Number: |
|-------------------------|-----------------|-------------------|

8. Authorization for release

(a) Does the variety require prior authorization for release under legislation concerning the protection of the environment, human and animal health?

Yes [] No []

(b) Has such authorization been obtained?

Yes [] No []

If the answer to (b) is yes, please attach a copy of the authorization.

9. Information on plant material to be examined.

9.1 The expression of a characteristic or several characteristics of a variety may be affected by factors, such as pests and disease, chemical treatment (e.g. growth retardants or pesticides), effects of tissue culture, different rootstocks, scions taken from different growth phases of a tree, etc.

9.2 The plant material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If the plant material has undergone such treatment, full details of the treatment must be given. In this respect, please indicate below, to the best of your knowledge, if the plant material to be examined has been subjected to:

(a) Microorganisms (e.g. virus, bacteria, phytoplasma) Yes [] No []

(b) Chemical treatment (e.g. growth retardant or pesticide) Yes [] No []

(c) Tissue culture Yes [] No []

(d) Other factors Yes [] No []

Please provide details of where you have indicated "yes":

.....

10. I hereby declare that, to the best of my knowledge, the information provided in this form is correct:

Applicant's name

Signature

Date

[End of document]