



TG/202/1 Rev. 2

ORIGINAL: English

DATE: 2003-04-09 + 2015-03-25

+ 2019-10-29

INTERNATIONAL UNION FOR THE PROTECTION OF NEW VARIETIES OF PLANTS

GENEVA

CITRUS L. – Group 2

ORANGES

GUIDELINES

FOR THE CONDUCT OF TESTS

FOR DISTINCTNESS, UNIFORMITY AND STABILITY

Alternative Names:

[See the list of alternative names and corresponding subgroups on page 2]

ASSOCIATED DOCUMENTS

These guidelines should be read in conjunction with document TG/1/3, “General Introduction to the Examination of Distinctness, Uniformity and Stability and the Development of Harmonized Descriptions of New Varieties of Plants” (hereinafter referred to as the “General Introduction”) and its associated “TGP” documents.

Other associated documents:

CITRUS L. – GROUP 1: TG/201 – Mandarins

CITRUS L. – GROUP 3: TG/203 – Lemons and Limes

CITRUS L. – GROUP 4: TG/204 – Grapefruit and Pummelo

CITRUS L. – GROUP 5: TG/83 – Trifoliate Orange

GROUP 2 – ALTERNATIVE NAMES AND CORRESPONDING SUBGROUPS*

<i>Latin</i>	<i>Subgroup</i>	<i>English</i>	<i>French</i>	<i>German</i>	<i>Spanish</i>
<i>Citrus aurantium</i> L.	SOR	Sour Orange	Bigaradier, Oranger amer	Sauerorange, Bitterorange, Pomeranze	Naranjo agrio, Naranjo amargo
<i>Citrus aurea</i> hort. ex Tanaka	SWO				
<i>Citrus canaliculata</i> hort. ex Yu. Tan	SOR				
<i>Citrus funadoko</i> hort. ex Yu. Tanaka	SWO				
<i>Citrus iriomotensis</i> hort. ex Tanaka	HOR				
<i>Citrus iyo</i> hort. ex Tanaka	SWO				
<i>Citrus luteoturgida</i> hort. ex Tanaka	SWO				
<i>Citrus maderaspatana</i> hort. ex Tan.	SOR				
<i>Citrus myrtifolia</i> Raf.	SOR				
<i>Citrus neoaurantium</i> Tanaka	SOR				
<i>Citrus oblonga</i> hort. ex Yu. Tanaka	SWO				
<i>Citrus papillaris</i> Blanco	HOR				
<i>Citrus pseudopapillaris</i> Tanaka	HOR				
<i>Citrus rokugatsu</i> hort. ex Yu. Tanaka	SOR				
<i>Citrus shunkokan</i> hort. ex Tanaka	SWO				
<i>Citrus sinensis</i> (L.) Osbeck	SWO	Sweet Orange	Oranger doux	Apfelsine	Naranjo dulce
<i>Citrus sinograndis</i> hort. ex Yu. Tanaka	SWO				
<i>Citrus taiwanica</i> Tanaka & Y. Shimada	HOR				
<i>Citrus tamurana</i> hort. ex Tanaka	SWO				
<i>Citrus tankan</i> Hayata	SWO				
<i>Citrus ujukitsu</i> Tanaka	SWO				
<i>Citrus yanbaruensis</i> hort. ex Tanaka	SOR				

* These names were correct at the time of the introduction of these Test Guidelines but may be revised or updated. [Readers are advised to consult the UPOV Code, which can be found on the UPOV Web Site (www.upov.int), for the latest information.]

<u>TABLE OF CONTENTS</u>	<u>PAGE</u>
1. SUBJECT OF THESE TEST GUIDELINES	4
2. MATERIAL REQUIRED	4
3. METHOD OF EXAMINATION.....	5
3.1 Duration of Tests.....	5
3.2 Testing Place.....	5
3.3 Conditions for Conducting the Examination.....	5
3.4 Test Design	5
3.5 Number of Plants / Parts of Plants to be Examined.....	5
3.6 Additional Tests	5
4. ASSESSMENT OF DISTINCTNESS, UNIFORMITY AND STABILITY.....	6
4.1 Distinctness	6
4.1.1 General Recommendations	6
4.1.2 Consistent Differences	6
4.1.3 Clear Differences.....	6
4.2 Uniformity.....	6
4.3 Stability	6
5. GROUPING OF VARIETIES AND ORGANIZATION OF THE GROWING TRIAL	6
6. INTRODUCTION TO THE TABLE OF CHARACTERISTICS	7
6.1 Categories of Characteristics.....	7
6.1.1 Standard Test Guidelines Characteristics	7
6.1.2 Asterisked Characteristics	7
6.2 States of Expression and Corresponding Notes.....	7
6.3 Types of Expression.....	7
6.4 Example Varieties	8
6.5 Legend.....	8
6.6 Abbreviations	8
7. TABLE OF CHARACTERISTICS	9
8. EXPLANATIONS ON THE TABLE OF CHARACTERISTICS	30
8.1 Explanations covering several characteristics	30
8.2 Explanations for individual characteristics	31
9. LITERATURE	37
10. TECHNICAL QUESTIONNAIRE.....	39
ANNEX: OVERALL TABLE OF CHARACTERISTICS	

1. Subject of these Test Guidelines

1.1 The following Test Guidelines have been developed from the standard Citrus Test Guidelines template. In particular, the Table of Characteristics has been selected from the overall set of citrus characteristics presented in the Annex.

1.2 These Test Guidelines apply to all varieties of the following group of the genus *Citrus* L. (*Rutaceae*), and their hybrids:

Group 2. ORANGES AND THEIR HYBRIDS

See page 2 for the list of species and their subgroups.

1.3 In the case of hybrids between species within the genus *Citrus* L., the Test Guidelines to be used should be those for which the overall appearance of fruit is most suited. However, if the variety cannot be clearly distinguished from all varieties covered by other Test Guidelines, those other Test Guidelines should also be used to examine the variety.

1.4 In the case of hybrids between species within the genus *Citrus* L., even where the variety is clearly distinguishable from all other varieties covered by other Test Guidelines, it may still be necessary to use additional citrus characteristics to examine the variety. In these circumstances the characteristics from the Test Guidelines covering the parent species, or characteristics from the overall set of citrus characteristics, presented in the Annex, may be particularly useful.

2. Material Required

2.1 The competent authorities decide on the quantity and quality of the plant material required for testing the variety and when and where it is to be delivered. Applicants submitting material from a State other than that in which the testing takes place must ensure that all customs formalities and phytosanitary requirements are complied with.

2.2 The material is to be supplied in the form of bud sticks of 6 to 10 mm in diameter (one year old), each cut just behind a typical fruit or, if required by the competent authorities, one-year-old grafted trees. In the case of rootstock varieties, rooted cuttings or polyembryonic seeds may be required in addition.

2.3 The minimum quantity of plant material, to be supplied by the applicant, should be:

10 bud sticks sufficient to establish 10 plants or,
if required by the competent authorities,
10 one-year-old grafted trees.

2.4 The plant material supplied should be visibly healthy, not lacking in vigor, nor affected by any important pest or disease. It should preferably not be obtained from *in vitro* propagation. If it has been produced by *in vitro* propagation this fact must be stated by the applicant.

2.5 The plant material must not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If it has been treated, full details of the treatment must be given.

3. Method of Examination

3.1 *Duration of Tests*

The minimum duration of tests should normally be two independent growing cycles. For the purposes of these Test Guidelines, a growing cycle refers to the fruiting cycle.

3.2 *Testing Place*

The tests should normally be conducted at one place. If any characteristics of the variety, which are relevant for the examination of DUS, cannot be seen at that place, the variety may be tested at an additional place.

3.3 *Conditions for Conducting the Examination*

3.3.1 The tests should be carried out under conditions ensuring satisfactory growth for the expression of the relevant characteristics of the variety and for the conduct of the examination. In particular, a satisfactory crop of fruit must be produced in at least two fruiting cycles. Where necessary for the examination of fruit varieties, a standard specified rootstock should be used for each group.

3.3.2 All observations should be made on plants of the same age not less than three years after planting. The age of the plants should be specified.

3.4 *Test Design*

3.4.1 The design of the tests should be such that plants or parts of plants may be removed for measurement or counting without prejudice to the observations which must be made up to the end of the growing cycle.

3.4.2 Each test should be designed to result in a total of, at least, 5 plants.

3.5 *Number of Plants / Parts of Plants to be Examined*

Unless otherwise indicated, all observations determined by measuring or counting should be made on 5 plants or 2 parts taken from each of 5 plants.

3.6 *Additional Tests*

Additional tests, for examining relevant characteristics, may be established.

4. Assessment of Distinctness, Uniformity and Stability

4.1 *Distinctness*

4.1.1 General Recommendations

It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding distinctness. However, the following points are provided for elaboration or emphasis in these Test Guidelines.

4.1.2 Consistent Differences

The minimum duration of tests recommended in section 3.1 reflects, in general, the need to ensure that any differences in a characteristic are sufficiently consistent.

4.1.3 Clear Differences

Determining whether a difference between two varieties is clear depends on many factors, and should consider, in particular, the type of expression of the characteristic being examined, i.e. whether it is expressed in a qualitative, quantitative, or pseudo-qualitative manner. Therefore, it is important that users of these Test Guidelines are familiar with the recommendations contained in the General Introduction prior to making decisions regarding distinctness.

4.2 *Uniformity*

It is of particular importance for users of these Test Guidelines to consult the General Introduction prior to making decisions regarding uniformity. However, the following points are provided for elaboration or emphasis in these Test Guidelines:

For the assessment of uniformity a population standard of 1% and an acceptance probability of at least 95% should be applied. In the case of a sample size of 5 plants no off-types are allowed.

4.3 *Stability*

4.3.1 In practice, it is not usual to perform tests of stability that produce results as certain as those of the testing of distinctness and uniformity. However, experience has demonstrated that, for many types of variety, when a variety has been shown to be uniform, it can also be considered to be stable.

4.3.2 Where appropriate, or in cases of doubt, stability may be tested, either by growing a further generation, or by testing a new plant stock to ensure that it exhibits the same characteristics as those shown by the previous material supplied.

5. Grouping of Varieties and Organization of the Growing Trial

5.1 The selection of varieties of common knowledge to be grown in the trial with the candidate varieties and the way in which these varieties are divided into groups to facilitate the assessment of distinctness is aided by the use of grouping characteristics.

5.2 Grouping characteristics are those in which the documented states of expression, even where produced at different locations, can be used, either individually or in combination with other such characteristics: (a) to select varieties of common knowledge that can be excluded from the growing trial used for examination of distinctness; and (b) to organize the growing trial so that similar varieties are grouped together.

5.3 The following have been agreed as useful grouping characteristics:

- (a) Fruit: length (characteristic 29)
- (b) Fruit: diameter (characteristic 30)
- (c) Fruit surface: predominant color(s) (characteristic 51)
- (d) Fruit: presence of navel (viewed internally) (characteristic 78)
- (e) Time of maturity of fruit for consumption (characteristic 93).

5.4 Guidance for the use of grouping characteristics, in the process of examining distinctness, is provided through the General Introduction.

6. Introduction to the Table of Characteristics

6.1 *Categories of Characteristics*

6.1.1 Standard Test Guidelines Characteristics

Standard Test Guidelines characteristics are those which are approved by UPOV for examination of DUS and from which members of the Union can select those suitable for their particular circumstances.

6.1.2 Asterisked Characteristics

Asterisked characteristics (denoted by *) are those included in the Test Guidelines which are important for the international harmonization of variety descriptions and should always be examined for DUS and included in the variety description by all members of the Union, except when the state of expression of a preceding characteristic or regional environmental conditions render this inappropriate.

6.2 *States of Expression and Corresponding Notes*

States of expression are given for each characteristic to define the characteristic and to harmonize descriptions. Each state of expression is allocated a corresponding numerical note for ease of recording of data and for the production and exchange of the description.

6.3 *Types of Expression*

An explanation of the types of expression of characteristics (qualitative, quantitative and pseudo-qualitative) is provided in the general Introduction.

6.4 *Example Varieties*

Where appropriate, example varieties are provided to clarify the states of expression of each characteristic. Each example variety is followed by the abbreviation of its subgroup in brackets.

6.5 *Legend*

- (*) Asterisked characteristic – see section 6.1.2
- QL Qualitative characteristic – see section 6.3
- QN Quantitative characteristic – see section 6.3
- PQ Pseudo-Qualitative characteristic – see section 6.3
- [###] Corresponding number of characteristic in the citrus overall table of characteristics (starting from 200)
- (a)-(f) See Explanations on the Table of Characteristics in Chapter 8, Section 8.1
- (+) See Explanations on the Table of Characteristics in Chapter 8, Section 8.2

6.6 *Abbreviations*

See page 2 for the list of species and their subgroups.

7. Table of Characteristics/Tableau des caractères/Merkmalstabelle/Tabla de caracteresticas

					Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
1.	Ploidy	Ploidie	Ploidie	Ploidía		
QL	diploid	diploïde	diploid	diploide		2
	triploid	triploïde	triploid	triploide		3
[201]	tetraploid	tétraploïde	tetraploid	tetraploide		4
2. (*) (+)	Tree: growth habit	Arbre: port	Baum: Wuchstyp	Árbol: porte		
PQ	upright	droit	aufrecht	erguido	Salustiana (SWO)	1
	spreading	étalé	breitbuschig	abierto	Valencia Late (SWO)	2
[202]	drooping	retombant	hängend	colgante	Washington Navel (SWO)	3
3.	Tree: density of spines	Arbre: densité des épines	Baum: Dichte der Stacheln	Árbol: densidad de las espinas		
QN	absent or sparse	absentes ou éparses	fehlend oder locker	ausentes o laxa	Washington Navel (SWO)	1
	intermediate	intermédiaires	mittel	media	Valencia Late (SWO)	2
[203]	dense	denses	dicht	densa	Navelate (SWO)	3
4.	Tree: length of spines	Arbre: longueur des épines	Baum: Länge der Stacheln	Árbol: longitud de las espinas		
QN	short	courtes	kurz	cortas		3
	medium	moyennes	mittel	medias		5
[204]	long	longues	lang	largas		7
5.	Leaf blade: length (apical leaflet in case of compound leaf)	Limbe: longueur des folioles apicales en cas de feuille composée	Blattspreite: Länge (apikales Teilblatt bei zusammen-gesetztem Blatt)	Limbo: longitud (foliolos atípicos en caso de hoja compuesta)		
QN (a)	short	court	kurz	corto	Valencia Late (SWO)	3
	medium	moyen	mittel	medio	Salustiana (SWO)	5
[210]	long	long	lang	largo	Navelate (SWO)	7

					Example Varieties Exemples Beispielssorten Variedades ejemplos	Note/ Nota
	English	français	deutsch	español		
6.	Leaf blade: width (as for 5)	Limbe: largeur (comme pour 5)	Blattspreite: Breite (wie für 5)	Limbo: anchura (como para 5)		
QN	(a) narrow	étroit	schmal	estrecho	Lanelate (SWO)	3
	medium	moyen	mittel	medio	Salustiana (SWO)	5
[211]	broad	large	breit	ancho	Washington Navel (SWO)	7
7.	Leaf blade: ratio length/width (as for 5)	Limbe: rapport longueur/largeur (comme pour 5)	Blattspreite: Verhältnis Länge/Breite (wie für 5)	Limbo: relación longitud/anchura (como para 5)		
QN	(a) small	faible	klein	pequeño	Navelate (SWO)	3
	medium	moyen	mittel	medio	Salustiana (SWO)	5
[212]	large	élevé	groß	grande	Lanelate (SWO)	7
8.	Leaf blade: shape in cross section (as for 5)	Limbe: forme en section transversale (comme pour 5)	Blattspreite: Form im Querschnitt (wie für 5)	Limbo: forma en sección transversal (como para 5)		
QN	(a) straight or weakly concave	droit ou légèrement concave	gerade oder leicht konkav	recto o ligeramente cóncavo	Salustiana (SWO)	1
	intermediate	intermédiaire	mittel	intermedio	Washington Navel (SWO)	2
[217]	strongly concave	fortement concave	stark konkav	fuertemente cóncavo	Sweet Navel (SWO)	3
9.	Leaf blade: twisting	Limbe: torsion	Blattspreite: Drehung	Limbo: torsión		
QN	(a) absent or weak	absente ou faible	fehlend oder gering	ausente o débil	Washington Navel (SWO)	1
	intermediate	intermédiaire	mittel	media		2
[218]	strong	forte	stark	fuerte		3
10.	Leaf blade: blistering	Limbe: cloquère	Blattspreite: Blasigkeit	Limbo: abullonado o ampollado		
QN	(a) absent or weak	absente ou faible	fehlend oder gering	ausente o débil	Washington Navel (SWO)	1
	intermediate	intermédiaire	mittel	medio	Summer Navel (SWO)	2
[219]	strong	forte	stark	fuerte	Navel Mas Baró (SWO)	3

					Example Varieties Exemples Beispielssorten Variedades ejemplos	Note/ Nota
	English	français	deutsch	español		
11.	Leaf blade: green color	Limbe: couleur verte	Blattspreite: Grünfärbung	Limbo: color verde		
QN (a)	light	claire	hell	claro	Valencia Late (SWO)	3
	medium	moyenne	mittel	medio	Washington Navel (SWO)	5
[220]	dark	foncée	dunkel	oscuro	Navelina (SWO)	7
12.	Leaf blade: undulation of margin	Limbe: ondulation du bord	Blattspreite: Randwellung	Limbo: ondulación del borde		
QN (a)	absent or weak	absente ou faible	fehlend oder gering	ausente o débil	Washington Navel (SWO)	1
	intermediate	intermédiaire	mittel	media		2
[222]	strong	forte	stark	fuerte		3
13.	Leaf blade: incisions of margin	Limbe: incisions du bord	Blattspreite: Randeinschnitte	Limbo: incisiones del borde		
PQ (a)	absent	absentes	fehlend	ausentes		1
	crenate	crénelées	gekerbt	crenadas		2
[223]	dentate	dentelées	gezähnt	dentadas		3
14.	Leaf blade: shape of apex	Limbe: forme de l'extrémité	Blattspreite: Form der Spitze	Limbo: forma del ápice		
(+)						
PQ (a)	acuminate	acuminée	mit aufgesetzter Spitze	acuminado		1
	acute	pointue	spitz	agudo	Salustiana (SWO)	2
	obtuse	obtuse	stumpf	obtuso		3
[224]	rounded	arrondie	abgerundet	redondeado	Navelate (SWO)	4
15.	Leaf blade: emargination at tip	Limbe: échancrure à l'extrémité	Blattspreite: Einkerbung an der Spitze	Limbo: emarginado en la parte superior		
(+)						
QL (a)	absent	absente	fehlend	ausente	Washington Navel (SWO)	1
[225]	present	présente	vorhanden	presente		9

					Example Varieties Exemples Beispielssorten Variedades ejemplos	Note/ Nota
	English	français	deutsch	español		
16.	Petiole: length	Pétiole: longueur	Blattstiel: Länge	Pecíolo: longitud		
QN	(a) short	court	kurz	corto	Lanelate (SWO)	3
	medium	moyen	mittel	medio	Valencia	5
[226]	long	long	lang	largo	Navelina (SWO)	7
17.	Petiole: presence of wings	Pétiole: présence d'ailes	Blattstiel: Vorhandensein von Flügeln	Pecíolo: presencia de alas		
QL	(a) absent	absentes	fehlend	ausentes	Salustiana (SWO)	1
[227]	present	présentes	vorhanden	presentes	Newhall (SWO)	9
18.	Varieties with petiole wings present only: Petiole: width of wings	Seulement les variétés présentant des ailes au pétiole: Pétiole: largeur des ailes	Nur Sorten mit vorhandenen Flügeln am Blattstiel: Blattstiel: Breite der Flügel	Sólo variedades con alas presentes en el pecíolo: Pecíolo: anchura de las alas		
QN	(a) narrow	étroites	schmal	estrechas	Newhall (SWO)	3
	medium	moyennes	mittel	medianas		5
[228]	broad	larges	breit	anchas		7
19.	Flower: diameter of calyx	Fleur: diamètre du calice	Blüte: Durchmesser des Kelches	Flor: diámetro del cáliz		
QN	(b) small	petit	klein	pequeño		3
	medium	moyen	mittel	medio		5
[231]	large	grand	groß	grande		7
20.	Flower: length of petal	Fleur: longueur du pétales	Blüte: Länge des Blütenblattes	Flor: longitud del pétalo		
QN	(b) short	court	kurz	corto	Newhall (SWO)	3
	medium	moyen	mittel	medio	Lanelate (SWO)	5
[232]	long	long	lang	largo	Salustiana (SWO)	7
21.	Flower: width of petal	Fleur: largeur du pétales	Blüte: Breite des Blütenblattes	Flor: anchura del pétalo		
QN	(b) narrow	étroit	schmal	estrecho	Newhall (SWO)	3
	medium	moyen	mittel	medio	Lanelate (SWO)	5
[233]	broad	large	breit	ancho	Salustiana (SWO)	7

					Example Varieties Exemples Beispielssorten Variedades ejemplos	Note/ Nota
	English	français	deutsch	español		
22.	Flower: ratio length/ width of petal	Fleur: rapport longueur/largeur du pétales	Blüte: Verhältnis Länge/Breite des Blütenblattes	Flor: relación longitud/anchura del pétalo		
QN	(b) small	faible	klein	pequeña	Summer Navel (SWO)	3
	medium	moyen	mittel	media	Washington Navel (SWO)	5
[234]	large	élevé	groß	grande	Sanguinelli (SWO)	7
23.	Flower: length of stamens	Fleur: longueur des étamines	Blüte: Länge der Staubfäden	Flor: longitud de los estambres		
QN	(b) short	courtes	kurz	cortos	Newhall (SWO)	3
	medium	moyennes	mittel	medios	Washington Navel (SWO)	5
[235]	long	longues	lang	largos	Valencia Late (SWO)	7
24.	Flower: basal union of stamens	Fleur: union basale des étamines	Blüte: Vereinigung der Staubfäden an der Basis	Flor: unión basal de los estambres		
QL	(b) absent	absente	fehlend	ausente		1
[236]	present	présente	vorhanden	presente	Valencia Late (SWO)	9
25.	Anther: color	Anthère: couleur	Anthere: Farbe	Antera: color		
PQ	(b) white	blanc	weiß	blanco		1
	light yellow	jaune clair	hellgelb	amarillo claro	Washington Navel (SWO)	2
[238]	medium yellow	jaune moyen	mittelgelb	amarillo medio	Valencia Late (SWO)	3
26.	Anther: pollen viability	Anthère: viabilité du pollen	Anthere: Keimfähigkeit des Pollen	Antera: viabilidad del polen		
(+)						
QN	(b) absent or low	absente ou faible	fehlend oder gering	ausente o baja	Valencia Late (SWO), Washington Navel (SWO)	1
	medium	moyenne	mittel	media	Olinda (SWO)	2
[239]	high	forte	stark	alta	Comuna (SWO)	3
27.	Style: length	Style: longueur	Griffel: Länge	Estilo: longitud		
QN	(b) short	court	kurz	corto		3
	medium	moyen	mittel	medio		5
[240]	long	long	lang	largo		7

					Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
	English	français	deutsch	español		
28.	Style: shape	Style: forme	Griffel: Form	Estilo: forma		
PQ	(b) straight	droit	gerade	recto	Washington Navel (SWO)	1
	arched	arqué	nach unten gebogen	arqueado		2
[241]	kinked	coudé	geknickt	sinuoso		3
29. (*)	Fruit: length	Fruit: longueur	Frucht: Länge	Fruto: longitud		
QN	(c) short	court	kurz	corto	Comuna (SWO)	3
	medium	moyen	mittel	medio	Valencia Late (SWO)	5
[244]	long	long	lang	largo	Newhall (SWO)	7
30. (*)	Fruit: diameter	Fruit: diamètre	Frucht: Durchmesser	Fruto: diámetro		
QN	(c) small	petit	klein	pequeño	Sanguinelli (SWO)	3
	medium	moyen	mittel	medio	Valencia Late (SWO)	5
[245]	large	grand	groß	grande	Washington Navel (SWO)	7
31. (*)	Fruit: ratio length/diameter	Fruit: rapport longueur/diamètre	Frucht: Verhältnis Länge/Durchmesser	Fruto: relación longitud/diámetro		
QN	(c) small	faible	klein	pequeña	Salustiana (SWO)	3
	medium	moyen	mittel	media	Valencia Late (SWO)	5
[246]	large	élevé	groß	grande	Newhall (SWO)	7
32. (*)	Fruit: position of broadest part	Fruit: position de la partie la plus large	Frucht: Position des breitesten Teils	Fruto: posición de la parte más ancha		
QN	(c) towards stalk end	vers l'extrémité pédonculaire	zum Stielende hin	hacia el extremo peduncular		1
	at middle	au milieu	in der Mitte	en el medio	Washington Navel (SWO)	2
[247]	towards distal end	vers la partie distale	zum distalen Ende hin	hacia el extremo distal		3

					Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
	English	français	deutsch	español		
33. (+)	Fruit: general shape of proximal part (excluding neck, collar and depression at stalk end)	Fruit: forme générale de la partie proximale (à l'exclusion du col, de la collerette et de la dépression à l'extrémité pédonculaire)	Frucht: allgemeine Form des proximalen Teils (ohne Hals, Kragen und Einsenkung am Stielende)	Fruto: forma general de la parte proximal (excluidos el cuello, el collar y la depresión del extremo peduncular)		
PQ (c)	flattened	aplatie	abgeflacht	aplanada	Salustiana (SWO)	1
	slightly rounded	légèrement arrondie	leicht abgerundet	ligeramente redondeada	Valencia Late (SWO)	2
	strongly rounded	fortement arrondie	stark abgerundet	fuertemente redondeada		3
[249]	tapered	effilée	spitz	afilada		4
34. (*) (+)	Only varieties without fruit neck: Fruit: presence of depression at stalk end	Seulement les variétés dont le fruit ne présente pas de col: Fruit: présence d'une dépression à l'extrémité pédonculaire	Nur Sorten ohne Fruchthals: Frucht: Vorhandensein einer Einsenkung am Stielende	Sólo variedades con fruto sin cuello: Fruto: presencia de una depresión en el extremo peduncular		
QL (c)	absent	absente	fehlend	ausente	Sanguinelli (SWO)	1
[253]	present	présente	vorhanden	presente	Washington Navel (SWO)	9
35.	Only varieties without fruit neck: Fruit: depth of depression at stalk end	Seulement les variétés dont le fruit ne présente pas de col: Fruit: profondeur de la dépression à l'extrémité pédonculaire	Nur Sorten ohne Fruchthals: Frucht: Tiefe der Einsenkung am Stielende	Sólo variedades con fruto sin cuello: Fruto: profundidad de la depresión en el extremo peduncular		
QN (c)	shallow	peu profonde	flach	poco profunda	Washington Navel (SWO)	3
	medium	moyenne	mittel	media		5
[254]	deep	profonde	tief	profunda		7

					Example Varieties Exemples Beispielssorten Variedades ejemplos	Note/ Nota
	English	français	deutsch	español		
36.	Fruit: number of radial grooves at stalk end	Fruit: nombre de cannelures radiales à l'extrémité pédonculaire	Frucht: Anzahl radialer Furchen am Stielende	Fruto: número de acanaladuras radiales en el extremo peduncular		
QN	(c) absent or few	absentes ou peu nombreuses	fehlend oder sehr gering	ausentes o bajo	Valencia Late (SWO)	1
	intermediate	moyennement nombreuses	mittel	medio	Lanelate (SWO)	2
[257]	many	nombreuses	groß	alto		3
37.	Fruit: length of radial grooves at stalk end	Fruit: longueur des cannelures radiales à l'extrémité pédonculaire	Frucht: Länge der radialen Furchen am Stielende	Fruto: longitud de las acanaladuras radiales en el extremo peduncular		
QN	(c) short	courtes	kurz	cortas		3
	medium	moyennes	mittel	medianas		5
[258]	long	longues	lang	largas		7
38.	Fruit: presence of collar	Fruit: présence d'une corolle	Frucht: Vorhanden- sein eines Kragens	Fruto: presencia de un collar		
(+)						
QL	(c) absent	absente	fehlend	ausente	Salustiana (SWO)	1
[260]	present	présente	vorhanden	presente		9
39.	Fruit: general shape of distal part (excluding nipple, bulging of navel and depression at distal end)	Fruit: forme générale de la partie distale (à l'exclusion du mamelon, de la courbure du fruit secondaire et de la dépression à l'extrémité distale)	Frucht: allgemeine Form des distalen Teils (ohne Warze, Wölbung der sekundären Frucht und Einsenkung am distalen Ende)	Fruto: forma general de la parte distal (excluidos el mamelón o pezón, el abultamiento del ombligo y la depresión en el extremo distal)		
(+)						
QN	(c) flattened	aplatie	abgeflacht	aplanada	Hamlin (SWO)	1
	slightly rounded	légèrement arrondie	leicht abgerundet	ligeramente redondeada	Valencia Late (SWO)	2
[264]	strongly rounded	fortement arrondie	stark abgerundet	fuertemente redondeada		3

					Example Varieties Exemples Beispielssorten Variedades ejemplar	Note/ Nota
	English	français	deutsch	español		
40. (*) (+)	Fruit: presence of depression at distal end	Fruit: présence d'une dépression à l'extrémité distale	Frucht: Vorhandensein einer Einsenkung am distalen Ende	Fruto: presencia de una depresión en el extremo distal		
QL [265]	(c) absent	absente	fehlend	ausente	Valencia Late (SWO)	1
	present	présente	vorhanden	presente		9
41. (*)	Fruit: presence of areola	Fruit: présence d'une aréole	Frucht: Vorhandensein einer Areola	Fruto: presencia de un areola		
QL [270]	(c) absent	absente	fehlend	ausente	Valencia Late (SWO)	1
	incomplete	incomplète	unvollständig	incompleta		2
	complete	complète	vollständig	completa	Peret (SWO)	3
42. (+)	Fruit: type of areola	Fruit: type d'aréole	Frucht: Typ der Areola	Fruto: tipo de areola		
QL [271]	(c) smooth	régulière	glatt	lisa	Peret (SWO)	1
	grooved	cannelée	gerieft	acanalada		2
	ridged	annelée	geringelt	acrestada		3
43.	Fruit: diameter of areola	Fruit: diamètre de l'aréole	Frucht: Durchmesser der Areola	Fruto: diámetro de la areola		
QN [272]	(c) small	petit	klein	pequeño		3
	medium	moyen	mittel	medio	Peret (SWO)	5
	large	grand	groß	grande		7
44.	Fruit: diameter of stylar scar	Fruit: diamètre de la cicatrice stylique	Frucht: Durchmesser der Griffelnarbe	Fruto: diámetro de la cicatriz estilar		
QN [273]	(c) small	petit	klein	pequeño	Salustiana (SWO)	3
	medium	moyen	mittel	medio		5
	large	grand	groß	grande		7

					Example Varieties Exemples Beispielssorten Variedades ejemplos	Note/ Nota
	English	français	deutsch	español		
45.	Fruit: persistence of style	Fruit: persistance du style	Frucht: Persistenz des Griffels	Fruto: persistencia del estilo		
PQ	(c) none	aucune	fehlend	ninguna	Valencia Late (SWO)	1
	partial	partielle	teilweise gegeben	parcial	Sangre Oval (SWO)	2
[275]	total	totale	vollständig gegeben	total		3
46.	Fruit: presence of navel opening	Fruit: présence d'une ouverture du fruit secondaire	Frucht: Vorhandensein der Öffnung der sekundären Frucht	Fruto: presencia de un apertura del ombligo		
PQ	(c) absent	absente	fehlend	ausente	Ricalate (SWO)	1
	occasionally present	présente occasionnellement	zuweilen vorhanden	ocasionalmente presente	Navelate (SWO)	2
[276]	always present	toujours présente	immer vorhanden	siempre presente	Washington Navel (SWO)	3
47.	Fruit: diameter of navel opening	Fruit: diamètre de l'ouverture du fruit secondaire	Frucht: Durchmesser der Öffnung der sekundären Frucht	Fruto: diámetro de la apertura del ombligo		
QN	(c) small	petit	klein	pequeño	Navelate (SWO)	3
	medium	moyen	mittel	medio	Lanelate (SWO)	5
[277]	large	grand	groß	grande	Washington Navel (SWO)	7
48.	Fruit: bulging of navel	Fruit: courbure du fruit secondaire	Frucht: Wölbung der sekundären Frucht	Fruto: abultamiento del ombligo		
QN	(c) absent or weak	absente ou faible	fehlend	ausente o débil	Washington Navel (SWO)	1
	intermediate	intermédiaire	mittel	medio		2
[278]	strong	forte	stark	fuerte		3
49.	Fruit: presence of radial grooves at distal end	Fruit: présence de cannelures radiales à l'extrémité distale	Frucht: Vorhandensein radialer Furchen am distalen Ende	Fruto: presencia de acanaladuras radiales en el extremo distal		
QL	(c) absent	absentes	fehlend	ausentes	Valencia Late (SWO)	1
[279]	present	présentes	vorhanden	presentes	Salustiana (SWO)	9

					Example Varieties	
	English	français	deutsch	español	Exemples	Note/ Nota
					Beispielssorten	
50.	Fruit: color variegation	Fruit: panachure de la couleur	Frucht: Panchierung der Farbe	Fruto: variegación del color		
QL	(c) absent	absente	fehlend	ausente		1
[281]	present	présente	vorhanden	presente		9
51. (*)	Fruit surface: predominant color(s)	Surface du fruit: couleur(s) prédominance(s)	Fruchtoberfläche: Hauptfarbe(n)	Superficie del fruto: color(es) predominante(s)		
PQ	(c) yellow orange	orange-jaune	gelborange	naranja amarillento	Pinalate (SWO)	1
	(d) medium orange	orange moyen	mittelorange	naranja medio	Valencia Late (SWO)	2
	dark orange	orange foncé	dunkelorange	naranja oscuro	Washington Navel (SWO)	3
	orange red	rouge orangé	orangerot	rojo anaranjado	Navelate (SWO)	4
[282]	red	rouge	rot	rojo	Sanguinelli (SWO)	5
52.	Fruit surface: roughness	Surface du fruit: rugosité	Fruchtoberfläche: Rauheit	Superficie del fruto: rugosidad		
QN	(c) smooth	lisse	glatt	lisa	Sangre Doblefina (SWO)	3
	(d) medium	intermédiaire	mittel	media	Valencia Late (SWO)	5
[286]	rough	rugueuse	rauh	rugosa		7
53.	Fruit surface: size of oil glands	Surface du fruit: taille des glandes à huile	Fruchtoberfläche: Größe der Öldrüsen	Superficie del fruto: tamaño de las glándulas de aceite		
PQ	(c) all more or less the same size	toutes plus ou moins de la même taille	alle mehr oder weniger gleich groß	todas más o menos del mismo tamaño		1
[287]	(d) larger ones interspersed by smaller ones	grandes et petites intercalées	größere vermischt mit kleineren	glándulas grandes intercaladas con otras más pequeñas		2
54.	Fruit surface: size of larger oil glands	Surface du fruit: taille des glandes à huile les plus grosses	Fruchtoberfläche: Größe der größeren Öldrüsen	Superficie del fruto: tamaño de las glándulas de aceite más grandes		
QN	(c) small	petites	klein	pequeñas		3
	(d) medium	moyennes	mittel	medianas		5
[288]	large	grosses	groß	grandes		7

					Example Varieties Exemples Beispielssorten Variedades ejemplos	Note/ Nota
	English	français	deutsch	español		
55.	Fruit surface: conspicuousness of larger oil glands	Surface du fruit: netteté des glandes à huile les plus grosses	Fruchtoberfläche: Sichtbarkeit der größeren Öldrüsen	Superficie del fruto: visibilidad de las glándulas de aceite más grandes		
QN	(c) weak	faible	gering	débil	Valencia Late (SWO)	3
	(d) medium	moyenne	mittel	media	Bonanza (SWO)	5
[289]	strong	forte	stark	fuerte		7
56.	Fruit surface: presence of pitting and pebbling (+)	Surface du fruit: présence de dépression et de protubérance	Fruchtoberfläche: Vorhandensein von Grübchen und Körnern	Superficie del fruto: presencia de picado y granulado		
PQ	(c) pitting and pebbling absent	dépression et protubérance absentes	Grübchen und Körner fehlend	picado y granulado ausentes		1
	(d) pitting absent, pebbling present	dépression absente, protubérance présente	Grübchen fehlend, Körner vorhanden	picado ausente, granulado presente		2
	pitting present, pebbling absent	dépression présente, protubérance absente	Grübchen vorhanden, Körner fehlend	picado presente, granulado ausente		3
[290]	pitting and pebbling present	dépression et protubérance présentes	Grübchen und Körner vorhanden	picado y granulado presentes		4
57.	Varieties with fruit surface: pitting present only: Fruit surface: density of pitting	Variétés avec sur- face du fruit: pré- sence seulement de dépression: Surface du fruit: densité de la dépression	Sorten mit Frucht- oberfläche: nur Grübchen vorhan- den: Fruchtober- fläche: Dichte der Grübchen	Variedades con superficie del fruto: presencia solamente de picado: Superficie del fruto: densidad del picado		
QN	(c) sparse	éparse	locker	dispersa		3
	(d) medium	moyenne	mittel	media		5
[291]	dense	dense	dicht	densa		7

				Example Varieties Exemples Beispielssorten Variedades ejemplos	Note/ Nota
	English	français	deutsch	español	
58.	Varieties with fruit surface: pebbling present only: Fruit surface: density of pebbling	Variétés avec surface du fruit: présence seulement de protubérance: Surface du fruit: densité de la protubérance	Sorten mit Frucht-oberfläche: nur Körner vorhanden: Fruchtoberfläche: Dichte der Körner	Variedades con superficie del fruto: presencia solamente de granulado: Superficie del fruto: densidad del granulado	
QN	(c) sparse	éparse	locker	dispersa	3
	(d) medium	moyenne	mittel	media	5
[293]	dense	dense	dicht	densa	7
59.	Varieties with fruit surface: pebbling present only: Fruit surface: degree of pebbling	Variétés avec surface du fruit: présence seulement de protubérance: Surface du fruit: degré de protubérance	Sorten mit Frucht-oberfläche: nur Körner vorhanden: Fruchtoberfläche: Grad der Körnerbildung	Variedades con superficie del fruto: presencia solamente de granulado: Superficie del fruto: nivel de granulado	
QN	(c) weak	faible	gering	débil	3
	(d) medium	moyen	mittel	medio	5
[294]	strong	fort	stark	fuerte	7
60. (*)	Fruit rind: thickness	Écorce du fruit: épaisseur	Fruchtschale: Dicke	Corteza del fruto: espesor	
QN	(c) thin	fine	dünn	delgada	Navelate (SWO) 3
	(d) medium	moyenne	mittel	media	Valencia Late (SWO) 5
[295]	thick	épaisse	dick	gruesa	Newhall (SWO) 7
61.	Fruit rind: strength	Écorce du fruit: rigidité	Fruchtschale: Festigkeit	Corteza del fruto: vigor	
QN	(c) weak	faible	gering	débil	3
	(d) medium	moyenne	mittel	medio	5
[297]	strong	forte	groß	fuerte	7

					Example Varieties	
	English	français	deutsch	español	Exemples	Note/ Nota
					Beispielssorten	
62.	Fruit: color of albedo	Fruit: couleur de l'albédo	Frucht: Farbe der Albedo	Fruto: color del albedo		
PQ	(c) greenish	verdâtre	grünlich	verdosado		1
	white	blanc	weiß	blanco		2
	light yellow	jaune clair	hellgelb	amarillo claro		3
	light orange	orange clair	hellorange	naranja claro		4
	pink	rose	rosa	rosa		5
[300]	reddish	rougeâtre	rötlich	rojizo		6
63.	Fruit: differently colored specks in flesh	Fruit: taches de couleurs différentes dans la chair	Frucht: unterschiedlich gefärbte Flecken am Fleisch	Fruto: manchas de distinto color en la pulpa		
QL	(c) absent	absentes	fehlend	ausentes	Valencia Late (SWO)	1
[305]	(e) present	présentes	vorhanden	presentes	Sanguinelli (SWO)	9
64.	Fruit: bicolored segments	Fruit: segments bicolores	Frucht: zweifarbig Segmente	Fruto: gajos bicolores		
QL	(c) absent	absents	fehlend	ausentes	Valencia Late (SWO)	1
[306]	(e) present	présents	vorhanden	presentes	Sanguinelli (SWO)	9
65.	Only varieties with Fruit: bicolored segments: present: Fruit: distribution of red coloration	Seulement variétés avec Fruit: segments bicolores: présents: Fruit: distribution de la coloration rouge	Nur Sorten mit Frucht: zweifarbig Segmente: vorhanden: Frucht: Verteilung der Rotfärbung	Sólo variedades con Fruto: gajos bicolores: presentes: Fruto: distribución de la coloración roja		
QL	(c) evenly distributed	uniformément répartis	gleichmäßig verteilt	repartido uniforme	Tarocco (SWO)	1
[306.1]	(e) mainly marginate	principalement au bord	vorwiegend am Rand	principalmente marginal	Sanguinelli (SWO)	2

					Example Varieties Exemples Beispielssorten Variedades ejemplos	Note/ Nota
	English	français	deutsch	español		
66. <small>(*)</small>	Fruit: main color of flesh	Fruit: couleur principale de la chair	Frucht: Hauptfarbe des Fleisches	Fruto: color principal de la pulpa		
PQ	(c) light orange	orange clair	hellorange	naranja claro	Valencia Late (SWO)	1
	(e) medium orange	orange moyen	mittelorange	naranja medio	Washington Navel (SWO)	2
	dark orange	orange foncé	dunkelorange	naranja oscuro		3
	orange red	rouge orangé	orangerot	rojo anaranjado	Caracara (SWO)	4
[307]	red	rouge	rot	rojo	Sanguinelli (SWO)	5
67.	Fruit: bitterness of flesh	Fruit: amertume de la chair	Frucht: Bitterkeit des Fleisches	Fruto: amargor de la pulpa		
QL	(c) absent	absente	fehlend	ausente		1
[308]	(e) present	présente	vorhanden	presente		9
68.	Fruit: filling of core	Fruit : structure du cœur	Frucht: Ausfüllung des inneren Fruchtfleisches	Fruto: relleno del hueco central		
QN	(c) absent or very sparse	absente ou très lâche	fehlend oder sehr locker	ausente o muy laxo		1
	(e) sparse	lâche	locker	laxo		3
	medium	intermédiaire	mittel	medio	Washington Navel (SWO)	5
	dense	dense	dicht	denso	Salustiana (SWO)	7
[309]	very dense	très dense	sehr dicht	muy denso		9
69.	Fruit: diameter of core	Fruit: diamètre du cœur	Frucht: Durchmesser des inneren Fruchtfleisches	Fruto: diámetro del hueco central		
QN	(c) small	petit	klein	pequeño	Salustiana (SWO)	3
	(e) medium	moyen	mittel	medio	Valencia Late (SWO)	5
[310]	large	grand	groß	grande	Navelate (SWO)	7

					Example Varieties	
	English	français	deutsch	español	Exemples	Note/ Nota
					Beispielssorten	
70.	Fruit: presence of rudimentary segments	Fruit: présence de segments rudimentaires	Frucht: Vorhandensein von unvollständigen Segmenten	Fruto: presencia de gajos rudimentarios		
QN	(c) absent or weak	nulle ou faible	fehlend oder schwach	nula o débil	Valencia Late (SWO)	1
	(e) intermediate	intermédiaire	mittel	intermedia		2
[311]	strong	forte	stark	fuerte		3
71.	Fruit: number of well developed segments	Fruit: nombre de segments bien développés	Frucht: Anzahl gut entwickelter Segmente	Fruto: número de gajos bien desarrollados		
QN	(c) few	peu	gering	bajo	Navelate (SWO)	3
	(e) medium	moyen	mittel	medio	Sanguinelli (SWO)	5
[312]	many	beaucoup	groß	alto		7
72.	Fruit: coherence of adjacent segment walls	Fruit: adhérence des parois des segments adjacents	Frucht: Zusammenhalt der neben-einanderliegenden Segmentwände	Fruto: adherencia entre los gajos adyacentes		
QN	(c) weak	faible	gering	débil	Navelina (SWO)	3
	(e) medium	moyenne	mittel	media	Valencia Late (SWO)	5
[313]	strong	forte	stark	fuerte		7
73.	Fruit: strength of segment walls	Fruit: rigidité des parois des segments	Frucht: Festigkeit der Segmentwände	Fruto: firmeza de las paredes de los gajos		
QN	(c) weak	faible	schwach	débil	Navelate (SWO)	3
	(e) medium	moyenne	mittel	media	Valencia Late (SWO)	5
[314]	strong	forte	stark	fuerte	Berna (SWO)	7
74.	Fruit: length of juice vesicles	Fruit: longueur des vésicules de jus	Frucht: Länge der Saftbläschen	Fruto: longitud de las vesículas de jugo		
QN	(c) short	courtes	kurz	corta	Salustiana (SWO)	3
	(e) medium	moyennes	mittel	media		5
[315]	long	longues	lang	larga	Washington Navel (SWO)	7

					Example Varieties	
	English	français	deutsch	español	Exemples	Note/ Nota
					Beispielssorten	
75.	Fruit: thickness of juice vesicles	Fruit: épaisseur des vésicules de jus	Frucht: Dicke der Saftbläschen	Fruto: grosor de las vesículas de jugo		
QN	(c) thin	fines	dünn	delgadas		3
	(e) medium	moyennes	mittel	medianas		5
[316]	thick	épaisses	dick	gruesas		7
76.	Fruit: conspicuousness of juice vesicle walls	Fruit: netteté des parois des vésicules de jus	Frucht: Sichtbarkeit der Saftbläschenwände	Fruto: visibilidad de las paredes de las vesículas de jugo		
QN	(c) low	faible	gering	baja		3
	(e) medium	moyenne	mittel	media		5
[317]	high	forte	stark	alta		7
77.	Fruit: coherence of juice vesicles	Fruit: adhérence des vésicules de jus	Frucht: Zusammenhalt der Saftbläschen	Fruto: coherencia de las vesículas de jugo		
QN	(c) weak	faible	gering	débil		3
	(e) medium	moyenne	mittel	media		5
[318]	strong	forte	stark	fuerte		7
78. (*)	Fruit: presence of navel (viewed internally)	Fruit: présence de fruit secondaire (vue interne)	Frucht: Vorhandensein von sekundären Früchten (von innen gesehen)	Fruto: presencia de ombligo (visto internamente)		
PQ	(c) absent or very rare	absent ou très rare	fehlend oder sehr selten	ausente o muy rara	Valencia Late (SWO)	1
	occasionally present	présent occasionnellement	zuweilen vorhanden	ocasionalmente presente		2
[319]	always present	toujours présent	immer vorhanden	siempre presente	Washington Navel (SWO)	3
79.	Fruit: size of navel (viewed internally)	Fruit: taille du fruit secondaire (vue interne)	Frucht: Größe der sekundären Frucht (von innen gesehen)	Fruto: tamaño del ombligo visto internamente		
QN	(c) small	petit	klein	pequeño		3
	medium	moyen	mittel	medio	Washington Navel (SWO)	5
[320]	large	grand	groß	grande	Navelate (SWO)	7

					Example Varieties Exemples Beispielssorten Variedades ejemplos	Note/ Nota
	English	français	deutsch	español		
80.	Fruit: juiciness	Fruit: succulence	Frucht: Saftigkeit	Fruto: contenido de jugo		
QN	(c) low	faible	gering	bajo		3
	medium	moyenne	mittel	medio	Washington Navel (SWO)	5
[321]	high	élevée	hoch	alto	Salustiana (SWO)	7
81.	Fruit juice: total soluble solids	Jus du fruit: quantité de solides solubles	Fruchtsaft: Gehalt an löslicher Trockensubstanz	Jugo del fruto: cantidad de sólidos solubles		
QN	(c) low	faible	niedrig	baja	Valencia Late (SWO)	3
	medium	moyenne	mittel	media	Washington Navel (SWO)	5
[322]	high	élevée	hoch	alta	Navelate (SWO)	7
82.	Fruit juice: acidity	Jus du fruit: acidité	Fruchtsaft: Säure	Jugo del fruto: acidez		
QN	(c) very low	très faible	sehr gering	muy baja	Sucreña (SWO)	1
	low	faible	gering	baja		3
	medium	moyenne	mittel	media	Washington Navel (SWO)	5
[323]	high	forte	stark	alta	Valencia Late (SWO)	7
83.	Fruit: strength of fibre	Fruit: rigidité des fibres	Frucht: Festigkeit der Fasern	Fruto: vigor de la fibra		
QN	(c) weak	faible	schwach	débil	Salustiana (SWO)	3
	medium	moyenne	mittel	medio	Washington Navel (SWO)	5
[324]	strong	forte	stark	fuerte		7
84. (+)	Fruit: number of seeds (open pollination)	Fruit: nombre de pépins (fécondation libre)	Frucht: Anzahl Samen (frei abhängig)	Fruto: número de semillas (polinización libre)		
QN	(c) absent or very few	absents ou très peu nombreux	fehlend oder sehr gering	ausentes o muy bajo	Salustiana (SWO)	1
	few	peu nombreux	gering	bajo	Valencia Late (SWO)	3
	medium	moyen	mittel	medium		5
[326]	many	nombreux	groß	alto	Pineapple (SWO)	7

					Example Varieties Exemples Beispielssorten Variedades ejemplos	Note/ Nota
	English	français	deutsch	español		
85. (*)	Seed: polyembryony	Pépin: polyembryonnie	Samen: Polyembryonie	Semilla: poliembrionía		
QL (f)	absent	absente	fehlend	ausente		1
[327]	present	présente	vorhanden	presente	Valencia Late (SWO)	9
86.	Seed: length	Pépin: longueur	Samen: Länge	Semilla: longitud		
QN (f)	short	court	kurz	corta		3
	medium	moyen	mittel	media		5
[328]	long	long	lang	larga		7
87.	Seed: width	Pépin: largeur	Samen: Breite	Semilla: anchura		
QN (f)	narrow	étroit	schmal	estrecha		3
	medium	moyen	mittel	media		5
[329]	broad	large	breit	ancha		7
88.	Seed: surface	Pépin: surface	Samen: Oberfläche	Semilla: superficie		
QL (f)	smooth	lisse	glatt	lisa		1
[330]	wrinkled	ridée	runzlig	rugosa		2
89.	Varieties with seed: surface wrinkled only: Seed: promi- nence of wrinkles	Variétés avec pépin: surface ridée seu- lement: Pépin: pro- éminence des rides	Sorten mit Samen: nur runzlige Ober- fläche: Samen: Ausprägung der Runzeln	Variedades con semilla: superficie solamente rugosa: Semilla: prominen- cia de la rugosidad		
QN (f)	weak	faible	schwach	débil		3
	medium	moyenne	mittel	media		5
[331]	strong	forte	stark	fuerte		7
90.	Seed: external color	Pépin: couleur externe	Samen: Außenfarbe	Semilla: color externo		
PQ (f)	greenish	verdâtre	grünlich	verdosado		1
	whitish	blanchâtre	weißlich	blanquecino	Comuna (SWO)	2
	yellowish	jaunâtre	gelblich	amarillento		3
	pinkish	rosâtre	rosa	rosado		4
[332]	brownish	brunâtre	bräunlich	amarronado		5

					Example Varieties Exemples Beispielssorten Variedades ejemplos	Note/ Nota
	English	français	deutsch	español		
91.	Seed: color of inner seed coat	Pépin: couleur du tégument interne	Samen: Farbe der inneren Samenschale	Semilla: color de la cubierta interna		
PQ	(f)	white	blanc	weiß	blanco	Sucreña (SWO) 1
		light yellow	jaune clair	hellgelb	amarillo claro	2
		light brown	marron clair	hellbraun	marrón claro	3
		medium brown	marron moyen	mittelbraun	marrón medio	Comuna (SWO) 4
		dark brown	marron foncé	dunkelbraun	marrón oscuro	5
		red	rouge	rot	rojo	6
[333]		purple	violet	purpur	púrpura	7
92.	Only varieties with seed: polyembryony present: Seed: color of cotyledons	Seulement variétés avec pépin: polyembryonie présente: Pépin: couleur des cotylédons	Nur Sorten mit Samen: Polyembryonie vorhanden: Samen: Farbe der Kotyledonen	Sólo variedades con semilla: poliembriónia presente: Semilla: color de los cotiledones		
PQ	(f)	white	blanc	weiß	blanco	Comuna (SWO) 1
		cream	crème	cremefarben	crema	2
		light green	vert clair	hellgrün	verde claro	3
[334]		dark green	vert foncé	dunkelgrün	verde oscuro	4
93. (*)	Time of maturity of fruit for consumption	Époque de maturité du fruit pour la consommation	Zeitpunkt der Genußreife	Época de madurez del fruto para su consumo		
QN	early	précoce	früh	temprana	Navelina (SWO)	3
	medium	moyenne	mittel	media	Salustiana (SWO)	5
[336]	late	tardive	spät	tardía	Valencia Late (SWO)	7
94. (*)	Fruit: parthenocarpic	Fruit: parthénocarpie	Frucht: Parthenokarpie	Fruto: partenocarpia		
QL	(c)	absent	absente	fehlend	ausente	Comuna (SWO) 1
[337]		present	présente	vorhanden	presente	Washington Navel (SWO) 9

					Example Varieties	
	English	français	deutsch	español	Exemples	Note/ Nota
					Beispielssorten	
95. (+)	Plant: self-incompatibility	Plante: auto-incompatibilité	Pflanze: Selbst-inkompatibilität	Planta: auto-incompatibilidad		
QL	absent	absente	fehlend	ausente	Pineapple (SWO)	1
[338]	present	présente	vorhanden	presente		9

8. Explanations on the Table of Characteristics

8.1 *Explanations covering several characteristics*

Characteristics containing the following key in the second column of the Table of Characteristics should be examined as indicated below:

(a) Leaf: Observations on the leaf should be made on fully developed leaves on the middle third of the youngest spring flush branch sections not showing signs of active growth.

(b) Flower: Unless otherwise indicated, observations on the flower bud and the flower should be made on the terminal flower bud and flower, at the time of full flowering of the variety.

Observations on the open flower should be made on the first day of opening.

(c) Fruit: Observations on the fruit should be made at the stage of optimum ripeness. The fruit should be tested weekly and harvested as soon as this stage has been reached.

All fruits for observation should be taken from the periphery of the tree and fruit misformed as a result of clustering should not be sampled.

(d) Fruit surface and fruit rind: Observations on the fruit surface and on the fruit rind should be made at the middle, between the base and apex of the fruit.

(e) Fruit flesh: Observations on the flesh of the fruit should be made on a cross section through the middle of the fruit.


(f) Seed: Observations on the seed should be made on the fresh seed.

8.2 Explanations for individual characteristics


Ad. 2 ([202]): Tree: growth habit

The observation on the growth habit of the tree should be made immediately after harvest.

Ad. 14 ([224]): Leaf blade: shape of apex



Ad. 15 ([225]): Leaf blade: emargination at tip



Ad. 26 ([239]): Anther: pollen viability


There is variability during development of the floral bud. It must be observed during the period of full flowering. From the two years of observations, the highest value should be taken, as this would indicate the highest potential for pollination.

Method to determine the percentage of pollen viability:


The pollen should be collected when the petals begin to open (but with the anthers closed). The anthers should be introduced into a Petri dish and placed inside a silica gel dryer at room temperature, for 20-48 hours of darkness. When the anthers are open they should be moved to an 8 °C chamber with a 70-80 % Relative Humidity for one hour. Afterwards, the pollen should be brushed onto a microscope slide with 2 ml of Brewbaker medium (Brewbaker and Kwack. 1963). Finally, the microscope slide should be placed in a 24 °C chamber with a 75 % RH for 20 hours.

The percentage of pollen fertility is calculated as the average of germinated pollen grains observed with a binocular microscope in 15 visual fields from 2 different microscope slides.


Ad. 33 ([249]): Fruit: general shape of proximal part (excluding neck, collar and depression at stalk end)




Ad. 34 ([253]): Only varieties without fruit neck: Fruit: presence of depression at stalk end




Ad. 38 ([260]): Fruit: presence of collar




Ad. 39 ([264]): Fruit: general shape of distal part (excluding nipple, bulging of navel and depression at distal end)



Ad. 40 ([265]): Fruit: presence of depression at distal end




Ad. 42 ([271]): Fruit: type of areola



Ad. 56 ([290]): Fruit surface: presence of pitting and pebbling

Observations should be made on the proximal half of the fruit.



Ad. 84 ([326]): Fruit: number of seeds (open pollination)

Open pollination means natural pollination between trees of any variety.

Ad. 95 ([338]): Plant: self-incompatibility

A variety is self-incompatible when the fertile pollen of its own flower or of other flowers of the same variety is not able to fertilize the ovary.

LIST OF EXAMPLE VARIETIES FOR ORANGES

Variety denomination	Subgroup
Bonanza	SWO
Caracara	SWO
Comuna	SWO
Hamlin	SWO
Lanelate	SWO
Navel Mas Baró	SWO
Navelate	SWO
Navelina	SWO
Newhall	SWO
Olinda	SWO
Peret	SWO
Pineapple	SWO
Pinalate	SWO
Ricalate	SWO
Salustiana	SWO
Sangre Doblefina	SWO
Sangre Oval	SWO
Sanguinelli	SWO
Sucreña	SWO
Summer Navel	SWO
Sweet Navel	SWO
Tarocco	SWO
Valencia Late	SWO
Washington Navel	SWO

9. Literature

Alexander, D. McE., 1983: "Some citrus species and varieties in Australia," Commonwealth Scientific and Industrial Research Organization, Australia, 64 pp.

Blondel, L., 1978: Botanical classification of species of the genus Citrus, Fruits 33 (11): pp. 695 - 720.

Bono, R., Soler, J., Fernandez de Cordova, L., 1986: "Variedades de agrios cultivadas en España". Generalitat Valenciana, 70 pp.

Brewbaker, J.L. and Kwack, B.H. 1963. The essential role of calcium ion in pollen germination and pollen tube growth. Amer. Jour. Botany. 50: 859-865.

Damigella, P., Tribulato, E., Calabrese, F., Crescimanno, F.G., Continella, G., 1980: "Gli Agrumi," Cultivar. R.E.D.A., Roma, Italy, pp. 9 - 70.

Ortiz Marcide, J.M., 1985: "Nomenclatura botánica de los cítricos". Levante Agrícola nº 259-260, pp. 71-79.

Ray, R., Walheim, L., 1980: "Citrus: How to select, grow and enjoy," HP Books, Tucson, USA, pp. 41 - 115.

Reuther, W. (Editor), 1973: "The Citrus Industry," Volume III, University of California, Division of Agricultural Sciences, 528 pp.

Reuther, W., Batchelor, L.D., Webber, H.J. (Editors), 1968: "The Citrus Industry," Volume II, University of California, Division of Agricultural Sciences, 398 pp.

Reuther, W., Webber, H.J., Batchelor, L.D. (Editors), 1967: "The Citrus Industry," Volume I, University of California, Division of Agricultural Sciences, 611 pp.

Saunt, J., 1990: "Citrus varieties of the world: an illustrated guide," Sinclair International Ltd., Norwich, England, 126 pp.

Soler, J., 1999: "Reconocimiento de variedades de cítricos en campo", Consellería de Agricultura, Pesca y Alimentación, Serie Divulgación Técnica, 191 pp.

Spina, P., Russo, F., Geraci, G., Martelli, S., 1980: "Schede per il registro varietale dei fruttiferi I - ARANCIO e MANDARINO," Ministero Agricoltura e Foreste - S.O.I., Roma, Italy, 92 pp.

Thornton, I.R., El-Zeftawi, B.M., 1983: "Culture of irrigated citrus fruits," Government Printer, State of Victoria, Australia, pp. 12 -25.

Zaragoza, S., Navarro, L., Cebolla, V., 1997: "Evaluation of the field Collection through the germo data-base," Proc. Sectorial meeting of the mediterranean citrus network (MECINET) en global cooperation for citrus germplasm conservation and use, 147-148, Acireale-Catania, Italia.

Zaragoza, S., Trenor, I., Alonso, E., Medina, A., Pina, J.A., Navarro, L., 1995: "Evaluación de la colección de variedades del Banco de Germoplasma de Cítricos del IVIA: Planteamiento y primeros resultados generales". Levante Agrícola nº 331, pp. 145-149.

10. Technical Questionnaire

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
		Application date: (not to be filled in by the applicant)
TECHNICAL QUESTIONNAIRE to be completed in connection with an application for plant breeders' rights		
<p>1. Subject of the Technical Questionnaire</p> <p>(a) Subgroup: (i) SOR ---- [] (ii) SWO ---- [] (iii) HOR ---- []</p> <p>(b) Species (please specify):</p>		
<p>2. Applicant: Name</p> <p>Address</p> <p>Telephone No.</p> <p>Fax No.</p> <p>E-mail address</p> <p>Breeder (if different from applicant)</p>		
<p>3. Proposed denomination and breeder's reference</p> <p>Proposed denomination (if available)</p> <p>Breeder's reference</p>		

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

4. Information on the breeding scheme and propagation of the variety

4.1 Breeding Scheme

4.1.1 Variety resulting from:

- (a) controlled cross []
(please state parent varieties:)
.....)
- (b) partially unknown cross []
(please state known parent variety(ies):)
.....)
- (c) totally unknown cross []

4.1.2 Mutation []
(please state parent variety:)

4.1.3 Discovery []
(please state where, when and how developed:)
.....)

4.1.4 Other []
(please provide details:)
.....)

4.2 Method of Propagating the Variety

4.3 Virus status

4.3.1 The plant material is virus-free []

4.3.2 The plant material is virus tested []
(indicate against which viruses:)
.....)

4.3.3 The virus status is unknown []

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

5. Characteristics of the variety to be indicated (the number in brackets refers to the corresponding characteristic in Test Guidelines; please mark the note which best corresponds).

Characteristics	Example Varieties	Note
5.1 Fruit: length (29)		
short	Comuna (SWO)	3[]
medium	Valencia Late (SWO)	5[]
long	Newhall (SWO)	7[]
5.2 Fruit: diameter (30)		
small	Sanguinelli (SWO)	3[]
medium	Valencia Late (SWO)	5[]
large	Washington Navel (SWO)	7[]
5.3 Fruit surface: predominant color(s) (51)		
yellow orange	Pinalate (SWO)	1[]
medium orange	Valencia Late (SWO)	2[]
dark orange	Washington Navel (SWO)	3[]
orange red	Navelate (SWO)	4[]
red	Sanguinelli (SWO)	5[]
5.4 Fruit: main color of flesh (66)		
light orange	Valencia Late (SWO)	1[]
medium orange	Washington Navel (SWO)	2[]
dark orange		3[]
orange red	Caracara (SWO)	4[]
red	Sanguinelli (SWO)	5[]

TECHNICAL QUESTIONNAIRE	Page {x} of {y}	Reference Number:
-------------------------	-----------------	-------------------

7. Additional information which may help in the examination of the variety

7.1 In addition to the information provided in sections 5 and 6, are there any additional characteristics which may help to distinguish the variety?

Yes [] No []

If yes, please provide details:.....
.....

7.2 Special conditions for the examination of the variety

7.2.1 Are there any special conditions for growing the variety or conducting the examination?

Yes [] No []

If yes, please provide details:

7.3 Other information

A representative color photograph of the variety should accompany the Technical Questionnaire

8. Authorization for release

(a) Does the variety require prior authorization for release under legislation concerning the protection of the environment, human and animal health?

Yes [] No []

(b) Has such authorization been obtained?

Yes [] No []

If the answer to (b) is yes, please attach a copy of the authorization.

9. I hereby declare that, to the best of my knowledge, the information provided in this form is correct:

Applicant's name

Signature

Date

ANNEX / ANNEXE / ANLAGE / ANEXO

CITRUS L.

**OVERALL TABLE OF CHARACTERISTICS / TABLEAU GÉNÉRAL DES CARACTÈRES /
ALLGEMEINE MERKMALSTABELLE / CONJUNTO GLOBAL DE CARACTERES**

- ◆ **Group 1:** MANDARINS and their hybrids / **Groupe 1 :** MANDARINIER et ses hybrides /
Gruppe 1 : MANDARINEN und ihre Hybriden / **Grupo 1:** MANDARINO y sus híbridos **TG/201/1 Rev. Corr.**
- ◆ **Group 2:** ORANGES and their hybrids / **Groupe 2 :** ORANGER et ses hybrides /
Gruppe 2 : ORANGEN und ihre Hybriden / **Grupo 2:** NARANJO y sus híbridos **TG/202/1 Rev. 2**
- ◆ **Group 3:** LEMONS and LIMES and their hybrids / **Groupe 3 :** CITRONNIER et LIMETTIER et leurs hybrides /
Gruppe 3 : ZITRONEN und LIMETTEN und ihre Hybriden / **Grupo 3:** LIMÓN y LIMA y sus híbridos **TG/203/1 Rev. Corr.**
- ◆ **Group 4:** GRAPEFRUIT and PUMMELO and their hybrids / **Groupe 4 :** POMELO et PAMPLEMOUSSIER et leurs hybrides /
Gruppe 4 : GRAPEFRUIT und PAMPELMUSE und ihre Hybriden / **Grupo 4:** POMELO Y PUMMELO y sus híbridos **TG/204/1 Rev. 2**
- ◆ **Group 5:** TRIFOLIATE ORANGE and its hybrids / **Groupe 5 :** ORANGER TRIFOLIÉ et ses hybrides/
Gruppe 5 : DREIBLÄTTRIGE ORANGE und ihre Hybriden / **Grupo 5:** NARANJO TRIFOLIADO y sus híbridos **TG/83/4 Rev. Corr.**

	Group/Groupe/Gruppe/Grupo					English	français	deutsch	español	Note/ Nota	Group/Groupe/Gruppe/Grupo				
	1	2	3	4	5						1	2	3	4	5
201. QL	1	1	1	1		Ploidy	Ploidie	Ploidie	Ploidía						
						diploid	diploïde	diploid	diploide	2					
						triploid	triploïde	triploid	triploide	3					
						tetraploid	tétraploïde	tetraploid	tetraploide	4					
202. (+) PQ	2	2	2	2	1	Tree: growth habit	Arbre: port	Baum: Wuchstyp	Árbol: porte						
	(*)	(*)	(*)	(*)	(*)	upright	droit	aufrecht	erguido	1					
						spreading	étalé	breitbuschig	abierto	2					
						drooping	retombant	hängend	colgante	3					
203. QN	3	3	3	3	2	Tree: density of spines	Arbre: densité des épines	Baum: Dichte der Stacheln	Árbol: densidad de las espinas						
						absent or sparse	absentes ou éparses	fehlend oder locker	ausentes o laxa	1					
						intermediate	intermédiaires	mittel	media	2					
						dense	denses	dicht	densa	3					
204. QN	4	4	4	4	3	Tree: length of spines	Arbre: longueur des épines	Baum: Länge der Stacheln	Árbol: longitud de las espinas						
						short	courtes	kurz	cortas	3					
						medium	moyennes	mittel	medias	5					
						long	longues	lang	largas	7					

	Group/Groupe/Gruppe/Grupo					English	français	deutsch	español	Note/Nota	Group/Groupe/Gruppe/Grupo				
	1	2	3	4	5						1	2	3	4	5
205. QN					4	Branch: expression of lenticels	Branche: expression des lenticelles	Zweig: Ausprägung der Lentizellen	Rama: expresión de las lenticelas						
						weak	faible	gering	débil	3					
						medium	moyenne	mittel	media	5					
						strong	forte	stark	fuerte	7					
206. QL		5	5	5		Young leaf: presence of anthocyanin coloration	Jeune feuille: présence de pigmentation anthocyanique	Junges Blatt: Vorhandensein von Anthocyanfärbung	Hoja joven: presencia de pigmentación antociánica						
		(*)	(*)	(*)		absent	absente	fehlend	ausente	1					
						present	présente	vorhanden	presente	9					
207. QN		6	6	6		Young leaf: intensity of anthocyanin coloration	Jeune feuille: intensité de la pigmentation anthocyanique	Junges Blatt: Intensität der Anthocyanfärbung	Hoja joven: intensidad de la pigmentación antociánica						
						weak	faible	gering	débil	3					
						medium	moyenne	mittel	media	5					
						strong	forte	stark	fuerte	7					
208. PQ					7 (*)	Leaf: caducity	Feuille: persistance	Blatt: frühzeitiges Abfallen	Hoja: caducidad						
						perennial	persistante	überdauernd	perenne	1					
						partially caducous	partiellement caduque	teilweise frühzeitig abfallend	parcialmente caduca	2					
						caducous	caduque	frühzeitig abfallend	caduca	3					

	Group/Groupe/Gruppe/Grupo					English	français	deutsch	español	Note/Nota	Group/Groupe/Gruppe/Grupo				
	1	2	3	4	5						1	2	3	4	5
209. PQ					8 (*)	Leaf: number of leaflets one only variable three only	Feuille: nombre de folioles une seule variable trois seulement	Blatt: Anzahl Teilblätter nur eines variabel nur drei	Hoja: número de foliolos sólo uno variable sólo tres						
210. QN	5	5	7	7	9	Leaf blade: length (apical leaflet in case of compound leaf) short medium long	Limbe: longueur (foliole apicale en cas de feuille composée) court moyen long	Blattspreite: Länge (apikales Teilblatt bei zusammengesetztem Blatt) kurz mittel lang	Limbo: longitud (foliolo atípico en caso de hoja compuesta) corto medio largo						
211. QN	6	6	8	8	10	Leaf blade: width (as for 210) narrow medium broad	Limbe: largeur (comme pour 210) étroit moyen large	Blattspreite: Breite (wie für 210) schmal mittel breit	Limbo: anchura (como para 210) estrecho medio ancho						
212. QN	7	7	9	9	11	Leaf blade: ratio length/width (as for 210) small medium large	Limbe: rapport longueur/largeur (comme pour 210) faible moyen élevé	Blattspreite: Verhältnis Länge/Breite (wie für 210) klein mittel groß	Limbo: relación longitud/anchura (como para 210) pequeña media grande						

	Group/Groupe/Gruppe/Grupo					English	français	deutsch	español	Note/Nota	Group/Groupe/Gruppe/Grupo				
	1	2	3	4	5						1	2	3	4	5
213. QN				12	Leaf blade: length (lateral leaflet in case of compound leaf)	Limbe: longueur (foliole latérale en cas de feuille composée)	Blattspreite: Länge (seitliches Teilblatt bei zusammengesetztem Blatt)	Limbo: longitud (foliololo lateral en caso de hoja compuesta)							
					short	court	kurz	corto	3						
					medium	moyen	mittel	medio	5						
					long	long	lang	largo	7						
214. QN				13	Leaf blade: width (as for 213)	Limbe: largeur (comme pour 213)	Blattspreite: Breite (wie für 213)	Limbo: anchura (como para 213)							
					narrow	étroit	schmal	estrecho	3						
					medium	moyen	mittel	medio	5						
					broad	large	breit	ancho	7						
215. QN				14	Leaf blade: ratio length/width (as for 213)	Limbe: rapport longueur/largeur (comme pour 213)	Blattspreite: Verhältnis Länge/Breite (wie für 213)	Limbo: relación longitud/anchura (como para 213)							
					small	faible	klein	pequeña	3						
					medium	moyen	mittel	media	5						
					large	élévé	groß	grande	7						

	Group/Groupe/Gruppe/Grupo					English	français	deutsch	español	Note/Nota	Group/Groupe/Gruppe/Grupo					
	1	2	3	4	5						1	2	3	4	5	
216.					15	Leaf blade: ratio length of blade of apical leaflet/length of blade of lateral leaflet	Limbe: rapport longueur du limbe de la foliole apicale/longueur du limbe de la foliole latérale	Blattspreite: Verhältnis Länge der Spreite des apikalen Teilblattes/ Länge der Spreite des seitlichen Teilblattes	Limbo: relación entre la longitud del limbo del foliolo apical y la longitud del limbo del foliolo lateral							
QN						small	faible	klein	pequeña	3						
						medium	moyen	mittel	media	5						
						large	élevé	groß	grande	7						
217.	8	8	10	10	16	Leaf blade: shape in cross section (as for 210)	Limbe: forme en section transversale (comme pour 210)	Blattspreite: Form im Querschnitt (wie für 210)	Limbo: forma en sección transversal (como para 210)							
QN						straight or weakly concave	droit ou légèrement concave	gerade oder leicht konkav	recto o ligeramente cóncavo	1						
						intermediate	intermédiaire	mittel	intermedio	2						
						strongly concave	fortement concave	stark konkav	fuertemente cóncavo	3						
218.	9	9	11	11	17	Leaf blade: twisting	Limbe: torsion	Blattspreite: Drehung	Limbo: torsión							
QN						absent or weak	absente ou faible	fehlend oder gering	ausente o débil	1						
						intermediate	intermédiaire	mittel	media	2						
						strong	forte	stark	fuerte	3						
219.	10	10		12	18	Leaf blade: blistering	Limbe: cloquère	Blattspreite: Blasigkeit	Limbo: abullonado o ampollado							
QN						absent or weak	absente ou faible	fehlend oder gering	ausente o débil	1						
						intermediate	intermédiaire	mittel	medio	2						
						strong	forte	stark	fuerte	3						

	Group/Groupe/Gruppe/Grupo					English	français	deutsch	español	Note/ Nota	Group/Groupe/Gruppe/Grupo				
	1	2	3	4	5						1	2	3	4	5
220. QN	11	11	12	13	19	Leaf blade: green color	Limbe: couleur verte	Blattspreite: Grünfärbung	Limbo: color verde						
						light	claire	hell	claro	3					
						medium	moyenne	mittel	medio	5					
						dark	foncée	dunkel	oscuro	7					
221. QN				14	20	Leaf blade: pubescence on lower side	Limbe: pilosité sur la face inférieure	Blattspreite: Behaarung an der Unterseite	Limbo: pubescencia en el envés						
						absent or weak	absente ou faible	fehlend oder gering	ausente o débil	1					
						intermediate	intermédiaire	mittel	media	2					
						strong	forte	stark	fuerte	3					
222. QN	12	12	13	15	21	Leaf blade: undulation of margin	Limbe: ondulation du bord	Blattspreite: Randwellung	Limbo: ondulación del borde						
						absent or weak	absente ou faible	fehlend oder gering	ausente o débil	1					
						intermediate	intermédiaire	mittel	media	2					
						strong	forte	stark	fuerte	3					
223. PQ	13	13	14	16	22 (*)	Leaf blade: incisions of margin	Limbe: incisions du bord	Blattspreite: Randeinschnitte	Limbo: incisiones del borde						
						absent	absentes	fehlend	ausentes	1					
						crenate	crénelées	gekerbt	crenadas	2					
						dentate	dentelées	gezähnt	dentadas	3					

	Group/Groupe/Gruppe/Grupo					English	français	deutsch	español	Note/ Nota	Group/Groupe/Gruppe/Grupo				
	1	2	3	4	5						1	2	3	4	5
224. PQ	14	14	15	17	23 (*)	Leaf blade: shape of apex	Limbe: forme de l'extrémité	Blattspreite: Form der Spitze	Limbo: forma del ápice						
						acuminate	acuminée	mit aufgesetzter Spitze	acuminado	1					
						acute	pointue	spitz	agudo	2					
						obtuse	obtuse	stumpf	obtuso	3					
						rounded	arrondie	abgerundet	redondeado	4					
225. QL	15	15	16	18	24	Leaf blade: emargination at tip	Limbe: échancrure à l'extrémité	Blattspreite: Einkerbung an der Spitze	Limbo: emarginado en la parte superior						
						absent	absente	fehlend	ausente	1					
						present	présente	vorhanden	presente	9					
226. QN	16	16	17	19	25	Petiole: length	Pétiole: longueur	Blattstiellänge	Pecíolo: longitud						
						short	court	kurz	corto	3					
						medium	moyen	mittel	medio	5					
						long	long	lang	largo	7					
227. QL	17	17	18	20	26 (*)	Petiole: presence of wings	Pétiole: présence d'ailes	Blattstiellänge	Pecíolo: presencia de alas						
						absent	absentes	fehlend	ausentes	1					
						present	présentes	vorhanden	presentes	9					

	Group/Groupe/Gruppe/Grupo					English	français	deutsch	español	Note/Nota	Group/Groupe/Gruppe/Grupo				
	1	2	3	4	5						1	2	3	4	5
228. QN	18	18	19	21	27	<u>Varieties with petiole wings present only:</u> Petiole: width of wings	<u>Seulement les variétés présentant des ailes au pétiole:</u> Pétiole: largeur des ailes narrow medium broad	<u>Nur Sorten mit vorhandenen Flügeln am Blattstiel:</u> Blattstiel: Breite der Flügel schmal mittel breit	<u>Sólo variedades con alas presentes en el pecíolo:</u> Pecíolo: anchura de las alas estrechas medias anchas						
229. QL			20	22	28	Flower bud: presence of anthocyanin coloration	Bouton floral: présence de pigmentation anthocyane absent present	Blütenknospe: Vorhandensein von Anthocyanfärbung fehlend vorhanden	Yema floral: presencia de pigmentación antociánica ausente presente						
230. QN			21	23	29	Flower bud: intensity of anthocyanin coloration	Bouton floral: intensité de la pigmentation anthocyane weak medium strong	Blütenknospe: Intensität der Anthocyanfärbung schwach mittel stark	Yema floral: intensidad de la pigmentación antociánica débil media fuerte						
231. QN	19	19	22	24	30	Flower: diameter of calyx	Fleur: diamètre du calice small medium large	Blüte: Durchmesser des Kelches klein mittel groß	Flor: diámetro del cáliz pequeño medio grande						

	Group/Groupe/Gruppe/Grupo					English	français	deutsch	español	Note/ Nota	Group/Groupe/Gruppe/Grupo				
	1	2	3	4	5						1	2	3	4	5
232. QN	20	20	23	25	31	Flower: length of petal	Fleur: longueur du pétales	Blüte: Länge des Blütenblattes	Flor: longitud del pétalo						
						short	court	kurz	corto	3					
						medium	moyen	mittel	medio	5					
						long	long	lang	largo	7					
233. QN	21	21	24	26	32	Flower: width of petal	Fleur: largeur du pétales	Blüte: Breite des Blütenblattes	Flor: anchura del pétalo						
						narrow	étroit	schmal	estrecho	3					
						medium	moyen	mittel	medio	5					
						broad	large	breit	ancho	7					
234. QN	22	22	25	27	33	Flower: ratio length/width of petal	Fleur: rapport longueur/largeur du pétales	Blüte: Verhältnis Länge/Breite des Blütenblattes	Flor: relación longitud/anchura del pétalo						
						small	faible	klein	pequeña	3					
						medium	moyen	mittel	media	5					
						large	élevé	groß	grande	7					
235. QN	23	23	26	28	34	Flower: length of stamens	Fleur: longueur des étamines	Blüte: Länge der Staubfäden	Flor: longitud de los estambres						
						short	courtes	kurz	cortos	3					
						medium	moyennes	mittel	medios	5					
						long	longues	lang	largos	7					

	Group/Groupe/Gruppe/Grupo					English	français	deutsch	español	Note/ Nota	Group/Groupe/Gruppe/Grupo				
	1	2	3	4	5						1	2	3	4	5
236. QL		24	27			Flower: basal union of stamens	Fleur: union basale des étamines	Blüte: Vereinigung der Staubfäden an der Basis	Flor: unión basal de los estambres						
						absent	absente	fehlend	ausente	1					
						present	présente	vorhanden	presente	9					
237. QL					35	Stamen: color of filament	Étamine: couleur du filament	Staubfaden: Farbe des Fadens	Estambre: color del filamento						
						white	blanc	weiß	blanco	1					
						pink	rose	rosa	rosa	2					
238. PQ	24	25	28	29	36	Anther: color	Anthère: couleur	Anthere: Farbe	Antera: color						
						white	blanc	weiß	blanco	1					
						light yellow	jaune clair	hellgelb	amarillo claro	2					
						medium yellow	jaune moyen	mittelgelb	amarillo medio	3					
239.- 1 QL			29		37	Anther: viable pollen	Anthère: pollen viable	Anthere: keimfähiger Pollen	Antera: polen viable						
						absent	absent	fehlend	ausente	1					
						present	présent	vorhanden	presente	9					
239.- 2 (+) QN		26		30		Anther: pollen viability	Anthère: viabilité du pollen	Anthere: Keimfähigkeit des Pollen	Antera: viabilidad del polen						
						absent or low	absente ou faible	fehlend oder gering	ausente o baja	1					
						medium	moyenne	mittel	media	2					
						high	forte	stark	alta	3					

	Group/Groupe/Gruppe/Grupo					English	français	deutsch	español	Note/ Nota	Group/Groupe/Gruppe/Grupo				
	1	2	3	4	5						1	2	3	4	5
240. QN	26	27	30	31	38	Style: length short medium long	Style: longueur court moyen long	Griffel: Länge kurz mittel lang	Estilo: longitud corto medio largo						
241. PQ		28				Style: shape straight arched kinked	Style: forme droit arqué coudé	Griffel: Form gerade nach unten gebogen geknickt	Estilo: forma recto arqueado sinuoso						
242. QL					39	Ovary: pubescence absent present	Ovaire: pilosité absente présente	Fruchtknoten: Behaarung fehlend vorhanden	Ovario: pubescencia ausente presente						
243. QL	27		31	32		Infructescence: cluster-ing of fruits absent present	Fructification: formation de grappes absente présente	Fruchtstand: Früchte in Büschen fehlend vorhanden	Infructescencia: arra-cimado de los frutos ausente presente						
244. QN	28 (*)	29 (*)	32 (*)	33 (*)	40 (*)	Fruit: length short medium long	Fruit: longueur court moyen long	Frucht: Länge kurz mittel lang	Fruto: longitud corto medio largo						

	Group/Groupe/Gruppe/Grupo					English	français	deutsch	español	Note/ Nota	Group/Groupe/Gruppe/Grupo				
	1	2	3	4	5						1	2	3	4	5
245.	29	30	33	34	41	Fruit: diameter	Fruit: diamètre	Frucht: Durchmesser	Fruto: diámetro						
QN	(*)	(*)	(*)	(*)		small	petit	klein	pequeño	3					
						medium	moyen	mittel	medio	5					
						large	grand	groß	grande	7					
246.	30	31	34	35	42	Fruit: ratio length/diameter	Fruit: rapport longueur/diamètre	Frucht: Verhältnis Länge/Durchmesser	Fruto: relación longitud/diámetro						
QN	(*)	(*)	(*)	(*)		small	faible	klein	pequeña	3					
						medium	moyen	mittel	media	5					
						large	élevé	groß	grande	7					
247.	31	32	35	36	43	Fruit: position of broadest part	Fruit: position de la partie la plus large	Frucht: Position des breitesten Teils	Fruto: posición de la parte más ancha						
QN	(*)	(*)	(*)	(*)		towards stalk end	vers l'extrémité pédonculaire	zum Stielende hin	hacia el extremo peduncular	1					
						at middle	au milieu	in der Mitte	en el medio	2					
						towards distal end	vers la partie distale	zum distalen Ende hin	hacia el extremo distal	3					
248.	32				44	Fruit: shape in transverse section	Fruit: forme en section transverse	Frucht: Form im Querschnitt	Fruto: forma en sección transversal						
(+)						circular	circulaire	rund	circular	1					
PQ						somewhat angular	légèrement anguleuse	etwas kantig	ligeramente angular	2					
						scalloped	dentelée	muschelförmig	festoneado	3					

	Group/Groupe/Gruppe/Grupo					English	français	deutsch	español	Note/ Nota	Group/Groupe/Gruppe/Grupo				
	1	2	3	4	5						1	2	3	4	5
249. PQ	33	33	36	37	45	Fruit: general shape of proximal part (excluding neck, collar and depression at stalk end)	Fruit: forme générale de la partie proximale (à l'exclusion du col, de la collerette et de la dépression à l'extrémité pédonculaire)	Frucht: allgemeine Form des proximalen Teils (ohne Hals, Kragen und Einsenkung am Stielende)	Fruto: forma general de la parte proximal (excluidos el cuello, el collar y la depresión del extremo peduncular)						
	(+)	(*)		(*)		flattened	aplatie	abgeflacht	aplanada	1					
						slightly rounded	légèrement arrondie	leicht abgerundet	ligeramente redondeada	2					
						strongly rounded	fortement arrondie	stark abgerundet	fuertemente redondeada	3					
						tapered	effilée	spitz	afilada	4					
250. QL	34		37		46	Fruit: presence of neck	Fruit: présence d'un col	Frucht: Vorhandensein eines Halses	Fruto: presencia de cuello						
(+)	(*)		(*)			absent	absent	fehlend	ausente	1					
						present	présent	vorhanden	presente	9					
251. QN	35		38		47	<u>Necked varieties only:</u> Fruit: length of neck	<u>Seulement les variétés dont le fruit présente un col:</u> Fruit: longueur du col	<u>Nur Sorten mit Fruchthals:</u> Frucht: Länge des Halses	<u>Sólo variedades con fruto con cuello:</u> Fruto: longitud del cuello						
						short	court	kurz	corto	3					
						medium	moyen	mittel	medio	5					
						long	long	lang	largo	7					

	Group/Groupe/Gruppe/Grupo					English	français	deutsch	español	Note/Nota	Group/Groupe/Gruppe/Grupo				
	1	2	3	4	5						1	2	3	4	5
252. QN	36					Necked varieties only: Fruit: thickness of neck thin medium thick	Seulement les variétés dont le fruit présente un col: Fruit: épaisseur du col mince moyen épais	Nur Sorten mit Fruchthals: Frucht: Dicke des Halses dünn mittel dick	Sólo variedades con fruto con cuello: Fruto: grosor del cuello delgado medio grueso	3 5 7					
253. (+) QL	37	34	39	38	49	Only varieties without fruit neck: Fruit: presence of depression at stalk end absent present	Seulement les variétés dont le fruit ne présente pas de col: Fruit: présence d'une dépression à l'extrémité pédonculaire absente présente	Nur Sorten ohne Fruchthals: Frucht: Vorhandensein einer Einsenkung am Stielende fehlend vorhanden	Sólo variedades con fruto sin cuello: Fruto: presencia de una depresión en el extremo peduncular ausente presente	1 9					
254. QN	38	35	40	39	50	Only varieties without fruit neck: Fruit: depth of depression at stalk end shallow medium deep	Seulement les variétés dont le fruit ne présente pas de col: Fruit: profondeur de la dépression à l'extrémité pédonculaire peu profonde moyenne profonde	Nur Sorten ohne Fruchthals: Frucht: Tiefe der Einsenkung am Stielende flach mittel tief	Sólo variedades con fruto sin cuello: Fruto: profundidad de la depresión en el extremo peduncular poco profunda media profunda	3 5 7					

	Group/Groupe/Gruppe/Grupo					English	français	deutsch	español	Note/Nota	Group/Groupe/Gruppe/Grupo				
	1	2	3	4	5						1	2	3	4	5
255. (+) QL	39				51	Fruit: presence of constriction at stalk end	Fruit: présence d'un étranglement à l'extrémité pédonculaire	Frucht: Vorhandensein einer Einschnürung am Stielende	Fruto: presencia de un estrechamiento en el extremo peduncular						
						absent	absent	fehlend	ausente	1					
						present	présent	vorhanden	presente	9					
256. QN	40				52	Fruit: expression of constriction at stalk end	Fruit: expression de l'étranglement à l'extrémité pédonculaire	Frucht: Ausprägung der Einschnürung am Stielende	Fruto: expresión del estrechamiento en el extremo peduncular						
						weak	faible	gering	débil	3					
						medium	moyenne	mittel	medio	5					
257. QN	41	36		40	53	Fruit: number of radial grooves at stalk end	Fruit: nombre de cannelures radiales à l'extrémité pédonculaire	Frucht: Anzahl radialer Furchen am Stielende	Fruto: número de acanaladuras radiales en el extremo peduncular						
						absent or few	absentes ou peu nombreuses	fehlend oder sehr gering	ausentes o bajo	1					
						intermediate	moyennement nombreuses	mittel	medio	2					
258. QN	42	37		41	54	Fruit: length of radial grooves at stalk end	Fruit: longueur des cannelures radiales à l'extrémité pédonculaire	Frucht: Länge der radialen Furchen am Stielende	Fruto: longitud de las acanaladuras radiales en el extremo peduncular						
						short	courtes	kurz	cortas	3					
						medium	moyennes	mittel	medianas	5					
						long	longues	lang	largas	7					

	Group/Groupe/Gruppe/Grupo					English	français	deutsch	español	Note/ Nota	Group/Groupe/Gruppe/Grupo				
	1	2	3	4	5						1	2	3	4	5
259.	43				55	Necked varieties only: Fruit: depression at stalk attachment	Seulement les variétés dont le fruit présente un col: Fruit: dépression à l'attache pédonculaire	Nur Sorten mit Fruchthals: Frucht: Einsenkung am Stielansatz	Sólo variedades con fruto con cuello: Fruto: depresión en la inserción peduncular						
QN						absent or shallow	absente ou peu profonde	fehlend oder flach	ausente o poco profunda	1					
						intermediate	intermédiaire	mittel	intermedia	2					
						deep	profonde	tief	profunda	3					
260.	44	38			56	Fruit: presence of collar	Fruit: présence d'une corolle	Frucht: Vorhandensein eines Kragens	Fruto: presencia de un collar						
(+)						absent	absente	fehlend	ausente	1					
QL						present	présente	vorhanden	presente	9					
261.	45					Fruit: height of collar	Fruit: hauteur de la corolle	Frucht: Höhe des Kragens	Fruto: altura del collar						
QN						low	basse	niedrig	bajo	3					
						medium	moyenne	mittel	medio	5					
						high	haute	hoch	alto	7					
262.	46					Fruit: diameter of collar	Fruit: diamètre de la corolle	Frucht: Durchmesser des Kragens	Fruto: diámetro del collar						
QN						small	petit	klein	pequeño	3					
						medium	moyen	mittel	medio	5					
						large	grand	groß	grande	7					

	Group/Groupe/Gruppe/Grupo					English	français	deutsch	español	Note/ Nota	Group/Groupe/Gruppe/Grupo						
	1	2	3	4	5						1	2	3	4	5		
263. QN	47					Fruit: abscission layer between floral disc and fruit absent or weakly developed intermediate strongly developed	Fruit: abscission entre le disque floral et le fruit absente ou faiblement développée intermédiaire fortement développée	Frucht: Trennschicht zwischen Blütenscheibe und Frucht fehlend oder schwach entwickelt mittel stark entwickelt	Fruto: capa de abscisión entre el disco floral y el fruto ausente o débilmente desarrollada intermedia fuertemente desarrollada	1 2 3							
264. QN	48 (+) (*)	39	41	42	58	Fruit: general shape of distal part (excluding nipple, bulging of navel and depression at distal end) flattened slightly rounded strongly rounded	Fruit: forme générale de la partie distale (à l'exclusion du mamelon, de la courbure du fruit secondaire et de la dépression à l'extrémité distale) aplatie légèrement arrondie fortement arrondie	Frucht: allgemeine Form des distalen Teils (ohne Warze, Wölbung der sekundären Frucht und Einsenkung am distalen Ende) abgeflacht leicht abgerundet stark abgerundet	Fruto: forma general de la parte distal (excluidos el mamelón o pezón, el abultamiento del ombligo y la depresión en el extremo distal) aplanada ligeramente redondeada fuertemente redondeada	1 2 3							
265. QL	49 (+) (*)	40		43	59	Fruit: presence of depression at distal end absent present	Fruit: présence d'une dépression à l'extrémité distale absente présente	Frucht: Vorhandensein einer Einsenkung am distalen Ende fehlend vorhanden	Fruto: presencia de una depresión en el extremo distal ausente presente	1 9							

	Group/Groupe/Gruppe/Grupo					English	français	deutsch	español	Note/ Nota	Group/Groupe/Gruppe/Grupo				
	1	2	3	4	5						1	2	3	4	5
266. QN	50			44	60	Fruit: depth of depression at distal end	Fruit: profondeur de la dépression à l'extrémité distale	Frucht: Tiefe der Einsenkung am distalen Ende	Fruto: profundidad de la depresión en el extremo distal						
						shallow	peu profonde	flach	poco profunda	3					
						medium	moyenne	mittel	media	5					
						deep	profonde	tief	profunda	7					
267. QN	51			45		Fruit: diameter of depression at distal end	Fruit: diamètre de la dépression à l'extrémité distale	Frucht: Durchmesser der Einstellung am distalen Ende	Fruto: diámetro de la depresión en el extremo distal						
						small	petit	klein	pequeño	3					
						medium	moyen	mittel	medio	5					
						large	grand	groß	grande	7					
268. (+) QL		42		61		Fruit: presence of nipple	Fruit: présence d'un mamelon	Frucht: Vorhandensein einer Warze	Fruto: presencia de un mamelón o pezón						
		(*)		(*)		absent	absent	fehlend	ausente	1					
						present	présent	vorhanden	presente	9					
269. QN		43		62		Fruit: prominence of nipple	Fruit: proéminence du mamelon	Frucht: Herausragen der Warze	Fruto: prominencia del mamelón o pezón						
						weak	faible	gering	débil	3					
						medium	moyenne	mittel	medio	5					
						strong	forte	stark	fuerte	7					

	Group/Groupe/Gruppe/Grupo					English	français	deutsch	español	Note/ Nota	Group/Groupe/Gruppe/Grupo				
	1	2	3	4	5						1	2	3	4	5
270.	52	41		46	63	Fruit: presence of areola	Fruit: présence d'une aréole	Frucht: Vorhandensein einer Areola	Fruto: presencia de un areola						
QL	(*)	(*)				absent	absente	fehlend	ausente	1					
						incomplete	incomplète	unvollständig	incompleta	2					
						complete	complète	vollständig	completa	3					
271.	53	42		47	64	Fruit: type of areola	Fruit: type d'aréole	Frucht: Typ der Areola	Fruto: tipo de areola						
(+)						smooth	régulière	glatt	lisa	1					
QL						grooved	cannelée	gerieft	acanalada	2					
						ridged	annelée	geringelt	acrestada	3					
272.	54	43		48	65	Fruit: diameter of areola	Fruit: diamètre de l'aréole	Frucht: Durchmesser der Areola	Fruto: diámetro de la areola						
QN						small	petit	klein	pequeño	3					
						medium	moyen	mittel	medio	5					
						large	grand	groß	grande	7					
273.	55	44	44	49	66	Fruit: diameter of styilar scar	Fruit: diamètre de la cicatrice stylaire	Frucht: Durchmesser der Griffelnarbe	Fruto: diámetro de la cicatriz estilar						
QN						small	petit	klein	pequeño	3					
						medium	moyen	mittel	medio	5					
						large	grand	groß	grande	7					

	Group/Groupe/Gruppe/Grupo					English	français	deutsch	español	Note/ Nota	Group/Groupe/Gruppe/Grupo				
	1	2	3	4	5						1	2	3	4	5
274. QN					67	Fruit: protruding stylar point small medium large	Fruit: point stylique protubérant petit moyen grand	Frucht: herausragende Griffelspitze klein mittel groß	Fruto: protuberancia del punto estilar pequeña media grande	3 5 7					
275. PQ	56	45	45		68	Fruit: persistence of style none partial total	Fruit: persistance du style aucune partielle totale	Frucht: Persistenz des Griffels fehlend teilweise gegeben vollständig gegeben	Fruto: persistencia del estilo ninguna parcial total	1 2 3					
276. PQ	57	46				Fruit: presence of navel opening absent occasionally present always present	Fruit: présence d'une ouverture du fruit secondaire absente présente occasionnellement toujours présente	Frucht: Vorhandensein der Öffnung der sekundären Frucht fehlend zuweilen vorhanden immer vorhanden	Fruto: presencia de un apertura del ombligo ausente ocasionalmente presente siempre presente	1 2 3					
277. QN	58	47				Fruit: diameter of navel opening small medium large	Fruit: diamètre de l'ouverture du fruit secondaire petit moyen grand	Frucht: Durchmesser der Öffnung der sekundären Frucht klein mittel groß	Fruto: diámetro de la apertura del ombligo pequeño medio grande	3 5 7					

	Group/Groupe/Gruppe/Grupo					English	français	deutsch	español	Note/ Nota	Group/Groupe/Gruppe/Grupo				
	1	2	3	4	5						1	2	3	4	5
278. QN		48			69	Fruit: bulging of navel	Fruit: courbure du fruit secondaire absent or weak intermediate strong	Frucht: Wölbung der sekundären Frucht absente ou faible intermédiaire forte	Fruto: abultamiento del ombligo ausente o débil medio fuerte		1	2	3	4	5
279. QL	59	49	46		70	Fruit: presence of radial grooves at distal end	Fruit: présence de cannelures radiales à l'extrémité distale absent present	Frucht: Vorhandensein radialer Furchen am distalen Ende absentes présentes	Fruto: presencia de acanaladuras radiales en el extremo distal ausentes presentes		1	9			
280. QN	60		47			Fruit: expression of radial grooves at distal end	Fruit: expression des cannelures radiales à l'extrémité distale weak medium strong	Frucht: Ausprägung der radialen Furchen am distalen Ende faible moyenne forte	Fruto: expresión de las acanaladuras radiales en el extremo distal débil media fuerte		3	5	7		
281. QL		50	48			Fruit: color variegation	Fruit: panachure de la couleur absent present	Frucht: Panachierung der Farbe fehlend vorhanden	Fruto: variegación del color ausente presente		1	9			

	Group/Groupe/Gruppe/Grupo					English	français	deutsch	español	Note/ Nota	Group/Groupe/Gruppe/Grupo				
	1	2	3	4	5						1	2	3	4	5
282.	61	51	49	50	71	Fruit surface: predominant color(s)	Surface du fruit: couleur(s) prédominante(s)	Fruchtoberfläche: Hauptfarbe(n)	Superficie del fruto: color(es) predominante(s)						
PQ	(*)	(*)	(*)			green	vert	grün	verde	1	1	1	1		
						yellow green	vert-jaune	gelbgrün	verde amarillento	2	2	1	2		
						greenish yellow	jaune verdâtre	grünlichgelb	amarillo verdoso			2			
						green and yellow	vert et jaune	grün und gelb	verde y amarillo			3			3
						light yellow	jaune clair	hellgelb	amarillo claro	3	3	3	4		
						medium yellow	jaune moyen	mittelgelb	amarillo medio	4	4	4	5		
						yellow orange	orange-jaune	gelborange	naranja amarillento	5	1	5	6		
						medium orange	orange moyen	mittelorange	naranja medio	6	2		7		
						dark orange	orange foncé	dunkelorange	naranja oscuro	7	3		8		
						green and orange	vert et orange	grün und orange	verde y naranja				9		
						yellow and orange	jaune et orange	gelb und orange	amarillo y naranja				10		
						orange red	rouge orangé	orangerot	rojo anaranjado	8	4		11		
						red	rouge	rot	rojo	9	5				
						yellow and red	jaune et rouge	gelb und rot	amarillo y rojo				12		
						orange and red	orange et rouge	orange und rot	naranja y rojo				13		
						light pink	rose pâle	hellrosa	rosa claro				5		
						medium pink	rose moyen	mittelrosa	rosa medio				6		
						dark pink	rose foncé	dunkelrosa	rosa oscuro				7		
						yellowish orange	orange jaunâtre	gelblich orange	naranja amarillento				8		

	Group/Groupe/Gruppe/Grupo					English	français	deutsch	español	Note/ Nota	Group/Groupe/Gruppe/Grupo				
	1	2	3	4	5						1	2	3	4	5
283. QL					72	Fruit surface: presence of pubescence	Surface du fruit: présence de pilosité	Fruchtoberfläche: Vorhandensein der Behaarung	Superficie del fruto: presencia de pubescencia						
					(*)	absent	absente	fehlend	ausente	1					
						present	présente	vorhanden	presente	9					
284. QN					73	Fruit surface: intensity of pubescence	Surface du fruit: intensité de la pilosité	Fruchtoberfläche: Intensität der Behaarung	Superficie del fruto: intensidad de la pubescencia						
					(*)	weak	faible	gering	débil	3					
						medium	moyenne	mittel	media	5					
						strong	forte	stark	fuerte	7					
285. QN	62 (*)	50 (*)	51			Fruit surface: glossiness	Surface du fruit: brillance	Fruchtoberfläche: Glanz	Superficie del fruto: brillo						
						absent or very weak	absente ou très faible	fehlend oder sehr gering	ausente o muy débil	1					
						weak	faible	gering	débil	3					
						medium	moyenne	mittel	medio	5					
						strong	forte	stark	fuerte	7					
						very strong	très forte	sehr stark	muy fuerte	9					
286. QN	63	52	51	52	74	Fruit surface: roughness	Surface du fruit: rugosité	Fruchtoberfläche: Rauheit	Superficie del fruto: rugosidad						
						smooth	lisse	glatt	lisa	3					
						medium	intermédiaire	mittel	media	5					
						rough	rugueuse	rauh	rugosa	7					

	Group/Groupe/Gruppe/Grupo					English	français	deutsch	español	Note/ Nota	Group/Groupe/Gruppe/Grupo				
	1	2	3	4	5						1	2	3	4	5
287. PQ	64	53	52	53	75	Fruit surface: size of oil glands	Surface du fruit: taille des glandes à huile	Fruchtoberfläche: Größe der Öldrüsen	Superficie del fruto: tamaño de las glándulas de aceite						
						all more or less the same size	toutes plus ou moins de la même taille	alle mehr oder weniger gleich groß	todas más o menos del mismo tamaño	1					
						larger ones interspersed by smaller ones	grandes et petites intercalées	größere vermischt mit kleineren	glándulas grandes intercaladas con otras más pequeñas	2					
288. QN	65	54	53	54	76	Fruit surface: size of larger oil glands	Surface du fruit: taille des glandes à huile les plus grosses	Fruchtoberfläche: Größe der größeren Öldrüsen	Superficie del fruto: tamaño de las glándulas de aceite más grandes						
						small	petites	klein	pequeñas	3					
						medium	moyennes	mittel	medianas	5					
						large	grosses	groß	grandes	7					
289. QN	66	55	54	55	77	Fruit surface: conspicuousness of larger oil glands	Surface du fruit: netteté des glandes à huile les plus grosses	Fruchtoberfläche: Sichtbarkeit der größeren Öldrüsen	Superficie del fruto: visibilidad de las glándulas de aceite más grandes						
						weak	faible	gering	débil	3					
						medium	moyenne	mittel	media	5					
						strong	forte	stark	fuerte	7					

	Group/Groupe/Gruppe/Grupo					English	français	deutsch	español	Note/Nota	Group/Groupe/Gruppe/Grupo				
	1	2	3	4	5						1	2	3	4	5
290.-1 PQ	67	55	56	78	Fruit surface: presence of pitting and pebbling on oil glands	Surface du fruit: présence de dépression et de protubérance sur les glandes à huile		Fruchtoberfläche: Vorhandensein von Grübchen und Körnern an den Öldrüsen	Superficie del fruto: presencia de picado y granulado en las glándulas de aceite						
						pitting and pebbling absent	dépression et protubérance absentes	Grübchen und Körner fehlend	picado y granulado ausentes	1					
						pitting absent, pebbling present	dépression absente, protubérance présente	Grübchen fehlend, Körner vorhanden	picado ausente, granulado presente	2					
						pitting present, pebbling absent	dépression présente, protubérance absente	Grübchen vorhanden, Körner fehlend	picado presente, granulado ausente	3					
						pitting and pebbling present	dépression et protubérance présentes	Grübchen und Körner vorhanden	picado y granulado presentes	4					
290.-2 (+) PQ	56				Fruit surface: presence of pitting and pebbling	Surface du fruit: présence de dépression et de protubérance		Fruchtoberfläche: Vorhandensein von Grübchen und Körnern	Superficie del fruto: presencia de picado y granulado						
						pitting and pebbling absent	dépression et protubérance absentes	Grübchen und Körner fehlend	picado y granulado ausentes	1					
						pitting absent, pebbling present	dépression absente, protubérance présente	Grübchen fehlend, Körner vorhanden	picado ausente, granulado presente	2					
						pitting present, pebbling absent	dépression présente, protubérance absente	Grübchen vorhanden, Körner fehlend	picado presente, granulado ausente	3					
						pitting and pebbling present	dépression et protubérance présentes	Grübchen und Körner vorhanden	picado y granulado presentes	4					

	Group/Groupe/Gruppe/Grupo					English	français	deutsch	español	Note/ Nota	Group/Groupe/Gruppe/Grupo				
	1	2	3	4	5						1	2	3	4	5
291.1	68		56	57	79	<u>Varieties with fruit surface: pitting on oil glands present only:</u> Fruit surface: density of pitting	<u>Variétés avec surface du fruit: présence seulement de dépression sur les glandes à huile:</u> Surface du fruit: densité de la dépression	<u>Sorten mit Frucht-oberfläche: nur Grübchen an den Öldrüsen vorhanden:</u> Fruchtoberfläche: Dichte der Grübchen	<u>Variedades con superficie del fruto: presencia solamente de picado en las glándulas de aceite:</u> Superficie del fruto: densidad del picado						
QN						sparse medium dense	éparse moyenne dense	locker mittel dicht	dispersa media densa	3 5 7					
291.2		57				<u>Varieties with fruit surface: pitting present only:</u> Fruit surface: density of pitting	<u>Variétés avec surface du fruit: présence seulement de dépression:</u> Surface du fruit: densité de la dépression	<u>Sorten mit Frucht-oberfläche: nur Grübchen vorhanden:</u> Fruchtoberfläche: Dichte der Grübchen	<u>Variedades con superficie del fruto: presencia solamente de picado:</u> Superficie del fruto: densidad del picado						
QN						sparse medium dense	éparse moyenne dense	locker mittel dicht	dispersa media densa	3 5 7					
292.			58	80		<u>Varieties with fruit surface: pitting on oil glands present only:</u> Fruit surface: depth of pitting	<u>Variétés avec surface du fruit: présence seulement de dépression sur les glandes à huile:</u> Surface du fruit: profondeur de la dépression	<u>Sorten mit Frucht-oberfläche: nur Grübchen an den Öldrüsen vorhanden:</u> Fruchtoberfläche: Tiefe der Grübchen	<u>Variedades con superficie del fruto: presencia solamente de picado en las glándulas de aceite:</u> Superficie del fruto: profundidad del picado						
QN						shallow medium deep	peu profonde moyenne profonde	flach mittel tief	poco profundo medio profundo	3 5 7					

	Group/Groupe/Gruppe/Grupo					English	français	deutsch	español	Note/Nota	Group/Groupe/Gruppe/Grupo				
	1	2	3	4	5						1	2	3	4	5
293.1	69			59	81	<u>Varieties with fruit surface: pebbling on oil glands present only:</u> Fruit surface: density of pebbling	<u>Variétés avec surface du fruit: présence seulement de protubérance sur les glandes à huile:</u> Surface du fruit: densité de la protubérance sparse medium dense	<u>Sorten mit Frucht-oberfläche: nur Körner an den Öldrüsen vorhanden:</u> Frucht-oberfläche: Dichte der Körner locker mittel dicht	<u>Variedades con superficie del fruto: presencia solamente de granulado en las glándulas de aceite:</u> Superficie del fruto: densidad del granulado dispersa media densa						
QN										3					
293.2		58				<u>Varieties with fruit surface: pebbling present only:</u> Fruit surface: density of pebbling	<u>Variétés avec surface du fruit: présence seulement de protubérance:</u> Surface du fruit: densité de la protubérance sparse medium dense	<u>Sorten mit Frucht-oberfläche: nur Körner vorhanden:</u> Fruchtoberfläche: Dichte der Körner locker mittel dicht	<u>Variedades con superficie del fruto: presencia solamente de granulado:</u> Superficie del fruto: densidad del granulado dispersa media densa						
QN										3					

	Group/Groupe/Gruppe/Grupo					English	français	deutsch	español	Note/ Nota	Group/Groupe/Gruppe/Grupo				
	1	2	3	4	5						1	2	3	4	5
296.	72			62	84	Fruit rind: adherence to flesh	Écorce du fruit: adhérence à la chair	Fruchtschale: Anhaftung am Fleisch	Corteza del fruto: adherencia a la pulpa						
QN	(*)			(*)		weak	faible	gering	débil	3					
						medium	moyenne	mittel	media	5					
						strong	forte	stark	fuerte	7					
297.	73	61			85	Fruit rind: strength	Écorce du fruit: rigidité	Fruchtschale: Festigkeit	Corteza del fruto: vigor						
QN						weak	faible	gering	débil	3					
						medium	moyenne	mittel	medio	5					
						strong	forte	groß	fuerte	7					
298.	74		58			Fruit rind: oiliness	Écorce du fruit: onctuosité	Fruchtschale: Öligkeit	Corteza del fruto: oleosidad						
QN						dry	sèche	trocken	seca	3					
						medium	moyenne	mittel	medianas	5					
						oily	grasse	ölig	oleosa	7					
299.	75				86	Fruit rind: conspicuousness of oil glands on inner surface	Écorce du fruit: netteté des glandes à huile sur la face interne	Fruchtschale: Sichtbarkeit der Öldrüsen an der inneren Oberfläche	Corteza del fruto: glándulas de aceite en la superficie interna						
QN						absent or weakly conspicuous	absentes ou peu nettes	fehlend oder schwach sichtbar	ausentes o poco visibles	1					
						intermediate	moyennes	mittel	intermedias	2					
						strongly conspicuous	très nettes	gut sichtbar	muy visibles	3					

	Group/Groupe/Gruppe/Grupo					English	français	deutsch	español	Note/ Nota	Group/Groupe/Gruppe/Grupo						
	1	2	3	4	5						1	2	3	4	5		
300. PQ	76 62	63	87	Fruit: color of albedo	Fruit: couleur de l'albédo	greenish	verdâtre	grünlich	verdosado	1	1				1		
						white	blanc	weiß	blanco	2	2		1	2			
						light yellow	jaune clair	hellgelb	amarillo claro	3	3			3			
						light orange	orange clair	hellorange	naranja claro	4	4			4			
						light pink	rose pâle	hellrosa	rosa claro			2					
						pink	rose	rosa	rosa	5	5	3	5				
						reddish	rougeâtre	rötlich	rojizo	6	6			6			
301.	77			88	Fruit: density of albedo	Fruit: densité de l'albédo	Fruit: densité de l'albédo	Frucht: Dichte der Albedo	Fruto: densidad del albedo								
QN						loose	faible	locker	laxo	3							
						medium	moyenne	mittel	medio	5							
						dense	dense	dicht	denso	7							
302.	78			89	Fruit: amount of albedo adhering to flesh (strands excluded)	Fruit: quantité d'albédo adhérente à la chair (à l'exclusion des brins)	Fruit: quantité d'albédo adhérente à la chair (à l'exclusion des brins)	Frucht: Umfang der am Fleisch anhaftenden Albedo (ohne Fasern)	Fruto: cantidad de albedo que se adhiere a la pulpa (excluidos los filamentos)								
QN (*)						absent or very small	absent ou très faible	fehlend oder sehr gering	ausente o muy pequeña	1							
						small	faible	gering	pequeña	3							
						medium	moyenne	mittel	media	5							
						large	forte	groß	grande	7							
						very large	très forte	sehr groß	muy grande	9							

	Group/Groupe/Gruppe/Grupo					English	français	deutsch	español	Note/ Nota	Group/Groupe/Gruppe/Grupo				
	1	2	3	4	5						1	2	3	4	5
303. QL	79				90	Fruit: presence of albedo strands	Fruit: présence de brins d'albédo	Frucht: Vorhandensein von Albedofasern	Fruto: presencia de filamentos de albedo						
						absent	absents	fehlend	ausentes	1					
						present	présents	vorhanden	presentes	9					
304. QN	80				91	Fruit: amount of albedo strands	Fruit: quantité de brins d'albédo	Frucht: Anzahl Albedofasern	Fruto: cantidad de filamentos de albedo						
						small	faible	klein	baja	3					
						medium	moyenne	mittel	media	5					
						large	forte	groß	alta	7					
305. QL		63		64		Fruit: differently colored specks in flesh	Fruit: taches de couleurs différentes dans la chair	Frucht: unterschiedlich gefärbte Flecken am Fleisch	Fruto: manchas de distinto color en la pulpa						
						absent	absentes	fehlend	ausentes	1					
						present	présentes	vorhanden	presentes	9					
306. QL		64		65		Fruit: bicolored segments	Fruit: segments bicolores	Frucht: zweifarbig Segmente	Fruto: gajos bicolores						
						absent	absents	fehlend	ausentes	1					
						present	présents	vorhanden	presentes	9					

	Group/Groupe/Gruppe/Grupo					English	français	deutsch	español	Note/ Nota	Group/Groupe/Gruppe/Grupo				
	1	2	3	4	5						1	2	3	4	5
306.1		65		66		<u>Only varieties with Fruit: bicolored segments: present: Fruit: distribution of red coloration</u>	<u>Seulement variétés avec Fruit: segments bicolores: présents: Fruit: distribution de la coloration rouge</u>	<u>Nur Sorten mit Frucht: zweifarbig Segmente: vorhanden: Frucht: Verteilung der Rotfärbung</u>	<u>Sólo variedades con Fruto: gajos bicolores: presentes: Fruto: distribución de la coloración roja</u>						
QL						evenly distributed	uniformément répartis	gleichmäßig verteilt	repartido uniforme	1					
						mainly marginate	principalement au bord	vorwiegend am Rand	principalmente marginal	2					

	Group/Groupe/Gruppe/Grupo					English	français	deutsch	español	Note/ Nota	Group/Groupe/Gruppe/Grupo				
	1	2	3	4	5						1	2	3	4	5
307.	81	66	59	67	92	Fruit: main color of flesh	Fruit: couleur principale de la chair	Frucht: Hauptfarbe des Fleisches	Fruto: color principal de la pulpa						
PQ	(*)	(*)	(*)	(*)		whitish	blanchâtre	weißlich	blanquecino	1					1
						greenish yellow	jaune verdâtre	grünlichgelb	amarillo verdoso					1	
						light green	vert clair	hellgrün	verde claro	2		1	2	2	
						light yellow	jaune clair	hellgelb	amarillo claro	3		2	3		
						medium yellow	jaune moyen	mittelgelb	amarillo medio	4				4	
						light orange	orange clair	hellorange	naranja claro	5	1			5	
						medium orange	orange moyen	mittelorange	naranja medio	6	2			6	
						dark orange	orange foncé	dunkelorange	naranja oscuro	7	3			7	
						orange red	rouge orangé	orangerot	rojo anaranjado			4			
						red	rouge	rot	rojo	8	5			8	
						purple	violet	purpur	púrpura	9				9	
						light pink	rose pâle	hellrosa	rosa claro				3		
						medium pink	rose moyen	mittelrosa	rosa medio			3	4		
						dark pink	rose foncé	dunkelrosa	rosa oscuro				5		
						whitish and pink	blanchâtre et rose	weißlich und rosa	blanquecino y rosa				6		
						yellow and red	jaune et rouge	gelb und rot	amarillo y rojo					10	
						orange	orange	orange	naranja				7		

	Group/Groupe/Gruppe/Grupo					English	français	deutsch	español	Note/Nota	Group/Groupe/Gruppe/Grupo					
	1	2	3	4	5						1	2	3	4	5	
308. QL		67		68	93	Fruit: bitterness of flesh	Fruit: amertume de la chair absent present	Frucht: Bitterkeit des Fleisches fehlend präsent	Fruto: amargor de la pulpa ausente presente							
309. QN	82	68	60	69	94	Fruit: filling of core	Fruit : structure du cœur absent or very sparse sparse medium dense very dense	Frucht: Ausfüllung des inneren Fruchtfleisches fehlend oder sehr locker lâche intermédiaire dense très dense	Fruto: relleno del hueco central ausente o muy laxo laxo medio denso muy denso							
310. QN	83	69	61	70		Fruit: diameter of core	Fruit: diamètre du cœur small medium large	Frucht: Durchmesser des inneren Fruchtfleisches klein mittel groß	Fruto: diámetro del hueco central pequeño medio grande							
311. QN	84	70	62	71	95	Fruit: presence of rudimentary segments	Fruit: présence de segments rudimentaires absent or weak intermediate strong	Frucht: Vorhandensein von unvollständigen Segmenten nulle ou faible intermédiaire forte	Fruto: presencia de gajos rudimentarios fehlend oder schwach mittel stark	Fruto: presencia de gajos rudimentarios nula a débil intermedia fuerte						

	Group/Groupe/Gruppe/Grupo					English	français	deutsch	español	Note/ Nota	Group/Groupe/Gruppe/Grupo				
	1	2	3	4	5						1	2	3	4	5
312. QN	85	71	63	72	96	Fruit: number of well developed segments	Fruit: nombre de segments bien développés	Frucht: Anzahl gut entwickelter Segmente	Fruto: número de gajos bien desarrollados						
						few	peu	gering	bajo	3					
						medium	moyen	mittel	medio	5					
						many	beaucoup	groß	alto	7					
313. QN	86	72				Fruit: coherence of adjacent segment walls	Fruit: adhérence des parois des segments adjacents	Frucht: Zusammenhalt der nebeneinander-liegenden Segmentwände	Fruto: adherencia entre los gajos adyacentes						
						weak	faible	gering	débil	3					
						medium	moyenne	mittel	media	5					
						strong	forte	stark	fuerte	7					
314. QN	87	73	64	73	97	Fruit: strength of segment walls	Fruit: rigidité des parois des segments	Frucht: Festigkeit der Segmentwände	Fruto: firmeza de las paredes de los gajos						
						weak	faible	schwach	débil	3					
						medium	moyenne	mittel	media	5					
						strong	forte	stark	fuerte	7					
315. QN	88	74	65	74		Fruit: length of juice vesicles	Fruit: longueur des vésicules de jus	Frucht: Länge der Saftbläschen	Fruto: longitud de las vesículas de jugo						
						short	courtes	kurz	corta	3					
						medium	moyennes	mittel	media	5					
						long	longues	lang	larga	7					

	Group/Groupe/Gruppe/Grupo					English	français	deutsch	español	Note/ Nota	Group/Groupe/Gruppe/Grupo				
	1	2	3	4	5						1	2	3	4	5
316. QN	89	75	66	75		Fruit: thickness of juice vesicles	Fruit: épaisseur des vésicules de jus	Frucht: Dicke der Saftbläschen	Fruto: grosor de las vesículas de jugo						
						thin	fines	dünn	delgadas	3					
						medium	moyennes	mittel	medianas	5					
						thick	épaisses	dick	gruesas	7					
317. QN	90	76	67	76		Fruit: conspicuousness of juice vesicle walls	Fruit: netteté des parois des vésicules de jus	Frucht: Sichtbarkeit der Saftbläschenwände	Fruto: visibilidad de las paredes de las vesículas de jugo						
						low	faible	gering	baja	3					
						medium	moyenne	mittel	media	5					
						high	forte	stark	alta	7					
318. QN	91	77	68	77		Fruit: coherence of juice vesicles	Fruit: adhérence des vésicules de jus	Frucht: Zusammenhalt der Saftbläschen	Fruto: coherencia de las vesículas de jugo						
						weak	faible	gering	débil	3					
						medium	moyenne	mittel	media	5					
						strong	forte	stark	fuerte	7					
319. PQ	92	78			98	Fruit: presence of navel (viewed internally)	Fruit: présence de fruit secondaire (vue interne)	Frucht: Vorhandensein von sekundären Früchten (von innen gesehen)	Fruto: presencia de ombligo (visto internamente)						
	(*)	(*)				absent or very rare	absent ou très rare	fehlend oder sehr selten	ausente o muy rara	1					
						occasionally present	présent occasionnellement	zuweilen vorhanden	ocasionalmente presente	2					
						always present	toujours présent	immer vorhanden	siempre presente	3					

	Group/Groupe/Gruppe/Grupo					English	français	deutsch	español	Note/ Nota	Group/Groupe/Gruppe/Grupo				
	1	2	3	4	5						1	2	3	4	5
320. QN	93	79				Fruit: size of navel (viewed internally)	Fruit: taille du fruit secondaire (vue interne)	Frucht: Größe der sekundären Frucht (von innen gesehen)	Fruto: tamaño del ombligo visto internamente						
						small	petit	klein	pequeño	3					
						medium	moyen	mittel	medio	5					
						large	grand	groß	grande	7					
321. QN	94	80	69	78	99	Fruit: juiciness	Fruit: succulence	Frucht: Saftigkeit	Fruto: contenido de jugo						
						low	faible	gering	bajo	3					
						medium	moyenne	mittel	medio	5					
						high	élevée	hoch	alto	7					
322. QN (*)	95	81	70	79		Fruit juice: total soluble solids	Jus du fruit: quantité de solides solubles	Fruchtsaft: Gehalt an löslicher Trocken-substanz	Jugo del fruto: cantidad de sólidos solubles						
						low	faible	niedrig	baja	3					
						medium	moyenne	mittel	media	5					
						high	élevée	hoch	alta	7					
323. QN	96	82	71	80		Fruit juice: acidity	Jus du fruit: acidité	Fruchtsaft: Säure	Jugo del fruto: acidez		1				
						very low	très faible	sehr gering	muy baja						
						low	faible	gering	baja	3	3	3	3		
						medium	moyenne	mittel	media	5	5	5	5		
						high	forte	stark	alta	7	7	7	7		

	Group/Groupe/Gruppe/Grupo					English	français	deutsch	español	Note/ Nota	Group/Groupe/Gruppe/Grupo				
	1	2	3	4	5						1	2	3	4	5
324. QN	97	83	72	80		Fruit: strength of fibre	Fruit: rigidité des fibres	Frucht: Festigkeit der Fasern	Fruto: vigor de la fibra						
						weak	faible	schwach	débil	3					
						medium	moyenne	mittel	medio	5					
						strong	forte	stark	fuerte	7					
325. (+) QN	98		73		100	Fruit: number of seeds (controlled manual self-pollination)	Fruit: nombre de pépins (autopolinisation manuelle contrôlée)	Frucht: Anzahl Samen (manuelle kontrollierte Selbstbestäubung)	Fruto: número de semillas (auto-polinización manual controlada)						
						absent or very few	absents ou très peu nombreux	fehlend oder sehr gering	ausentes o muy bajo	1					
						few	peu nombreux	gering	bajo	3					
						medium	moyen	mittel	medio	5					
						many	nombreux	groß	alto	7					
						very many	très nombreux	sehr groß	muy alto	9					
326. (+) QN	99	84	74	82	101	Fruit: number of seeds (open pollination)	Fruit: nombre de pépins (fécondation libre)	Frucht: Anzahl Samen (frei abblühend)	Fruto: número de semillas (polinización libre)						
						absent or very few	absents ou très peu nombreux	fehlend oder sehr gering	ausentes o muy bajo	1					
						few	peu nombreux	gering	bajo	3					
						medium	moyen	mittel	medio	5					
						many	nombreux	groß	alto	7					

	Group/Groupe/Gruppe/Grupo					English	français	deutsch	español	Note/ Nota	Group/Groupe/Gruppe/Grupo					
	1	2	3	4	5						1	2	3	4	5	
327. QL	100 (*)	85 (*)	75 (*)	83 (*)	102 (*)	Seed: polyembryony absent present	Pépin: polyembryonnie absente présente	Samen: Polyembryonie fehlend vorhanden	Semilla: poliembrionía ausente presente		1 9					
328. QN	101	86		84	103	Seed: length short medium long	Pépin: longueur court moyen long	Samen: Länge kurz mittel lang	Semilla: longitud corta media larga		3 5 7					
329. QN	102	87		85	104	Seed: width narrow medium broad	Pépin: largeur étroit moyen large	Samen: Breite schmal mittel breit	Semilla: anchura estrecha media ancha		3 5 7					
330. QL	103	88		86	105	Seed: surface smooth wrinkled	Pépin: surface lisse ridée	Samen: Oberfläche glatt runzlig	Semilla: superficie lisa rugosa		1 2					
331. QN	104	89		87		Varieties with seed: surface wrinkled only: Seed: prominence of wrinkles weak medium strong	Variétés avec pépin: surface ridée seulement: Pépin: proéminence des rides faible moyenne forte	Sorten mit Samen: nur runzlige Oberfläche: Samen: Ausprägung der Runzeln schwach mittel stark	Variedades con semilla: superficie solamente rugosa: Semilla: pro- minencia de la rugosidad débil media fuerte		3 5 7					

	Group/Groupe/Gruppe/Grupo					English	français	deutsch	español	Note/Nota	Group/Groupe/Gruppe/Grupo				
	1	2	3	4	5						1	2	3	4	5
332. PQ	105	90	88	106	Seed: external color	Pépin: couleur externe	Samen: Außenfarbe	Semilla: color externo							
					greenish	verdâtre	grünlich	verdosado	1						
					whitish	blanchâtre	weißlich	blanquecino	2						
					yellowish	jaunâtre	gelblich	amarillento	3						
					pinkish	rosâtre	rosa	rosado	4						
					brownish	brunâtre	bräunlich	amarronado	5						
333. PQ	106	91	89	107	Seed: color of inner seed coat	Pépin: couleur du tégument interne	Samen: Farbe der inneren Samenschale	Semilla: color de la cubierta interna							
					white	blanc	weiß	blanco	1						
					light yellow	jaune clair	hellgelb	amarillo claro	2						
					light brown	marron clair	hellbraun	marrón claro	3						
					medium brown	marron moyen	mittelbraun	marrón medio	4						
					dark brown	marron foncé	dunkelbraun	marrón oscuro	5						
					red	rouge	rot	rojo	6						
					purple	violet	purpur	púrpura	7						

	Group/Groupe/Gruppe/Grupo					English	français	deutsch	español	Note/Nota	Group/Groupe/Gruppe/Grupo				
	1	2	3	4	5						1	2	3	4	5
334. PQ	107 92		90	108	<u>Only varieties with seed: polyembryony present:</u> Seed: color of cotyledons	<u>Seulement variétés avec pépin: polyembryonnie présente:</u> Pépin: couleur des cotylédons	<u>Nur Sorten mit Samen: Polyembryonie vorhanden:</u> Samen: Farbe der Kotyledonen	<u>Sólo variedades con semilla: poliembrionía presente:</u> Semilla: color de los cotiledones							
						white	blanc	weiß	blanco	1					
						cream	crème	cremefarben	crema	2					
						light green	vert clair	hellgrün	verde claro	3					
						dark green	vert foncé	dunkelgrün	verde oscuro	4					
335. QL			76 (*)	91 (*)	109 (*)	Flowering habit	Floraison	Blühverhalten	Tipo de floración						
						flowering once	une seule	einmal blühend	una floración	1					
						flowering more than once	plusieurs	mehr als einmal blühend	más de una floración	2					
336. QN	108 (*)	93 (*)	77 (*)	92 (*)	110	Time of maturity of fruit for consumption	Époque de maturité du fruit pour la consommation	Zeitpunkt der Genußreife	Época de madurez del fruto para su consumo						
						early	précoce	früh	temprana	3					
						medium	moyenne	mittel	media	5					
						late	tardive	spät	tardía	7					
337. QL	109 (*)	94 (*)	78 (*)	93 (*)		Fruit: parthenocarpny	Fruit: parthénocarpie	Frucht: Parthenokarpie	Fruto: partenocarpia						
						absent	absente	fehlend	ausente	1					
						present	présente	vorhanden	presente	9					

	Group/Groupe/Gruppe/Grupo					English	français	deutsch	español	Note/ Nota	Group/Groupe/Gruppe/Grupo				
	1	2	3	4	5						1	2	3	4	5
338. (+) QL	110	95	79	94	111	Plant: self-incompatibility absent present	Plante: auto-incompatibilité absente présente	Pflanze: Selbstinkompatibilität fehlend vorhanden	Planta: autoincompatibilidad ausente presente		1				
339. (+) QN	25					Anther: viable pollen absent or very low low medium high very high	Anthère: pollen viable absent ou très faible faible moyen élevé très élevé	Anthere: keimfähiger Pollen fehlend oder sehr gering gering mittel hoch sehr hoch	Antera: polen viable ausente o muy bajo bajo medio alto muy alto		1				

[End of document/
 Fin du document/
 Ende des Dokuments/
 Fin del documento]