



TG/187/1

INTERNATIONAL UNION
FOR THE PROTECTION
OF NEW VARIETIES OF
PLANTS

UNION INTERNATIONALE
POUR LA PROTECTION
DES OBTECTIONS
VÉGÉTALES

INTERNATIONALER
VERBANDZUMSCHUTZ
VON PFLANZEN -
ZÜCHTUNGEN

UNIÓN INTERNACIONAL
PARA LA PROTECCIÓN
DE LAS OBTECCIONES
VEGETALES

GUIDELINES
FOR THE CONDUCT OF TESTS
FOR DISTINCTNESS, UNIFORMITY AND STABILITY

PRUNUS ROOTSTOCKS

(Prunus L.)

GENEVA
2002

Copies of this document are available on request at the price of 10 Swiss francs each, including surface mail, from the Office of UPOV, 34, chemin des Colombettes, P.O. Box 18, 1211 Geneva 20, Switzerland

This document or parts of it may be reproduced, translated and published without obtaining the specific consent of UPOV, provided that the source is acknowledged.



TG/187/1

ORIGINAL: English

DATE: 2002-04-17

INTERNATIONAL UNION
FOR THE PROTECTION
OF NEW VARIETIES OF
PLANTS

UNION INTERNATIONALE
POUR LA PROTECTION
DES OBTENTIONS
VÉGÉTALES

INTERNATIONALER
VERBANDZUMSCHUTZ
VON PFLANZEN -
ZÜCHTUNGEN

UNIÓN INTERNACIONAL
PARA LA PROTECCIÓN
DE LAS OBTENCIONES
VEGETALES

GUIDELINES
FOR THE CONDUCT OF TESTS
FOR DISTINCTNESS, UNIFORMITY AND STABILITY

PRUNUS ROOTSTOCKS

(Prunus L.)

These Guidelines should be read in conjunction with document TG/1/2, which contains explanatory notes on the general principles on which the Guidelines have been established.

<u>TABLEOFCONTENTS</u>	<u>PAGE</u>
I. SubjectoftheseGuidelines	3
II. MaterialRequired	3
III. ConductofTests	3
IV. MethodsandObservations	4
V. GroupingofVarieties	4
VI. CharacteristicsandSymbols	4
VII. TableofCharacteristics	6
VIII. ExplanationsontheTableofCharacteristics	15
IX. Literature	21
X. TechnicalQuestionnaire	22

I. Subject of these Guidelines

These Test Guidelines apply to all varieties used as rootstocks of all species of *Prunus* L. If characteristics of the flower, the fruit or the seed are necessary to examine the varieties, the Test Guidelines for Almond TG/56/3, Apricot TG/70/3, Cherry TG/35/6, European Plum TG/41/4, Japanese Plum TG/84/3, Mume (Japanese Apricot) TG/160/1 or Peach, Nectarine TG/53/6 should be used for those characteristics, as appropriate.

II. Material Required

1. The competent authorities decide when, where and in what quantity and quality the plant material required for testing the variety is to be delivered. Applicants submitting material from a State other than that in which the testing takes place must make sure that all customs formalities are complied with. As a minimum, the following quantity of plant material is recommended:

- (a) 10 rooted cuttings, for vegetatively propagated varieties, or
- (b) 40 one-year-old seedlings for seed propagated varieties.

2. In the case of seed, the seed should meet the minimum requirements for germination, species and analytical purity, health and moisture content, specified by the competent authority. In cases where the seed is to be stored, the germination capacity should be as high as possible and should be stated by the applicant.

3. The plant material supplied should be visibly healthy, not lacking in vigor or affected by any important pest or disease. It should preferably not be obtained from *in vitro* propagation.

4. The plant material must not have undergone any treatment unless the competent authorities allow or request such treatment. If it has been treated, full details of the treatment must be given.

III. Conduct of Tests

1. The minimum duration of the tests should normally be two independent growing cycles.

2. The tests should normally be conducted at one place. If any important characteristic of the variety cannot be seen at that place, the variety may be tested at an additional place.

3. The tests should be carried out under conditions ensuring satisfactory growth for the expression of the relevant characteristics of the variety and for the conduct of the examination.

4. Additional tests for special purposes may be established.

IV. Methods and Observations

1. Unless otherwise stated, all observations determined by measurement, weighing and counting should be made on 10 plants or parts taken from each of 10 plants for vegetatively propagated varieties, or on 40 plants or parts taken from each of 40 plants for seed propagated varieties.
2. For the assessment of uniformity of:
 - (a) vegetatively propagated varieties, a population standard of 1% and an acceptance probability of at least 95% should be applied. In the case of a sample size of 10 plants, the maximum number of off-types allowed would be 1;
 - (b) self-pollinated seedling varieties, a population standard of 2% and an acceptance probability of at least 95% should be applied. In the case of a sample size of 40 plants, the maximum number of off-types allowed would be 2.
 - (c) cross-pollinated seedling varieties, the assessment should be according to the recommendations in the General Introduction.
3. Unless otherwise stated, all observations on the plant and the leaf should be made during early summer.
4. Unless otherwise stated, all observations on the one-year-old shoot should be made during the dormant season.

V. Grouping of Varieties

1. The collection of varieties to be grown should be divided into groups to facilitate the assessment of distinctness. Characteristics which are suitable for grouping purposes are those which are known from experience not to vary, or to vary only slightly, within a variety. Their various states of expression should be fairly evenly distributed throughout the collection.
2. It is recommended that the competent authorities group the varieties according to the botanical species and use the following characteristics for grouping varieties:
 - (a) Plant: vigor (characteristic 1)
 - (b) Leaf blade: length (characteristic 15)
 - (c) Leaf blade: shape (characteristic 18)
 - (d) Plant: flowers (characteristic 39)

VI. Characteristics and Symbols

1. To assess distinctness, uniformity and stability, the characteristics and their states as given in the Table of Characteristics should be used.

2. Notes (numbers), for the purposes of electronic data processing, are given opposite the states of expression for each characteristic.

3. Legend:

(*) Characteristics that should be used on all varieties in every growing period over which examinations are made and always be included in the variety descriptions, except when the state of expression of a preceding characteristic or regional environmental conditions render this impossible.

(+) See Explanations on the Table of Characteristics in Chapter VIII.

VII. Table of Characteristics/ Tableaux de caractères/ Merkmalstabelle/ Tabla de caracteres

English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
1. Plant:vigor (* (+)	Plante:vigueur	Pflanze: Wuchsstärke	Planta:vigor		
weak	faible	gering	débil	Edabriz, Ferlenain	3
medium	moyenne	mittel	medio	Brokforest, GM 61/1	5
strong	forte	stark	fuerte	Alkavo, F 12/1	7
2. Plant:habit (*	Plante:port	Pflanze:Wuchsform	Planta:porte		
upright	dressé	aufrecht	erecto	Colt	1
spreading	étalé	breitwüchsig	rastrero	Gisela5	3
drooping	retombant	hängend	colgante	<i>Prunusbesseyi</i>	5
3. Plant:branching	Plante:ramification	Pflanze: Verzweigung	Planta:ramificación		
weak	faible	gering	débil	F 12/1, Ferciana	3
medium	moyenne	mittel	media	Pixy	5
strong	forte	stark	fuerte	Gisela 5	7
4. One-year-oldshoot: thickness	Poussed'unan: grosueur	EinjährigerTrieb: Dicke	Ramadeunaño: espesor		
thin	fine	dünn	delgada	Edabriz, Gisela5	3
medium	moyenne	mittel	media	Colt, Pixy	5
thick	grosse	dick	gruesa	Brooks-60, F 12/1	7
5. One-year-oldshoot: lengthofinternode (middlethirdof shoot)	Poussed'unan: longueurde l'entre-nœud(tiers médiandelapousse)	EinjährigerTrieb: Internodienlänge(im mittlerenDritteldes Triebes)	Ramadeunaño: longituddel entrenudo(enel terciomediodela rama)		
short	court	kurz	corto	SL 64	3
medium	moyen	mittel	medio	Colt	5
long	long	lang	largo	F 12/1	7

English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedadesejemplo	Note/ Nota
6. One-year-oldshoot: pubescence(upper third)	Poussed'unan: pilosité(tiers supérieur)	EinjährigerTrieb: Behaarung(oberes Drittel)	Ramadeunaño: pubescencia(enel terciosuperior)		
absent	absente	fehlend	ausente	Pixy	1
present	présente	vorhanden	presente	SL 64	9
7. One-year-oldshoot: numberoflenticels	Poussed'unan: nombredelenticelles	EinjährigerTrieb: AnzahlLentizellen	Ramadeunaño: númerodelenticelas		
few	petit	gering	pequeño	Colt,Fereley	3
medium	moyen	mittel	medio	Gisela4,Pixy	5
many	grand	groß	grande	SL 64	7
8. One-year-oldshoot: anthocyanin colorationofapex	Poussed'unan: pigmentation anthocyaniquedu sommet	EinjährigerTrieb: Anthocyanfärbung derSpitze	Ramadeunaño: pigmentación antociánicadelápice		
absentorveryweak	nulleoutrèsfaible	fehlendodersehr gering	ausenteomuydébil	F 12/1	1
weak	faible	gering	débil	Fereley	3
medium	moyenne	mittel	media	Pixy	5
strong	forte	stark	fuerte	Hamyra	7
verystrong	trèsforte	sehrstark	muyfuerte	Ferciana	9
9. One-year-oldshoot: positionof vegetativebudin relationtoshoot (+)	Poussed'unan: positiondu bourgeonàboispar rapportaurameau	EinjährigerTrieb: Stellungder vegetativenKnospe imVerhältniszum Trieb	Ramadeunaño: posicióndelayema demaderaen relaciónconlarama		
adpressed	appliqué	anliegend	alineada	Hamyra	1
slightlyheldout	légèremetdécollé	leichtabstehend	ligeramente divergente	Gisela 5	2
markedlyheldout	nettementdécollé	deutlichabstehend	fuertemente divergente	F 12/1	3

English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedades ejemplo	Note/ Nota
10. One-year-old shoot: size of vegetative bud	Poussed'unan: tailedubourgeon àbois	Einjähriger Trieb: Größeder vegetativen Knospe	Ramadeunaño: tamaño delayema dem adera		
small	petit	klein	pequeña	SL 64	3
medium	moyen	mittel	media	F 12/1	5
large	gros	groß	grande	Piku 1	7
11. One-year-old shoot: (* shape of apex of (+) vegetative bud	Poussed'unan: formedusommetdu bourgeonàbois	Einjähriger Trieb: Form der Spitzeder vegetativen Knospe	Ramadeunaño: formadelápicedela yemademadera		
acute	pointu	spitz	agudo	Hamyra, Pixy	1
obtuse	obtus	stumpf	obtus	Gisela 5	2
rounded	arrondi	abgerundet	redondeado	F 12/1	3
12. One-year-old shoot: size of vegetative (+) bud support	Poussed'unan: tailedusupportdu bourgeonàbois	Einjähriger Trieb: Größedes Wulstes der vegetativen Knospe	Ramadeunaño: tamaño delsoporte delayemademadera		
small	petit	klein	pequeño	Hamyra	3
medium	moyen	mittel	medio	F 12/1	5
large	grand	groß	grande		7
13. One-year-old shoot: (* branching (at the end of summer)	Poussed'unan: ramification (à la fin del'été)	Einjähriger Trieb: Verzweigung (zum Ende des Sommers)	Ramadeunaño: ramificación (al final del verano)		
weak	faible	gering	débil	Felinem, Mayor	3
medium	moyenne	mittel	media	Adafuel	5
strong	forte	stark	fuerte	GF 677	7

English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedadesejemplo	Note/ Nota
14. Youngshoot: intensityof anthocyanin colorationofyoung leaf(duringrapid growth)	Jeunerameau: intensitédela pigmentation anthocyaniquedela jeunefeuille (pendantla croissancerapide)	JungerTrieb: Intensitätder Anthocyanfärbung desjungenBlattes (währenddes schnellen Wachstums)	Ramajoven: intensidadde la pigmentación antocíanicadela hojajoven(durante elcrecimiento rápido)		
weak	faible	gering	débil	Edabriz,Fereley, Hamyra	3
medium	moyenne	mittel	media	F 12/1	5
strong	forte	stark	fuerte	Colt	7
15. Leafblade:length (*)	Limbe:longueur	Blattspreite:Länge	Limbo:longitud		
veryshort	trèscourt	sehrkurz	muycorto	Myrobalan B	1
short	court	kurz	corto	Edabriz,Weito T6	3
medium	moyen	mittel	medio	Piku 1	5
long	long	lang	largo	F 12/1	7
verylong	trèslong	sehrlang	muylargo	GF 677	9
16. Leafblade:width	Limbe:largeur	Blattspreite:Breite	Limbo:anchura		
verynarrow	trèsétroit	sehrschmal	muyestrecho	GF 677	1
narrow	étroit	schmal	estrecho	Myrobalan B	3
medium	moyen	mittel	medio	Fereley	5
broad	large	breit	ancho	Broksec-60, F 12/1	7
verybroad	trèslarge	sehrbreit	muyancho	Colt	9
17. Leafblade:ratio length/width	Limbe:rapport longueur/largeur	Blattspreite: Verhältnis Länge/Breite	Limbo:relación entrelalongitudyla anchura		
verysmall	trèspetit	sehrklein	muypequeña	GM 61/1	1
small	petit	klein	pequeña	Gisela 5	3
medium	moyen	mittel	media	F 12/1,Pixy	5
large	grand	groß	grande	Piku 3	7
verylarge	trèsgrand	sehrgroß	muygrande	GF 677	9

English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedadesejemplo	Note/ Nota
18. Leafblade:shape (* (+)	Limbe:forme	Blattspreite:Form	Limbo:forma		
narrowelliptic	elliptiqueétroit	schmalelliptisch	elípticoestrecho	GF 677	1
elliptic	élliptique	elliptisch	elíptico	Colt,Fereley,Pixy	2
circular	circulaire	kreisförmig	circular	Adara,SL 64	3
ovate	ovale	eiförmig	oval	Edabriz,Gisela 5	4
obovate	obovale	verkehrteiförmig	oboval		5
19. Leafblade:angleof apex(excludingtip) (+)	Limbe:angleausommet(hors extrémité)	Blattspreite:WinkelanderSpitze(ohne aufgesetzteSpitze)	Limbo:ángulodelápice(excluyendoel extremo)		
acute	aigu	spitz	agudo	GF 677,Pixy	1
right-angled	droit	rechtwinklig	enángulorecto	Edabriz	2
obtuse	obtus	stumpf	obtusos	Colt,Fereley	3
20. Leafblade:lengthof tip (* (+)	Limbe:longueur delapointe	Blattspreite:Länge deraufgesetzten Spitze	Limbo:longitud delápice		
short	courte	kurz	corto	Fereley	3
medium	moyenne	mittel	medio	GM 61/1	5
long	longue	lang	largo	Colt,Ferlenain	7
21. Leafblade:shapeof base (* (+)	Limbe:formedelabase	Blattspreite:Form derBasis	Limbo:formadela base		
acute	pointue	spitz	aguda	Colt	1
obtuse	obtuse	stumpf	obtusa	F 12/1,Ferlenain	2
truncate	tronquée	gerade	truncada	SL 64	3

English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedadesejemplo	Note/ Nota
22. Leafblade:colorof upperside	Limbe:couleurdela facesupérieure	Blattspreite:Farbe derOberseite	Limbo:colordel haz		
lightgreen	vertclair	hellgrün	verdeclaro	Gisela 5, Pixy	1
darkgreen	vertfoncé	dunkelgrün	verdeoscuro	Colt	2
red	rouge	rot	rojo	Citation	3
reddishbrown	brunrougeâtre	rötlichbraun	marrónrojizo	Rubira	4
23. Leafblade: glossinessofupper side	Limbe:brillancede lafacesupérieure	Blattspreite:Glanz derOberseite	Limbo:brillodel haz		
weak	faible	gering	débil	Hamyra	3
medium	moyenne	mittel	medio	Fereley, Gisela 5	5
strong	élevée	stark	fuerte	Colt	7
24. Leafblade: pubescenceoflower sideat apex	Limbe:pilositédel a faceinférieurede l'apex	Blattspreite: Behaarungder Unterseiteander Spitze	Limbo:pubescencia del envésenelápice		
weak	faible	gering	débil	Hamyra	3
medium	moyenne	mittel	media	Pixy	5
strong	forte	stark	fuerte	Weito T6	7
25. Leafblade:incisions (*) ofmargin (+)	Limbe:incisions dubord	Blattspreite: Randeinschnitte	Limbo:incisionesdel borde		
onlycrenate	seulemencrénelées	nur gekerbt	solamentecrenadas	Pixy	1
bothcrenateand serrate	créneléesetendents descie	gekerbtundgesägt	crenadasyaserradas	Adesoto, GF 1869	2
onlyserrate	seulemendententsde scie	nur gesägt	solamenteaserradas	Gisela 5	3
26. Leafblade:depthof incisionsofmargin	Limbe:profondeur desincisionsdubord	Blattspreite:Tiefe derRandeinschnitte	Limbo:profundidad delasincisionesdel borde		
shallow	peu profondes	flach	poco profundas	Edabriz	3
medium	moyennes	mittel	medianas	Piku 3	5
deep	profondes	tief	profundas	Colt	7

English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedadesejemplo	Note/ Nota
27. Petiole:length (*)	Pétiole:longueur	Blattstiel:Länge	Pecíolo:l ongitud		
short	court	kurz	corto	Piku 3	3
medium	moyen	mittel	medio	Pixy	5
long	long	lang	largo	GF 677	7
28. Petiole:presenceof pubescenceofupper side	Pétiole:présencede pilositédelaface supérieure	Blattstiel: Vorhandenseinvon Behaarungde r Oberseite	Pecíolo:presenciade pubescenciadela partesuperior		
absent	absente	fehlend	ausente	F 12/1	1
present	présente	vorhanden	presente	Weito T6	9
29. Petiole:intensityof pubescenceofupper side	Pétiole:intensité delapilositédela facesupérieure	Blattstiel:Stärkeder Behaarungander Oberseite	Pecíolo:intensidad delapubescenciade lapartesuperior		
weak	faible	gering	débil	Colt	3
medium	moyenne	mittel	media	Hamyra	5
strong	forte	stark	fuerte	Weito T6	7
30. Petiole: depthof groove (+)	Pétiole:profondeur du canal	Blattstiel:Tiefeder Rinne	Pecíolo: profundidaddela acanaladura		
shallow	peuprofond	flach	pocoprofunda	F 12/1	3
medium	moyen	mittel	media	Gisela 5	5
deep	profond	tief	profunda	Myrobalan B	7
31. Leaf:ratiolengthof leafblade/lengthof petiole	Feuille:rapport longueurdu limbe/longueurdu pétiole	Blatt:Verhältnis Längeder Blattspreite/Länge desBlattstiels	Hoja:relaciónentre lalongitudellimbo ylalongituddel pecíolo		
small	petit	klein	pequeña	Piku 1	3
medium	moyen	mittel	media	Colt	5
large	grand	groß	grande	Fereley,GF 677	7

English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedadesejemplo	Note/ Nota
32. Leaf:presenceof stipules	Feuille:présence destipules	Blatt: Vorhandenseinvon Nebenblättern	Hoja:presenciade estípulas		
absent	absentes	fehlend	ausentes	Hamyra	1
present	présentes	vorhanden	presentes	F 12/1, Weito T6	9
33. Stipule:length	Stipule:longueur	Nebenblatt:Länge	Estípulas:longitud		
short	court	kurz	cortas	Weito T6	3
medium	moyen	mittel	medianas	Gisela 5, Pixy	5
long	long	lang	largas	F 12/1	7
34. Leaf:presenceof (*) nectaries	Feuille:présence denectaires	Blatt: Vorhandenseinvon Nektarien	Hoja:presenciade nectarios		
absent	absents	fehlend	ausentes	Ferlenain, Hamyra	1
present	présents	vorhanden	presentes	GF 677, Pixy, St. Julien A	9
35. Varietieswith (*) nectariesonly : Leaf:predominant numberofnectaries	Uniquementles variétésànectaires : Feuille:nombre prédominantde nectaires	NurSortenmit Nektarien:Blatt: vorwiegendeAnzahl Nektarien	Sólovariedadescon nectarios:Hoja: número predominantede nectarios		
one	un	eins	uno	Weiroot 158	1
two	deux	zwei	dos	Gisela 5, Pixy	2
morethantwo	plusededeux	mehralszwei	másdedos	Weito T6	3
36. Leaf:positionof nectaries	Feuille:positiondes nectaires	Blatt:Stellungder Nektarien	Hoja:posicióndelos nectarios		
predominantlyon baseofblade	prédominanceàla basedulimbe	vorwiegendander BasisderSpreite	predominantementeenlabasedellimbo	Gisela 5	1
equallydistributedon baseofbladeand petiole	égalementrépartieà labasedulimbeetsur lepétiole	gleichermaßenverteilt anderBasisder Spreiteundam Blattstiel	igualmente distribuido enlabasedellimboy enelpecíolo	Colt	2
predominantlyon petiole	prédominancesurle pétiole	vorwiegendam Blattstiel	predominantementeenelpecíolo	F 12/1	3


English	français	deutsch	español	Example Varieties Exemples Beispielssorten Variedadesejemplo	Note/ Nota
37. Nectary:color (*)	Nectaire:couleur	Nektarie:Farbe	Nectario:color		
green	vert	grün	verde	Pixy	1
yellow	jaune	gelb	amarillo	Weito T6	2
red	rouge	rot	rojo	Weiroot 158	3
violet	violet	violett	violeta	Colt	4
38. Nectary:shape (*)	Nectaire:forme	Nektarie:Form	Nectario:forma		
round	arrondi	rund	redonda	Gisela 5	1
reniform	réniforme	nierenförmig	reniforme	Colt	2
39. Plant:flowers (*)	Plante:fleurs	Pflanze:Blüten	Planta:flore s		
absent	absentes	fehlend	ausentes	Brokforest	1
present	présentes	vorhanden	presentes	Colt	9

VIII. ExplanationsontheTableofCharacteristics


Ad.1:Plant:vigor

The vigor of the plant should be considered as the overall abundance of vegetative growth.


Ad.9:One -year-oldshoot:positionofvegetativebudinrelationtoshoot



1
adpressed




2
slightlyheldout




3
markedlyheldout


Ad.11:One -year-oldshoot:shapeofapexofvegetativebud



1
acute




2
obtuse




3
rounded


Ad.12:One -year-oldshoot:sizeofvegetativebudsupport



3
small




5
medium




7
large


Ad.18: Leafblade:shape




1
narrowelliptic




2
elliptic



3
circular




4
ovate




5
obovate


Ad.19: Leafblade:angleofapex(excludingtip)



1
acute




2
right-angled




3
obtuse


Ad.20: Leafblade:lengthoftip



3
short




5
medium




7
long


Ad.21: Leafblade:shapeofbase



1
acute




2
obtuse



3
truncate

Ad.25: Leafblade:incisionsofmargin



1
only crenate




2
both crenate and serrate




3
only serrate


Ad.30: Petiole: depth of groove



3
shallow



5
medium



7
deep

Explanations on the Reference Varieties

Varietydenomination	Species
Adafuel	<i>Prunusdulcis</i> (Mill.)D.A.Webbx <i>P. persica</i> (L.)Batsch.
Adara	<i>Prunuscerasifera</i> Ehrh.,openpollinated
Adesoto	<i>Prunusdomestica</i> L.ssp.insititia (L.) Schneid
Alkavo	(syn. Altenweddinge Kaukasische Vgelkirsche) <i>Prunus avium</i> (L.)L.
Brokforest	(syn.MxM14) <i>Prunusmahaleb</i> L.x <i>Prunusavium</i> (L.)L.
Brooks-60	(syn.Broksec,MxM60) <i>Prunusmahaleb</i> L.x <i>Prunusavium</i> (L.)L.
Citation	<i>Prunusdomestica</i> L.x <i>P.persica</i> (L.)Batsch.
Colt	<i>Prunusavium</i> (L.)L.x <i>P.pseudocerasus</i> Lindl.
Edabriz	<i>Prunuscerasus</i> L.
F12/1	<i>Prunusavium</i> (L.)L.
Felinem	<i>Prunuspersica</i> (L.)Batsch.x <i>P.dulcis</i> (Mill.)D.A.Webb
Ferciana	(<i>Prunuscerasifera</i> Ehrh.x <i>P. salicina</i> Lindl.)x(<i>P.domestica</i> L. x <i>P.persica</i> (L.)Batsch.)
Fereley	(<i>Prunussalicina</i> Lindl.x <i>P.cerasifera</i> Ehrh.)x <i>P.spinosa</i> L.
Ferlenain	<i>Prunusbesseyi</i> L.H.Baileyx <i>P.cerasifera</i> Ehrh.
GF677	<i>Prunuspersica</i> (L.)Batsch.x <i>P.dulcis</i> (Mill.)D.A.Webb
GF1869	<i>Prunusdomestica</i> (L.)x <i>P.persica</i> (L.)Batsch.
Gisela4	(syn.473/10) <i>Prunusavium</i> (L.)L.x <i>P.fruticosa</i> Pall.
Gisela5	(syn. 148/2) <i>Prunuscerasus</i> L.x <i>P.canescens</i> Bois
GM61/1	<i>Prunusdawyckensis</i> Sealy
Hamyra	<i>Prunus cerasifera</i> Ehrh.
Mayor	<i>Prunuspersica</i> (L.)Batsch.x <i>P.dulcis</i> (Mill.)D.A.Webb
MyrobalanB	<i>Prunuscerasifera</i> Ehrh.
Piku1	(syn. Pi-Ku4,20) <i>Prunusavium</i> (L.)L. x (<i>P.canescens</i> Bois x <i>P.tomentosa</i> Thunb.exMurr.)
Piku3	(syn. Pi-Ku4,83) <i>Prunus.pseudocerasus</i> Lindl. x (<i>P.canescens</i> Bois x <i>P.incisa</i> Thunb.exMurr.)
Pixy	<i>Prunusdomestica</i> L.ssp. insititia (L.)Schneid.
Rubira	<i>Prunuspersica</i> (L.)Batsch.
SL64	(syn. 'SaintLucie64') <i>Prunusmahaleb</i> L.
St.JulienA	<i>Prunusdomestica</i> L.s sp.insititia (L.)Schneid.
Weiroot158	<i>Prunuscerasus</i> L.
WeitoT6	<i>Prunustomentosa</i> Thunb.exMurr.

IX. Literature

Anonymous: The Brooks and Olmo Register of Fruit & Nut Varieties. Alexandria VA, USA, ASHS Press, 3rd edition, 1997, 744p.

De Haas, P.G.: Die Unterlagen - und Baumformen des Kern - und Steinobstes. Stuttgart: Ulmer Verlag, 1976.

Friedrich, G.: Handbuch des Obstbaus. Radebeul: Neumann Verlag, 1993.

Kester, D.E. and C. Grasselly: Almond rootstocks, pp. 265 -293 in: Roy C. Rom and Robert F. Carlson: Rootstocks for Fruit Crops. J. Wiley and Sons, 1987.

Layne, R. E. C.: Peach rootstocks, pp. 185 -216 in: Roy C. Rom and Robert F. Carlson: Rootstocks for Fruit Crops. J. Wiley and Sons, 1987.

Maurer, E.: Die Unterlagen der Obstgehölze. Berlin: Parey Verlag, 1939.

Okie, W. R.: Plum rootstocks, pp. 321 -360 in: Roy C. Rom and Robert F. Carlson: Rootstocks for Fruit Crops. J. Wiley and Sons, 1987.

Perry, R. L.: Cherry rootstocks, pp. 217 -264 in: Roy C. Rom and Robert F. Carlson: Rootstocks for Fruit Crops. J. Wiley and Sons, 1987.

Raynaud, P. C. and J.M. Audergon: Apricot rootstocks, pp. 295 -320 in: Roy C. Rom and Robert F. Carlson: Rootstocks for Fruit Crops. J. Wiley and Sons, 1987.

Salesses, G., Grasselly, C., Renaud, R., Claverie, J.: Les porte greffes des espèces fruitières à noyau du genre *Prunus*. "Amélioration des espèces végétales cultivées. Objectifs et critères de sélection", pp. 768, A. Gallais, H. Bannerot I.N.R.A. Paris, France, 605 -619, 1992.

Wertheim, S.J. : Rootstock Guide. Fruit Research Station Wilhelminadorp, Publication no. 25, 1998.

X. TechnicalQuestionnaire

		ReferenceNumber (nottobefilledinbytheapplicant)
<p>TECHNICALQUESTIONNAIRE tobecompletedinconnectionwithanapplicatio nforPlantBreeders’Rights</p>		
1.1. Genus	<i>Prunus</i> L. PrunusRootstocks	
1.2. Species	<i>P.armeniaca</i> L. 1[] <i>P.avium</i> (L.)L. 2[] <i>P.cerasifera</i> Ehrh. 3[] <i>P.cerasus</i> L. 4[] <i>P.domestica</i> L. 5[] <i>P.dulcis</i> (Mill.)D.A.Webb(<i>P.amygdalus</i> Batsch) 6[] <i>P.mahaleb</i> L. 7[] <i>P.persica</i> (L.)Batsch 8[] <i>P.salicina</i> Lindl. 9[] otherspecies (pleasespecify) 10[] interspecifichybrid (pleasespecify) 11[]	
2.	Applicant(Nameandaddress)	
3.	Proposeddenominationorbreeder’sreference	

4. Information on origin, maintenance and reproduction of the variety

4.1 Origin

(a) Seedling of unknown parentage

(b) Produced by controlled pollination (indicate parent varieties)

-Seed bearing parent

.....

-Pollen parent

.....

(c) Produced by open pollination of (indicate seed bearing parent plant)

.....

(d) Mutation or sport from (indicate original parent variety)

.....

(e) Discovery (indicate where and when)

.....

4.2 *In vitro* propagation

The plant material of the candidate variety has been obtained by *in vitro* propagation

yes

no

4.3 Other type of multiplication (seed, leaf cutting, hardwood cutting, layer)



4.4 Virusstatus

Theplantmaterialisvirus tested yes
no

Iftheanswertothatquestionisyes,pleaseindicateagainstwhichviruses

.....

4.5 Otherinformation

5. Characteristics of the variety to be indicated (the number in brackets refers to the corresponding characteristic in Test Guidelines; please mark the state of expression which best corresponds).

Characteristics	Example Varieties	Note
5.1 Plant:vigor (1)		
weak	Edabriz,Ferlenain	3[]
medium	Brokforest,GM 61/1	5[]
strong	Alkavo,F 12/1	7[]
5.2 Leafblade:length (15)		
veryshort	Myrobalan B	1[]
short	Edabriz,Weito T6	3[]
medium	Piku 1	5[]
long	F 12/1	7[]
verylong	GF 677	9[]
5.3 Leafblade:shape (18)		
narrowelliptic	GF 677	1[]
elliptic	Colt,Fereley,Pixy	2[]
circular	Adara,SL 64	3[]
ovate	Edabriz,Gisela 5	4[]
obovate		5[]
5.4 Plant:flowers (39)		
absent	Brokforest	1[]
present	Colt	9[]

6. Similar varieties and differences from these varieties

Denomination of similar variety	Characteristic in which the similar variety is different ^{o)}	State of expression of similar variety	State of expression of candidate variety
---------------------------------	--	--	--

^{o)} In the case of identical states of expressions of both varieties, please indicate the size of the difference.

7. Additional information which may help to distinguish the variety

7.1 Resistances to pests and diseases

7.2 Utilization as rootstock for

<i>P. armeniaca</i> L.	1 <input type="checkbox"/>
<i>P. avium</i> (L.) L.	2 <input type="checkbox"/>
<i>P. cerasifera</i> Ehrh.	3 <input type="checkbox"/>
<i>P. cerasus</i> L.	4 <input type="checkbox"/>
<i>P. domestica</i> L.	5 <input type="checkbox"/>
<i>P. dulci</i> (Mill.) D.A. Webb (<i>P. amygdalus</i> Batsch)	6 <input type="checkbox"/>
<i>P. mahaleb</i> L.	7 <input type="checkbox"/>
<i>P. persica</i> (L.) Batsch	8 <input type="checkbox"/>
<i>P. salicina</i> Lindl.	9 <input type="checkbox"/>
other species (please specify)	10 <input type="checkbox"/>

7.3 Special conditions for the examination of the variety

7.4 Other information

A representative color photo of the variety should be included in the Technical Questionnaire.

8. Authorizationforrelease

- (a) Does the variety require prior authorization for release under legislation concerningthe protectionoftheenvironment,humanandanimalhealth?

Yes No

- (b) Hassuchauthorizationbeenobtained?

Yes No

Iftheanswertothatquestionisyes,pleaseattachacopyofsuchanauthorization.

[Endofdocument]